

# Sesquiterpene lactones from *Calea pinnatifida*: absolute configuration and structural requirements for antitumor activity

Lha é A. Caldas<sup>1</sup>, Mariana T. Rodrigues<sup>2</sup>, Andrea N. L. Batista<sup>3</sup>, João M. Batista Jr.<sup>4</sup>, João H.G. Lago<sup>5</sup>, Marcelo J. P. Ferreira<sup>6</sup>, Ileana G. S. Rubio<sup>1,2</sup>, Patr éia Sartorelli<sup>1\*</sup>.




Figure S1:  $^1\text{H}$  NMR spectrum of (**1**) (300 MHz; Benzene  $d_6$ )




Figure S2:  $^{13}\text{C}$  NMR spectrum of (**1**) (75 MHz; Benzene  $d_6$ )




Figure S3: HMBC NMR spectrum of (**1**) (300 MHz; Benzene  $d_6$ )




Figure S4: NOESY NMR spectrum of (**1**) (300 MHz; Benzene  $d_6$ )




Figure S5:  $^1\text{H}$  NMR spectrum of (2) (300 MHz;  $\text{CDCl}_3$ )




Figure S6:  $^{13}\text{C}$  NMR spectrum of (2) (75 MHz;  $\text{CDCl}_3$ )




Figure S7: NOESY NMR spectrum of (2) (300 MHz;  $\text{CDCl}_3$ )




Figure S8:  $^1\text{H}$  NMR spectrum of (3) (300 MHz;  $\text{CDCl}_3$ )




Figure S9:  $^{13}\text{C}$  NMR spectrum of (3) (75 MHz;  $\text{CDCl}_3$ )