


In Vitro and In Vivo Screening of Wild Bitter Melon Leaf for Anti-Inflammatory Activity against *Cutibacterium acnes*

Lu-Te Chuang ¹, Ya-Hsin Shih ², Wen-Cheng Huang ², Lie-Chwen Lin ³, Chin Hsu ⁴, Jong-Ho Chyuan ⁵, Tsung-Hsien Tsai ^{6,*} and Po-Jung Tsai ^{2,7,*}

- ¹ Department of Biotechnology and Pharmaceutical Technology, Yuanpei University of Medical Technology, Hsinchu 3, Taiwan; ltchuang@mail.ypu.edu.tw
² Department of Human Development and Family Studies, National Taiwan Normal University, Taipei 1, Taiwan; yayayungchieh@hotmail.com (Y.-H. S.); wencheng7373@gmail.com (W.-C. H.)
³ National Research Institute of Chinese Medicine, Ministry of Health and Welfare, Taipei 1, Taiwan; lclin@nricm.edu.tw
⁴ Department of Exercise Health Science, National Taiwan University of Sport, Taichung 4, Taiwan; jeanhhsu@ntupes.edu.tw
⁵ Hualien District Agricultural Research and Extension Station, Hualien 9, Taiwan; jonghoc@hdares.gov.tw
⁶ Department of Dermatology, Taipei Municipal Wan Fang Hospital and Taipei Medical University, Taipei 1, Taiwan
⁷ Program of Nutrition Science, School of Life Science, National Taiwan Normal University, Taipei 1, Taiwan
* Correspondence: thtsai2@yahoo.com.tw (T.-H. T.); pjtsai@ntnu.edu.tw (P.-J. T.); Tel.: +886-2-7749-1455 (P.-J. T.)

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Figure S1: Gas chromatography–mass spectrometry (GC-MS) fragmentation patterns of β -ionone and dihydroactinidiolide.



Supplementary Figure 1. Gas chromatography–mass spectrometry (GC-MS) fragmentation patterns of β -ionone (a) and dihydroactinidiolide (b). The validity of two compounds was examined using respective authentic standards.

