

*Article*

# The HDAC/HSP90 Inhibitor G570 Attenuated Blue Light-Induced Cell Migration in RPE Cells and Neovascularization in Mice through Decreased VEGF Production

Tai-Ju Hsu <sup>1,†</sup>, Kunal Nepali <sup>1,†</sup>, Chi-Hao Tsai <sup>1,2</sup>, Zuha Imtiyaz <sup>1</sup>, Fan-Li Lin <sup>3</sup>, George Hsiao <sup>3</sup>, Mei-Jung Lai <sup>4,5,\*</sup>, and Yu-Wen Cheng <sup>1,4,\*</sup>

<sup>1</sup> School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei 100301, Taiwan; nick010519@tmu.edu.tw (T.-J.H.); nepali@tmu.edu.tw (K.N.); d01447001@ntu.edu.tw (C.-H.T.); zuhaiimtiyaz@tmu.edu.tw (Z.I.)

<sup>2</sup> Department of Ophthalmology, School of Medicine, the University of North Carolina at Chapel Hill, Chapel Hill 27599, USA.

<sup>3</sup> Department of Pharmacology, School of Medicine, College of Medicine, Taipei Medical University, Taipei 100301, Taiwan; fllin@tmu.edu.tw (F.-L.L.); geohsiao@tmu.edu.tw (G.H.)

<sup>4</sup> Ph.D. Program in Drug Discovery and Development Industry, College of Pharmacy, Taipei Medical University, Taipei 100301, Taiwan

<sup>5</sup> Biomedical Commercialization Center, Taipei Medical University, Taipei 100301, Taiwan

\* Correspondence: [mjl@tmu.edu.tw](mailto:mjl@tmu.edu.tw) (M.-J.L.); [ywcheng@tmu.edu.tw](mailto:ywcheng@tmu.edu.tw) (Y.-W.C.); Tel.: +886-2-27361661 (ext. 7686) (M.-J.L.); +886-2-27361661 (ext. 6123) (Y.-W.C.); Fax: +886-2-27374622 (Y.-W.C.);

† These authors contributed equally to this work.

## Contents:

HPLC purity data.....	SI-2
<sup>1</sup> H NMR Spectra for compounds 1-8.....	SI-3-10



## HPLC Purity Data of compounds 1-8



### HPLC purity determination:


HPLC purity was determined using an Hitachi 2000 series HPLC system using C-18 column (Agilent ZORBAX Eclipse XDB-C18 5 µm. 4.6 mm × 150 mm). Elution conditions: Mobile phase A-Acetonitrile; Mobile phase B-Water containing 0.1% formic acid + 10 mmol NH<sub>4</sub>OAc. The flow-rate was 0.5 mL/min and the injection volume was 5 µl. The system operated at 25 °C. Peaks were detected at 254 nm.

Compound	Retention time (min)	Purity (%)
<b>1</b>	3.104	96.281
<b>2</b>	3.108	99.189
<b>3</b>	3.173	99.431
<b>4</b>	16.54	97.618


<b>5</b>	17.67	96.475
<b>6</b>	18.84	98.886
<b>7</b>	18.75	98.959
<b>8</b>	16.93	97.929


<sup>1</sup>H Spectra for compound 1.<sup>1</sup>H Spectra for compound 2.

<sup>1</sup>H Spectra for compound 3.





<sup>1</sup>H NMR of compound 4.







<sup>1</sup>H NMR of compound 5.







<sup>1</sup>H NMR of compound 6.





<sup>1</sup>H NMR of compound 7.





<sup>1</sup>H NMR of compound 8.

