

Supplementary materials

Anti-allergic activity of 6-deoxy-2-O-methyl-6-(*N*-hexadecanoyl)amino-L-ascorbic acid

Kaori Miura ¹, Hiroaki Matsuno ², Yuji Iwaoka ³, Hideyuki Ito ³ and Akihiro Tai ^{2,4,*}

¹ LAIMU Corporation, 3-6-12 Shinyokohama, Kohoku-ku, Yokohama, Kanagawa 222-0033, Japan; miura@laimu.jp

² Faculty of Life and Environmental Sciences, Prefectural University of Hiroshima, 5562 Nanatsuka-cho, Shobara, Hiroshima 727-0023, Japan; q623021jw@ed.pu-hiroshima.ac.jp (H.M.)

³ Faculty of Health and Welfare Science, Okayama Prefectural University, 111 Kuboki, Soja, Okayama 719-1197, Japan; iwaoka@fhw.oka-pu.ac.jp (Y.I); hito@fhw.oka-pu.ac.jp (H.I.)

⁴ Graduate School of Technology, Industrial and Social Sciences, Tokushima University, 2-1 Minami-josanjima, Tokushima 770-8506, Japan

* Correspondence: atai@tokushima-u.ac.jp

Contents:

Figure S1. ¹H NMR spectrum of 2-Me-6-Palm-AA

Figure S2. ¹³C NMR spectrum of 2-Me-6-Palm-AA

Figure S3. ¹H NMR spectrum of 2-Me-6-*N*-Palm-AA

Figure S4. ¹³C NMR spectrum of 2-Me-6-*N*-Palm-AA

¹H NMR (CD₃OD, 600 MHz)
2-Me-6-Palm-AA









Figure S1. ¹H NMR spectrum of 2-Me-6-Palm-AA

¹³C NMR (CD_3OD , 150 MHz)
2-Me-6-Palm-AA








Figure S4. ^{13}C NMR spectrum of 2-Me-6-*N*-Palm-AA