







## Article

# Hydrolysis of Element (White) Phosphorus under the Action of Heterometallic Cubane-Type Cluster $\{\text{Mo}_3\text{PdS}_4\}$

Airat M. Kuchkaev <sup>1,2</sup>, Nikita Y. Shmelev <sup>1,3,4</sup> , Aidar M. Kuchkaev <sup>1,2</sup>, Aleksandr V. Sukhov <sup>1,2</sup>, Vasily M. Babaev <sup>1</sup> , Khasan R. Khayarov <sup>2</sup> , Artem L. Gushchin <sup>3,\*</sup> , Maxim N. Sokolov <sup>3,4</sup> , Oleg G. Sinyashin <sup>1</sup> and Dmitry G. Yakhvarov <sup>1,2,\*</sup> 

<sup>1</sup> FRC Kazan Scientific Center of RAS, Arbuzov Institute of Organic and Physical Chemistry, Arbuzov Str. 8, 420088 Kazan, Russia; 95ka@bk.ru (A.M.K.); nikulek94@mail.ru (N.Y.S.); kuchkaev95@mail.ru (A.M.K.); alex.suhoff@rambler.ru (A.V.S.); babaev@iopc.ru (V.M.B.); oleg@iopc.ru (O.G.S.)

<sup>2</sup> Department of Organic Chemistry, Kazan Federal University, Kremlyovskaya Str. 18, 420008 Kazan, Russia; khayarov.kh@gmail.com

<sup>3</sup> Nikolaev Institute of Inorganic Chemistry SB RAS Lavrentiev av. 3, 630090 Novosibirsk, Russia; caesar@niic.nsc.ru

<sup>4</sup> Department of Natural Sciences, Novosibirsk State University, Pirogov Str. 2, 630090 Novosibirsk, Russia

\* Correspondence: gushchin@niic.nsc.ru (A.L.G.); yakhvar@iopc.ru (D.G.Y.); Tel.: +7-(843)-2732253 (D.G.Y.)

**Abstract:** Reaction of heterometallic cubane-type cluster complexes— $[\text{Mo}_3\{\text{Pd}(\text{dba})\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$ ,  $[\text{Mo}_3\{\text{Pd}(\text{tu})\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{Cl}$  and  $[\text{Mo}_3\{\text{Pd}(\text{dba})\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$ , where dba—dibenzylideneacetone, dbbpy—4,4'-di-*tert*-butyl-2,2'-bipyridine, tu—thiourea, acac—acetylacetonate, py—pyridine, with white phosphorus ( $\text{P}_4$ ) in the presence of water leads to the formation of phosphorous acid  $\text{H}_3\text{PO}_3$  as the major product. The crucial role of the Pd atom in the cluster core  $\{\text{Mo}_3\text{PdS}_4\}$  has been established in the hydrolytic activation of  $\text{P}_4$  molecule. The main intermediate of the process, the cluster complex  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$  with coordinated  $\text{P}(\text{OH})_3$  molecule and phosphine  $\text{PH}_3$ , have been detected by  $^{31}\text{P}$  NMR spectroscopy in the reaction mixture.

**Keywords:** white phosphorus; heterometallic cubane-type clusters; molybdenum; palladium; phosphorous acid; phosphine



**Citation:** Kuchkaev, A.M.; Shmelev, N.Y.; Kuchkaev, A.M.; Sukhov, A.V.; Babaev, V.M.; Khayarov, K.R.; Gushchin, A.L.; Sokolov, M.N.; Sinyashin, O.G.; Yakhvarov, D.G. Hydrolysis of Element (White) Phosphorus under the Action of Heterometallic Cubane-Type Cluster  $\{\text{Mo}_3\text{PdS}_4\}$ . *Molecules* **2021**, *26*, 538. <https://doi.org/10.3390/molecules26030538>

Academic Editors: Ana Lourenço and Jorge Gominho

Received: 28 December 2020

Accepted: 18 January 2021

Published: 21 January 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Both organic and inorganic phosphorus-containing compounds have become widespread agents for various industrial applications. Traditional methods for the preparation of phosphorus compounds involve oxidation and chlorination of the element (white) phosphorus ( $\text{P}_4$ ) and use the phosphorus chlorides as phosphorylating agents for the synthesis of various organophosphorus substrates. It should be noted that direct activation and transformation of  $\text{P}_4$  is a very harsh and risky process that involves toxic and hazardous reagents and waste, and which negatively impacts on the environment.

The oxidation of  $\text{P}_4$  in the presence of  $\text{H}_2\text{O}$  usually leads to the formation of phosphoric acid  $\text{H}_3\text{PO}_4$ , while phosphorous acid  $\text{H}_3\text{PO}_3$  is more interesting and important phosphorus precursor which can be used as a phosphorylating agent as it contains a functionally capable P-H bond. Moreover, the current methods for the preparation of  $\text{H}_3\text{PO}_3$  leave much to desire and require the use of toxic and hazardous phosphorus trichloride producing a huge amount of corrosive gaseous hydrogen chloride and dangerous phosphorus chloro-derivatives.

We have previously demonstrated that palladium complexes can be efficiently used for the preparation of phosphorous acid  $\text{H}_3\text{PO}_3$  directly from  $\text{P}_4$  by its mild hydrolysis in the coordination sphere of the metal [1,2]. However, the main limitation of these catalytic systems is the formation of catalytically inactive palladium phosphides and palladium black [3]. It is important to note, that ruthenium-mediated  $\text{P}_4$  hydrolysis was thoroughly studied in Florence in the scientific group of M. Peruzzini and P. Stoppioni. It

was shown that the hydrolysis of  $[\text{CpRu}(\text{PPh}_3)_2(\eta^1\text{-P}_4)]$  ( $\text{Cp}$  = cyclopentadienyl) complex leads to the formation of phosphine  $\text{PH}_3$  and phosphorous acid  $\text{H}_3\text{PO}_3$  [4,5]. Moreover, it has been established that the mechanism of this process may involve the formation of binuclear intermediate, when  $\eta^2\text{-P}_4$  unit is doubly coordinated to two  $\{\text{CpRu}(\text{PPh}_3)_2\}$  moieties, which is hydrolyzed with the formation of  $\text{H}_3\text{PO}_3$  and previously unknown 1-hydroxytriphosphane ( $\text{PH}(\text{OH})\text{P}(\text{OH})\text{P}(\text{OH})_2$ ) as the intermediate of the overall process [6]. Later this process was improved and the kinetic of  $\text{P}_4$  hydrolysis was investigated using ruthenium complexes with water-soluble phosphine ligands [7]. There is also a notable example of stabilization of phosphorous acid  $\text{H}_3\text{PO}_3$  in its tautomeric form  $\text{P}(\text{OH})_3$  on the Ru site [8]. Hence, further development of new methods of white phosphorus hydrolysis is of high interest.

Heterometallic cubane-type cluster complexes with  $\{\text{M}_3\text{PdS}_4\}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core that were first described by Hidai's group in Japan [9,10], possess a number of attractive properties including unordinary reactivity and catalytic activity of the Pd site [11–22]. To cite the most recent example,  $\{\text{Mo}_3\text{PdS}_4\}$  cluster complexes react with fullerene  $\text{C}_{60}$  to form hybrid compounds containing a fullerene molecule coordinated to palladium in the cluster core [23]. The  $\{\text{M}_3\text{PdS}_4\}$  clusters catalyze allylation of aromatics [24–26] and nucleophilic addition to triple bonds [10,27]. However, until now there have been no examples of  $\text{P}_4$  molecule coordination and activation in the coordination sphere of the  $\{\text{Mo}_3\text{PdS}_4\}$  cluster core, despite reported ability of such clusters to stabilize the unstable species or less-favoured tautomers, such as  $\text{As}(\text{OH})_3$ ,  $\text{P}(\text{OH})_3$ ,  $\text{PhP}(\text{OH})_2$ ,  $\text{Ph}_2\text{P}(\text{OH})$ ,  $\text{HP}(\text{OH})_2$  through coordination to the palladium site, which indicates high affinity of the latter for pnictogens. [28,29]. From these earlier studies, we have assumed that  $\{\text{Mo}_3\text{PdS}_4\}$  complexes could be involved in the process of  $\text{P}_4$  activation and its transformation.

In this work we investigated the reactivity of heterometallic complexes  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$  (**1**),  $[\text{Mo}_3\{\text{Pd}(\text{tu})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{Cl}$  (**2**) and  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  (**3**) towards element (white) phosphorus.

## 2. Results and Discussion


### 2.1. Synthesis and Characterization

The trinuclear cluster complexes with the  $\{\text{M}_3\text{S}_4\}$  ( $\text{M} = \text{Mo}, \text{W}$ ) core are able to incorporate a range of transition metals in low oxidation state (from 0 to +2), affording heterometallic cubane-type derivatives  $\{\text{M}_3\text{M}'\text{S}_4\}$ , where  $\text{M}' = \text{Cu}, \text{Ni}, \text{Pd}, \text{Pt}$ , etc. [11,30–35].

The common approach for synthesis of the  $\{\text{M}_3\text{M}'\text{S}_4\}$  clusters involves the reaction of a low-valent metal precursor with a  $\{\text{M}_3\text{S}_4\}$  trinuclear complex in a desired coordination environment [12]. The same synthetic approach has been applied in current work, where  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  was used as the palladium source. The synthetic routes to new cluster complexes **1** and **3** used in this work are depicted in Scheme 1. The cluster complex **2** has been obtained according to a previously published procedure [36].

The reaction of  $[\text{Mo}_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$  with  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  (2:1 molar ratio) in  $\text{CH}_2\text{Cl}_2$  gives complex **1** in 54% yield. The complex **3** was prepared in a similar way in 47% yield from  $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  as starting compound. The formation and identity of **1** and **3** were confirmed by microanalysis and spectroscopic data.

The IR spectrum of **1** demonstrates the characteristic composite vibration bands,  $\nu(\text{C}-\text{C})$  and  $\nu(\text{C}-\text{N})$  at  $1610\text{ cm}^{-1}$ ;  $\nu(\text{C}-\text{C})$ ,  $\delta(\text{C}-\text{H})$ ,  $\delta(\text{C}-\text{N})$  and  $\delta(\text{Mo}-\text{N})$  in the  $1480\text{--}1410\text{ cm}^{-1}$  range;  $\gamma(\text{C}-\text{H})$  and  $\delta(\text{C}-\text{H})$  in the  $870\text{--}830\text{ cm}^{-1}$  and  $1380\text{--}1310\text{ cm}^{-1}$  regions, and ring breathing bands from the coordinated dbbpy ligands in the  $1028\text{--}906$  and  $766\text{--}423\text{ cm}^{-1}$  regions. The spectral data are in agreement with those reported for complex **2** [36]. The intensive band ( $\nu(\text{C}=\text{O})$ ) at  $1625\text{ cm}^{-1}$  testifies the coordination of dba ligand to the palladium atom, while bands at  $839$  and  $555\text{ cm}^{-1}$  belong to the hexafluorophosphate anion.



**Scheme 1.** Synthesis of cubane-type cluster complexes **1** and **3**.

In the case of **3**, the IR spectrum is complicated by a significant overlap between bands from acac and py ligands. The band at  $1603\text{ cm}^{-1}$  ( $\nu(\text{C}=\text{C} + \text{C}-\text{N})$ ) relates to the bonded pyridine, and the two characteristic bands ( $\nu(\text{C}=\text{C} + \text{C}=\text{O})$ ) at  $1578$  and  $1523\text{ cm}^{-1}$  are associated with the acetylacetonate ligand. The two intensive bands at  $838$  and  $556\text{ cm}^{-1}$  are explained by the presence of  $\text{PF}_6^-$  group. The highly intensive band at  $1627\text{ cm}^{-1}$  reveals the presence of coordinated dba [37].

The  $^1\text{H}$  NMR spectra of both **1** and **3** in  $\text{CDCl}_3$  demonstrate a complicated pattern due to the presence of overlapping signals in the 7–9 ppm area, that originate from the protons of various aromatic rings present in dbbpy, py, and dba ligands. There are also signals related to alkene fragments in dba ( $\delta \sim 7$  ppm). The characteristic signals generated by protons of the tert-butyl group ( $\delta 1.40\text{--}1.45$  ppm) in dbbpy ligand are detected in the case of **1**. The spectrum of **3** contains peaks associated with  $\text{CH}_3$ - ( $\delta 1.85$  ppm) and  $\text{CH}$ -groups ( $\delta 5.36$  ppm) in acac.


The ESI-MS (+) of **1** contains the peaks with  $m/z$  ratio = 1669 ( $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ) (pseudomolecular peak), 1436 ( $[\text{Mo}_3\text{PdS}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ), 1476 ( $[\text{Mo}_3\{\text{Pd}(\text{CH}_3\text{CN})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ), 1327 ( $[\text{Mo}_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ) which result from  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$  fragmentation under experimental conditions. The signals with  $m/z$  ratio of 1057.8 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})_3]^+$ ), 978.6 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})_2]^+$ ), 899.6 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})]^+$ ), 951.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_3]^+$ ), 870.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_2]^+$ ), 793.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})]^+$ ), 714.7 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3]^+$ ) are detected in the spectrum of **3**. The de-coordination of the pyridine ligands is expectable under ionization conditions and was also observed for the trinuclear  $\{\text{Mo}_3\text{S}_4\}$  precursor [38]. All the peaks have been assigned both from  $m/z$  and characteristic isotope patterns.

## 2.2. Interaction with $\text{P}_4$

The reactivity of heterometallic cluster complexes  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$  (**1**),  $[\text{Mo}_3\{\text{Pd}(\text{tu})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{Cl}$  (**2**) and  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  (**3**) towards  $\text{P}_4$  was investigated both in the absence and in the presence of water. According to  $^{31}\text{P}$  NMR spectra, addition of the equimolar amount of  $\text{P}_4$  to the solutions of these complexes in DMF, THF and  $\text{CH}_2\text{Cl}_2$  does not lead to transformation of  $\text{P}_4$  molecule, and new signals from phosphorus-containing species were detected.

However, addition of water to the DMF solutions containing complexes **1** or **2** and  $\text{P}_4$  has led to the appearance of the signals associated with inorganic oxo-acids  $\text{H}_3\text{PO}_3$  ( $\delta 2.1$  ppm) and  $\text{H}_3\text{PO}_4$  ( $\delta 0.9$  ppm) with integral ratio of 3.4:1.0 (for **1**) and 7.5:1.0 (for **2**). Additionally, new signals around 110 ppm were detected. These signals correspond to the formation of  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$  (Figure 1), in which the Pd atoms bears the tautomeric form of phosphorous acid  $\text{H}_3\text{PO}_3$  ( $\text{P}(\text{OH})_3$ ) formed by the hydrolysis

of white phosphorus. These results nicely fit with the previously published data [28,29], where the signals around 115 ppm in  $^{31}\text{P}$  NMR spectra were attributed to the complexes  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4(\text{H}_2\text{O})_{9-x}\text{Cl}_x]^{(4-x)+}$  obtained by the reaction of  $[\text{Mo}_3\{\text{PdCl}\}\text{S}_4(\text{H}_2\text{O})_9]^{3+}$  with  $\text{PCl}_3$  or  $\text{H}_3\text{PO}_3$  in 4M HCl. The total conversion of white phosphorus (by  $^{31}\text{P}$  NMR spectroscopy) in these reactions was 96.7% for complex **1** and 68.0% for complex **2**.



**Figure 1.** Structure of  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ .

In order to boost the activity of clusters **1** and **2**, we attempted modification of their ligand surrounding by the substitution of the chloride-ions from the first and the second coordination spheres with weakly coordinated ions, and in this way to increase the electrophilic properties of Pd. Indeed, addition of  $\text{TiNO}_3$  as a halide scavenger increased the reactivity of the cluster towards  $\text{P}_4$ . As a result, increased intensity of the signals related to  $\text{H}_3\text{PO}_3$  and  $\text{H}_3\text{PO}_4$ , and the decreased intensity of the  $\text{P}_4$  signal were observed in  $^{31}\text{P}$  NMR spectra. In the case of complex **1**, full conversion of  $\text{P}_4$  was accomplished with a 72.0% yield of  $\text{H}_3\text{PO}_3$ . Complex **2** gave 74.4% conversion of  $\text{P}_4$  and 53.1% yield of  $\text{H}_3\text{PO}_3$ .

Addition of  $\text{H}_2\text{O}$  to the reaction mixture containing  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  (**3**) in DMF and  $\text{P}_4$  also allowed for the appearance of the signals associated with  $\text{H}_3\text{PO}_3$  and  $\text{H}_3\text{PO}_4$ . The conversion of  $\text{P}_4$  (34.4%) was, however, substantially lower than with **1** and **2** and no signal attributed to  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4(\text{acac})_3(\text{py})_3]^+$  was observed in  $^{31}\text{P}$  NMR spectrum. This can be explained by assuming  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4(\text{acac})_3(\text{py})_3]^+$  being less stable than  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ , and quickly releases  $\text{P}(\text{OH})_3$ , which tautomerizes into the final product—phosphorous acid. The observed yield of  $\text{H}_3\text{PO}_3$  was only 20.1%.

The solvent influence on the reactivity of  $\text{P}_4$  and its hydrolysis was also investigated, using the complex **1** as the benchmark. Addition of an excess of  $\text{H}_2\text{O}$  to the reaction mixture containing **1** and  $\text{P}_4$  in  $\text{CH}_2\text{Cl}_2$  allowed for detection of the signals associated with  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$  ( $\delta$  110 ppm),  $\text{H}_3\text{PO}_3$  ( $\delta$  4.7 ppm) and  $\text{H}_3\text{PO}_4$  ( $\delta$  1.1 ppm). Adding  $\text{TiNO}_3$  increased the signal intensities, and the observed molar ratio of  $\text{H}_3\text{PO}_3$ : $\text{H}_3\text{PO}_4$  was 3:2. The signal at  $\delta$ -243.6 ppm related to the formation of  $\text{PH}_3$  was also observed, as minor peak. The conversion of  $\text{P}_4$  was 43.1% with only 8.6% yield of  $\text{H}_3\text{PO}_3$ .

In case of THF as the solvent, the addition of  $\text{H}_2\text{O}$  to the reaction mixture containing **1** and  $\text{P}_4$  yielded the signal associated with  $[\text{Mo}_3\{\text{PdP}(\text{OH})_3\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$  with  $\delta$  +113.8 ppm in  $^{31}\text{P}$  NMR spectrum. It is worth noting that there were no signals of any phosphorus-containing acids in  $^{31}\text{P}$  NMR spectra in this case. However, the activation of the complex with  $\text{TiNO}_3$  caused both the signal growth and the appearance of new signals related to  $\text{H}_3\text{PO}_3$  ( $\delta$  3.3 ppm),  $\text{H}_3\text{PO}_4$  ( $\delta$  1.1 ppm), and  $\text{PH}_3$  ( $\delta$ -244.4 ppm). The integrated intensity ratio  $\text{H}_3\text{PO}_3$ : $\text{H}_3\text{PO}_4$  was 8:3, and the observed conversion of  $\text{P}_4$  was 85.5% with 24.4% yield of  $\text{H}_3\text{PO}_3$ .

The summary of the results obtained in the reaction of  $\{\text{Mo}_3\text{PdS}_4\}$  complexes with  $\text{P}_4$  is presented in Table 1.

**Table 1.** Interaction of {Mo<sub>3</sub>PdS<sub>4</sub>} complexes 1–3 with P<sub>4</sub>.

Complex.	Solvent	Complex + P <sub>4</sub> + H <sub>2</sub> O		Complex + P <sub>4</sub> + H <sub>2</sub> O + TiNO <sub>3</sub>	
		Conversion of P <sub>4</sub> (%)	Yield of H <sub>3</sub> PO <sub>3</sub> (%) / H <sub>3</sub> PO <sub>4</sub> (%)	Conversion of P <sub>4</sub> (%)	Yield of H <sub>3</sub> PO <sub>3</sub> (%) / H <sub>3</sub> PO <sub>4</sub> (%)
1	DMF	96.7	61.5/20.5	100.0	72.0/27.9
	CH <sub>2</sub> Cl <sub>2</sub>	33.6	6.1/1.2	43.1	8.6/4.3
	THF	14	0/0	85.5	24.4/10.4
2	DMF	68.8	37.1/3.7	74.4	53.1/16.2
3	DMF	34.4	20.1/11.6	-	-

It should be noted, that the presence of the Pd site in the cluster moiety to realise the hydrolysis of P<sub>4</sub> molecule is mandatory, as this reaction does not proceed with Pd-free trinuclear cluster complexes [Mo<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]Cl and [Mo<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]PF<sub>6</sub>. This fact confirms that the transformation and followed hydrolysis of P<sub>4</sub> requires the presence of Pd center. Moreover, the use of cluster core {Mo<sub>3</sub>PdS<sub>4</sub>} for hydrolysis of white phosphorus tetrahedron and its transformation into phosphorous acid (H<sub>3</sub>PO<sub>3</sub>) allows to avoid the formation of the insoluble and inactive Pd-black that is very important for the further use of these {Mo<sub>3</sub>PdS<sub>4</sub>} clusters as catalysts for the hydrolysis of white phosphorus.

### 3. Conclusions

Based on the experimental data, we can conclude that heterometallic cubane-type clusters [Mo<sub>3</sub>{Pd(dba)}S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]PF<sub>6</sub> (1), [Mo<sub>3</sub>{Pd(tu)}S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]Cl (2) and [Mo<sub>3</sub>{Pd(dba)}S<sub>4</sub>(acac)<sub>3</sub>(py)<sub>3</sub>]PF<sub>6</sub> (3) efficiently promote the hydrolysis of P<sub>4</sub> molecule leading to the formation of H<sub>3</sub>PO<sub>3</sub> as the major product. The complexes 1 and 2 bearing dbbpy ligand demonstrate higher activity in comparison with complex 3 containing acac ligand. Moreover, removal chloride anions from the coordination sphere of the cluster core with TiNO<sub>3</sub> increases both the activity of the cluster complexes in P<sub>4</sub> activation process and the yield of H<sub>3</sub>PO<sub>3</sub>. The use of the cluster with embedded Pd atom allows to avoid the Pd black formation which occurs when non-cluster Pd complexes are used for white phosphorus hydrolysis process. Thus, this work opens up prospects for studying the potential of heterometallic cubane-type clusters as catalysts for the selective conversion of white phosphorus to phosphorous acid. Further studies are in progress.

### 4. Experimental Section

**CAUTION:** White phosphorus and phosphine mentioned in this communication are hazardous compounds. White phosphorus needs to be stored under water in a well-ventilated dark place. White phosphorus is highly toxic and burns spontaneously when exposed to air. In an emergency, white phosphorus can be treated with aqueous copper(II) sulfate solution or sand. On contact with skin, white phosphorus gives highly painful, badly healing burns. In case of skin burns, washing with diluted aqueous solutions of KMnO<sub>4</sub> or CuSO<sub>4</sub> is advised. The continuous inhaling of white phosphorus vapors results in disease of the bone tissue, loss of teeth, and necrosis of parts of the jaw. An aqueous copper(II) sulfate solution (2%) can be used as an immediate antidote for poisoning. All reactions and handling of phosphine and white phosphorus must be carried out under an inert atmosphere in a well-ventilated hood.

All experiments related to the synthesis of the complexes, preparation of the solutions, solvents, and the manipulations with white phosphorus and all chemical reagents were performed under nitrogen atmosphere using standard Schlenk-line techniques.

Trinuclear precursors [Mo<sub>3</sub>S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]X (X = Cl<sup>−</sup>, PF<sub>6</sub><sup>−</sup>; dbbpy—4,4′-di-*tert*-Bu-2,2′-bipyridine) [39] and [Mo<sub>3</sub>S<sub>4</sub>(acac)<sub>3</sub>(py)<sub>3</sub>]PF<sub>6</sub> (acac—acetylacetonate, py—pyridine) [40] used to synthesize complexes 1, 2 and 3 as well as complex [Mo<sub>3</sub>{Pd(tu)}S<sub>4</sub>Cl<sub>3</sub>(dbbpy)<sub>3</sub>]Cl (2) (tu—thiourea) [36], were prepared according to the published procedures. Commercially



available reagents  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  (dba—dibenzylideneacetone) (Sigma-Aldrich, Steinheim, Germany), thallium(I) nitrate (99.5%, ACROS Organics, Geel, Belgium) were used as purchased. White phosphorus was stored under a protective nitrogen atmosphere in a flask filled with water in a dark place and was washed sequentially in ethanol, acetone, and diethyl ether prior to use. 0.05 M THF solution was prepared by dissolving  $\text{P}_4$  in a required amount of solvent. Organic solvents were distilled before used.

Elemental C, H, N analyses were performed with a EuroEA3000 Eurovector analyzer (EurovectorSpA, Milano, Italy). IR spectra of samples in KBr pellets were recorded in the  $4000\text{--}400\text{ cm}^{-1}$  range with a Perkin-Elmer System 2000 FTIR spectrometer (PerkinElmer, Waltham, Massachusetts, USA).  $^1\text{H}$  and  $^{31}\text{P}\{^1\text{H}\}$  NMR spectra were registered at room temperature on a Bruker Avance III 400 MHz spectrometer (Bruker, New York, New York, USA) at frequencies of 400.0 ( $^1\text{H}$ ) and 161.9 ( $^{31}\text{P}$ ) MHz. UV-vis spectra were recorded with a Specord M40 (Carl Zeiss, Jena, Germany), Helios  $\gamma$  spectrophotometer (ThermoFisher Scientific, Waltham, Massachusetts, USA) in the 200–900 nm range in the  $\text{CH}_3\text{CN}$  solution. A mass spectrometer (Agilent, 6130 Quadrupole MS, 1260 infinity LC, Santa Clara, California, USA) was utilized for the ESI measurements of **1**. The drying gas was nitrogen at a  $300\text{ L h}^{-1}$  flow rate. The sample solution (approx.  $5 \times 10^{-5}\text{ M}$ ) in acetonitrile was infused through a syringe pump directly into the interface at a flow rate of  $0.4\text{ mL min}^{-1}$ . The temperature of the source block was set to  $120\text{ }^\circ\text{C}$  and the interface to  $150\text{ }^\circ\text{C}$ . A capillary voltage of 2.0 kV was used in the positive scan mode, and low values of the cone voltage ( $U_c = 5\text{--}10\text{ V}$ ) were used to control the extent of fragmentation. The observed isotopic pattern of each compound perfectly matched the theoretical isotope pattern calculated from their elemental composition by using the MassLynx 4.1 program (Waters Corporation, Milford, Massachusetts, USA). ESI measurements of **3** were performed using an AmazonX (Bruker Daltonics, Bremen, Germany) ion trap mass spectrometer in positive mode in the mass range of 200–3000 Da. The sample solution (approx.  $5 \times 10^{-5}\text{ M}$ ) in DMF was infused through a syringe pump directly into the interface at a flow rate of  $0.2\text{ mL min}^{-1}$ . The ESI-MS conditions were as follows: capillary voltage, 2.5 kV; nitrogen drying gas,  $10\text{ L/min}$ ,  $250\text{ }^\circ\text{C}$ . Data processing was performed by DataAnalysis software (Bruker Daltonik GmbH, Version 4.0 SP4, Bremen, Germany).

Synthesis of  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$  (**1**). A mixture of  $[\text{Mo}_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{PF}_6$  (0.2 g, 0.13 mmol) and  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  (0.070 g, 0.068 mmol) was stirred in  $\text{CH}_2\text{Cl}_2$  (20 mL) for 24 h. An excess of *n*-hexane was layered onto the resulting brown solution to give a brown product of **1** that was washed by *n*-hexane and diethyl ether. Yield: 0.133 g (54%).

IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 3395 w, 3158 m, 3123 m, 2960 s, 2903 s, 2869 s, 1626 s, 1618 s, 1549 m, 1477 m, 1464 m, 1415 s, 1311 w, 1295 w, 1254 m, 1201 w, 1158 w, 1129 w, 1020 m, 985 s, 975 w, 903 m, 885 w, 859 s, 830 m, 741 w, 720 w, 604 w, 555 s, 485 w, 427 w.

NMR  $^1\text{H}$  (400 MHz,  $\text{CDCl}_3$ , 293 K):  $\delta = 9.93$  (d, 3H), 9.06 (3H), 8.49 (d, 3H), 8.31 (d, 3H), 7.80 (2H, dba), 7.28–7.65 (10H, dba), 7.31–7.39 (m, 6H, dbbpy), 7.01 (2H, dba), 1.45 (s, 27H,  $^t\text{Bu}$ ), 1.40 (s, 27H,  $^t\text{Bu}$ ) ppm.

ESI-MSI (+,  $\text{CH}_3\text{CN}$ ):  $m/z = 1671$  ( $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ), 1437 ( $[\text{Mo}_3\text{PdS}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ), 1331 ( $[\text{Mo}_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]^+$ ), 1313 ( $[\text{Mo}_3\text{S}_4\text{Cl}_2(\text{OH})(\text{dbbpy})_3]^+$ ), 1383 ( $[\text{Mo}_3\text{PdS}_4(\text{OH})_3(\text{dbbpy})_3]^+$ ), 1460 ( $[\text{Mo}_3\{\text{Pd}(\text{CH}_3\text{CN})\}\text{S}_4\text{Cl}_2(\text{OH})(\text{dbbpy})_3]^+$ ), 1277 ( $[\text{Mo}_3\text{S}_4(\text{OH})_3(\text{dbbpy})_3]^+$ ).

Anal. Calc. for  $\text{C}_{71}\text{H}_{86}\text{Cl}_3\text{F}_6\text{Mo}_3\text{N}_6\text{OPdS}_4$ : C, 47.0; H, 4.8; N, 4.6%. Found: C, 46.7; H, 4.4; N, 5.0%.

Synthesis of  $[\text{Mo}_3\{\text{Pd}(\text{tu})\}\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{Cl}$  (**2**). A mixture of  $[\text{Mo}_3\text{S}_4\text{Cl}_3(\text{dbbpy})_3]\text{Cl}$  (0.1 g, 0.74 mmol),  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  (0.038 g, 0.37 mmol), and thiourea (0.056 g, 0.74 mmol) in 20 mL of  $\text{CH}_2\text{Cl}_2$  was stirred for 24 h. An excess of hexane was layered onto the resulting brown solution to give a greenish-brown product of **2**. Yield: 0.080 g (71%). The obtained analytical data nicely fit with previously published results [36].

Synthesis of  $[\text{Mo}_3\{\text{Pd}(\text{dba})\}\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  (**3**). A mixture of  $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_3]\text{PF}_6$  (0.2 g, 0.17 mmol) and  $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$  (0.089 g, 0.086 mmol) were stirred in  $\text{CH}_2\text{Cl}_2$

(20 mL) for 24 h. An excess of *n*-hexane was layered onto the resulting brown solution to give a green-brown product of **3** that was washed by *n*-hexane and diethyl ether. Yield: 0.121 g (47%).

IR ( $\nu$ ,  $\text{cm}^{-1}$ ): 2918 m, 1627s, 1611m, 1603 m, 1597s, 1578 m, 1568 m, 1523 s, 1485 m, 1444 s, 1416 m, 1366 s, 1280 s, 1177 w, 1151 w, 1066 s, 1021 s, 1013 s, 975 w, 876 s, 838 s, 782 w, 755 s, 698 s, 672 m, 622 m, 551 s, 540 m, 490s, 425 s.

NMR  $^1\text{H}$  (400 MHz,  $\text{CDCl}_3$ , 293 K):  $\delta$  = 9.58, 9.47 (6H,  $\alpha$ -py), 8.15, 8.13, 8.10 (3H,  $\gamma$ -py), 7.80–7.30 (2H + 10H, dba; 6H,  $\beta$ -py), 7.03 (2H, dba) 5.36 ( $\gamma$ -CH, acac), 1.85 ( $\text{CH}_3$ , acac) ppm.

ESI-MSI (+, DMF):  $m/z$  = 1057.8 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})_3]^+$ ), 978.6 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})_2]^+$ ), 899.6 ( $[\text{Mo}_3\text{PdS}_4(\text{acac})_3(\text{py})]^+$ ), 951.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_3]^+$ ), 870.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})_2]^+$ ), 793.8 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3(\text{py})]^+$ ), 714.7 ( $[\text{Mo}_3\text{S}_4(\text{acac})_3]^+$ ).

Anal. Calc. for  $\text{C}_{47}\text{H}_{50}\text{F}_6\text{Mo}_3\text{N}_3\text{O}_7\text{PPdS}_4$ : C, 39.3; H, 3.5; N, 2.9%. Found: C, 39.0; H, 3.1; N, 3.3%.

Interaction of **1**, **2**, and **3** with  $\text{P}_4$  in the presence of  $\text{H}_2\text{O}$ . A solution of  $\text{P}_4$  (0.0025 g, 0.02 mmol) in THF (0.4 mL) was added at room temperature to a solution of complex **1**, **2**, or **3** (0.02 mmol) in 1 mL of solvent (DMF,  $\text{CH}_2\text{Cl}_2$ , or THF). Then,  $\text{H}_2\text{O}$  (0.043 mL, 2.4 mmol) was added dropwise to the reaction mixture. After 12 h of stirring,  $\text{TiNO}_3$  (0.021 g, 0.08 mmol) was added. An analysis of the reaction mixture was provided by  $^{31}\text{P}$  NMR spectroscopy after each of the consequent steps.

**Author Contributions:** Conceptualization, A.L.G. and D.G.Y.; methodology, A.L.G. and D.G.Y.; validation, A.L.G., M.N.S., O.G.S. and D.G.Y.; formal analysis, A.V.S., V.M.B. and K.R.K.; investigation, A.M.K. (Airat M. Kuchkaev), N.Y.S. and A.M.K. (Aidar M. Kuchkaev); data curation, A.M.K. (Airat M. Kuchkaev), N.Y.S. and A.M.K. (Aidar M. Kuchkaev); writing—original draft preparation, A.M.K. (Airat M. Kuchkaev) and N.Y.S.; writing—review and editing, A.L.G., M.N.S., O.G.S. and D.G.Y.; supervision, A.L.G. and D.G.Y.; project administration, A.L.G. and D.G.Y. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research was funded by the Russian Foundation for Basic Research, grant numbers 19-33-50136 and 18-33-20056.

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** Data sharing is not applicable to this article.

**Acknowledgments:** We acknowledge the financial support from the Russian Foundation for Basic Research (grants 19-33-50136 and 18-33-20056). The authors are thankful to the Russian Government Program of Competitive Growth of Kazan Federal University (Reg. Nr. AAAA-A19-119053190012-8) for performing of the NMR experiments in KFU and to the Government assignment for FRC Kazan Scientific Center of RAS (Reg. Nr. AAAA-A18-118041760011-2) for supporting this research activity dealing with the synthesis of the cluster complexes and phosphorus derivatives. The authors also gratefully acknowledge the CSF-SAC FRC KSC RAS for providing necessary facilities to carry out this work.

**Conflicts of Interest:** The authors declare no conflict of interest.

**Sample Availability:** Samples of the compounds **1**–**3** are available from the authors.

## References

1. Yakhvarov, D.G.; Kagiroy, R.M.; Sinyashin, O.G. Method of Producing Phosphorous Acid from White Phosphorus. Russian Patent 2460687, 25 February 2011.
2. Kagiroy, R.M.; Voloshin, A.V.; Rizvanov, I.K.; Sinyashin, O.G.; Yakhvarov, D.G. Activation and transformation of white phosphorus by palladium(II) complexes. *Russ. Chem. Bull.* **2010**, *59*, 1116–1118. [\[CrossRef\]](#)
3. Kagiroy, R.M.; Voloshin, A.V.; Kadirov, M.K.; Nizameev, I.R.; Sinyashin, O.G.; Yakhvarov, D.G. Selective synthesis of nanosized palladium phosphides from white phosphorus. *Mendeleev Commun.* **2011**, *21*, 201–203. [\[CrossRef\]](#)
4. Di Vaira, M.; Peruzzini, M.; Stoppioni, P.  $d^6$  metal systems for white phosphorus activation. *Comptes Rendus Chim.* **2010**, *13*, 935–942. [\[CrossRef\]](#)
5. Di Vaira, M.; Frediani, P.; Costantini, S.S.; Peruzzini, M.; Stoppioni, P. Easy hydrolysis of white phosphorus coordinated to ruthenium. *Dalton Trans.* **2005**, 2234–2236. [\[CrossRef\]](#) [\[PubMed\]](#)

6. Barbaro, P.; Di Vaira, M.; Peruzzini, M.; Seniori Costantini, S.; Stoppioni, P. Controlling the activation of white phosphorus: Formation of phosphorous acid and ruthenium-coordinated 1-hydroxytriphosphane by hydrolysis of doubly metalated P<sub>4</sub>. *Angew. Chem. Int. Ed.* **2008**, *47*, 4425–4427. [\[CrossRef\]](#)
7. Caporali, M.; Gonsalvi, L.; Kagirow, R.; Mirabello, V.; Peruzzini, M.; Sinyashin, O.; Stoppioni, P.; Yakhvarov, D. The first water-soluble tetraphosphorus ruthenium complex. Synthesis, characterization and kinetic study of its hydrolysis. *J. Organomet. Chem.* **2012**, *714*, 67–73. [\[CrossRef\]](#)
8. Akbayeva, D.N.; Di Vaira, M.; Seniori Costantini, S.; Peruzzini, M.; Stoppioni, P. Stabilization of the tautomers HP(OH)<sub>2</sub> and P(OH)<sub>3</sub> of hypophosphorous and phosphorous acids as ligands. *Dalton Trans.* **2006**, 389–395. [\[CrossRef\]](#)
9. Murata, T.; Gao, H.; Mizobe, Y.; Nakano, F.; Motomura, S.; Tanase, T.; Yano, S.; Hidai, M. Synthesis of Mixed-Metal Sulfido Clusters with a Cuboidal Mo<sub>3</sub>PdS<sub>4</sub> Core Which Coordinate Alkene to the Unique Palladium Site Surrounded by Sulfido Ligands. *J. Am. Chem. Soc.* **1992**, *114*, 8287–8288. [\[CrossRef\]](#)
10. Murata, T.; Mizobe, Y.; Gao, H.; Ishii, Y.; Wakabayashi, T.; Nakano, F.; Hidai, M.; Echizen, I.; Nanikawa, H.; Motomura, S.; et al. Syntheses of Mixed-Metal Sulfide Cubane-Type Clusters with the Novel PdMo<sub>3</sub>S<sub>4</sub> Core and Reactivities of the Unique Tetrahedral Pd Site Surrounded by Sulfide Ligands toward Alkenes, CO, tBuNC, and Alkynes. *J. Am. Chem. Soc.* **1994**, *116*, 3389–3398. [\[CrossRef\]](#)
11. Gushchin, A.L.; Laricheva, Y.A.; Sokolov, M.N.; Llusar, R. Tri- and tetranuclear molybdenum and tungsten chalcogenide clusters: On the way to new materials and catalysts. *Russ. Chem. Rev.* **2018**, *87*, 670–706. [\[CrossRef\]](#)
12. Sokolov, M.N.; Fedin, V.P.; Sykes, A.G. *Comprehensive Coordination Chemistry II*; McCleverty, J.A., Meyer, T.J., Eds.; Elsevier Ltd.: Amsterdam, The Netherlands, 2004; Volume 4, pp. 761–823.
13. Herbst, K.; Zanello, P.; Corsini, M.; D’Amelio, N.; Dahlenburg, L.; Brorson, M. A complete family of isostructural cluster compounds with cubane-like M<sub>3</sub>S<sub>4</sub>M’ cores (M = Mo, W; M’ = Ni, Pd, Pt): Comparative crystallography and electrochemistry. *Inorg. Chem.* **2003**, *42*, 974–981. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Hernandez-molina, R.; Sokolov, M.N.; Sykes, A.G. Behavioral Patterns of Heterometallic Cuboidal Derivatives of [M<sub>3</sub>Q<sub>4</sub>(H<sub>2</sub>O)<sub>9</sub>]<sup>4+</sup> (M = Mo, W; Q = S, Se). *Acc. Chem. Res.* **2001**, *34*, 223–230. [\[CrossRef\]](#) [\[PubMed\]](#)
15. Seino, H.; Hidai, M. Catalytic functions of cubane-type M<sub>4</sub>S<sub>4</sub> clusters. *Chem. Sci.* **2011**, *2*, 847–857. [\[CrossRef\]](#)
16. Algarra, A.G.; Fernández-Trujillo, M.J.; Safont, V.S.; Hernández-Molina, R.; Basallote, M.G. Combined kinetic and DFT studies on the stabilization of the pyramidal form of H<sub>3</sub>PO<sub>2</sub> at the heterometal site of [Mo<sub>3</sub>M’S<sub>4</sub>(H<sub>2</sub>O)<sub>10</sub>]<sup>4+</sup> clusters (M’ = Pd, Ni). *Dalton Trans.* **2009**, *4*, 1579–1586. [\[CrossRef\]](#)
17. Kryuchkova, N.A.; Ryzhikov, M.R.; Syrovashin, M.M. Interatomic Interactions in Heterometallic Cubane-Type Clusters with {Mo<sub>3</sub>S<sub>4</sub>M’} (M’ = Cu, Ni, Pd) Core. *J. Clust. Sci.* **2020**. [\[CrossRef\]](#)
18. Herbst, K.; Monari, M.; Brorson, M. Molecular metal sulfide cluster model for substrate binding to oil-refinery hydrodesulfurization catalysts. *Inorg. Chem.* **2002**, *41*, 1336–1338. [\[CrossRef\]](#)
19. Shibahara, T. Syntheses of sulphur-bridged molybdenum and tungsten coordination compounds. *Coord. Chem. Rev.* **1993**, *123*, 73–147. [\[CrossRef\]](#)
20. Hidai, M.; Kuwata, S.; Mizobe, Y. Synthesis and reactivities of cubane-type sulfido clusters containing noble metals. *Acc. Chem. Res.* **2000**, *33*, 46–52. [\[CrossRef\]](#)
21. Llusar, R.; Vicent, C. Trinuclear Molybdenum and Tungsten Cluster Chalcogenides: From Solid State to Molecular Materials. *Inorg. Chem. Focus III* **2006**, 105–120. [\[CrossRef\]](#)
22. Algarra, A.G.; Feliz, M.; Fernández-Trajiillo, M.J.; Llusar, R.; Safont, V.S.; Vicent, C.; Basallote, M.G. Unprecedented Solvent-assisted reactivity of hydrido W<sub>3</sub>CuS<sub>4</sub> cubane clusters: The Non-innocent behaviour of the Cluster-core unit. *Chem. A Eur. J.* **2009**, *15*, 4582–4594. [\[CrossRef\]](#)
23. Laricheva, Y.A.; Shmelev, N.Y.; Gushchin, A.L.; Sokolov, M.N. The study of the interaction of fullerene C<sub>60</sub> with palladium-containing molybdenum clusters {CS<sub>4</sub>} by spectroscopic and calculation methods. *Russ. J. Coord. Chem.* **2021**, in press.
24. Tao, Y.; Wang, B.; Zhao, J.; Song, Y.; Qu, L.; Qu, J. Friedel-Crafts-type allylation of nitrogen-containing aromatic compounds with allylic alcohols catalyzed by a [Mo<sub>3</sub>S<sub>4</sub>Pd(η<sup>3</sup>-allyl)] cluster. *J. Org. Chem.* **2012**, *77*, 2942–2946. [\[CrossRef\]](#) [\[PubMed\]](#)
25. Tao, Y.; Wang, B.; Wang, B.; Qu, L.; Qu, J. Highly efficient and regioselective allylation with allylic alcohols catalyzed by [Mo<sub>3</sub>S<sub>4</sub>Pd(η<sup>3</sup>-allyl)] clusters. *Org. Lett.* **2010**, *12*, 2726–2729. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Tao, Y.; Zhou, Y.; Qu, J.; Hidai, M. Highly efficient and regioselective allylic amination of allylic alcohols catalyzed by [Mo<sub>3</sub>PdS<sub>4</sub>] cluster. *Tetrahedron Lett.* **2010**, *51*, 1982–1984. [\[CrossRef\]](#)
27. Wakabayashi, T.; Ishii, Y.; Ishikawa, K.; Hidai, M. A Novel Catalyst with a Cuboidal PdMo<sub>3</sub>S<sub>4</sub> Core for the Cyclization of Alkynoic Acids to Enol Lactones. *Angew. Chem.* **1996**, *35*, 2123–2124. [\[CrossRef\]](#)
28. Sokolov, M.N.; Virovets, A.V.; Dybtsev, D.N.; Chubarova, E.V.; Fedin, V.P.; Fenske, D. Phosphorous acid and arsenious acid as ligands. *Inorg. Chem.* **2001**, *40*, 4816–4817. [\[CrossRef\]](#)
29. Sokolov, M.N.; Chubarova, E.V.; Virovets, A.V.; Llusar, R.; Fedin, V.P. Reactivity of Mo<sub>3</sub>PdS<sub>4</sub><sup>4+</sup> cluster: Evidence for new ligands PhP(OH)<sub>2</sub> and Ph<sub>2</sub>P(OH) and structural characterization of [Mo<sub>3</sub>(Pd(PPh<sub>3</sub>))S<sub>4</sub>(H<sub>2</sub>O)<sub>5</sub>Cl<sub>4</sub>].0.5CH<sub>3</sub>OH.3H<sub>2</sub>O. *J. Clust. Sci.* **2003**, *14*, 227–235. [\[CrossRef\]](#)
30. Oriwaki, K.M.; Oshida, R.Y.; Kashi, H.A. X-ray Structure Analysis Online Synthesis and Crystal Structure of a Sulfur-bridged Incomplete Cubane-type Molybdenum Cluster with a Tris(pyrazolyl)methanesulfonate Ligand. *X-ray Struct. Anal. Online* **2014**, *30*, 11–12. [\[CrossRef\]](#)



31. Ohki, Y.; Hara, R.; Munakata, K.; Tada, M.; Takayama, T.; Sakai, Y.; Cramer, R.E. Synthesis of  $[\text{Mo}_3\text{S}_4]$  Clusters from Half-Sandwich Molybdenum(V) Chlorides and Their Application as Platforms for  $[\text{Mo}_3\text{S}_4\text{Fe}]$  Cubes. *Inorg. Chem.* **2019**, *58*, 5230–5240. [\[CrossRef\]](#)
32. Pedrajas, E.; Sorribes, I.; Guillamón, E.; Junge, K.; Beller, M.; Llusar, R. Efficient and Selective N-Methylation of Nitroarenes under Mild Reaction Conditions. *Chem. A Eur. J.* **2017**, *23*, 13205–13212. [\[CrossRef\]](#)
33. Ohki, Y.; Uchida, K.; Hara, R.; Kachi, M.; Fujisawa, M.; Tada, M.; Sakai, Y.; Sameera, W.M.C. Cubane-Type  $[\text{Mo}_3\text{S}_4\text{M}]$  Clusters with First-Row Groups 4–10 Transition-Metal Halides Supported by  $\text{C}_5\text{Me}_5$  Ligands on Molybdenum. *Chem. A Eur. J.* **2018**, *24*, 17138–17147. [\[CrossRef\]](#) [\[PubMed\]](#)
34. Petrov, P.A.; Sukhikh, T.S. Phosphine-substituted Cubane Clusters with the  $\text{Mo}_3\text{S}_4\text{Ga}$  Core. *Russ. J. Coord. Chem.* **2019**, *45*, 333–339. [\[CrossRef\]](#)
35. Gushchin, A.L.; Sokolov, M.N.; Naumov, D.Y.; Fedin, V.P. Crystal structure of  $\text{K}_2[\text{Mo}_3(\text{PdPPh}_3)\text{S}_4(\text{C}_2\text{O}_4)_3(\text{H}_2\text{O})_3] \cdot 0.5\text{H}_2\text{O}$ . *J. Struct. Chem.* **2008**, *49*, 748–752. [\[CrossRef\]](#)
36. Laricheva, Y.A.; Gushchin, A.L.; Abramov, P.A.; Sokolov, M.N. Novel mixed-metal cubane-type  $\{\text{Mo}_3\text{NiS}_4\}$  and  $\{\text{Mo}_3\text{PdS}_4\}$  clusters coordinated with 2,2'-bipyridine type ligands. *Polyhedron* **2018**, *154*, 202–208. [\[CrossRef\]](#)
37. Algarra, G.; Gushchin, A.L.; Abramov, P.A.; Bustelo, E.; Llusar, R. Kinetics Aspects of the Reversible Assembly of Copper in Heterometallic  $\text{Mo}_3\text{CuS}_4$  Clusters with 4,4'-Di-tert-butyl-2,2'-bipyridine. *Inorg. Chem.* **2016**, *55*, 9912–9922. [\[CrossRef\]](#)
38. Ukai, T.; Kawazura, H.; Ishii, Y.; Bonnet, J.J.; Ibers, J.A. Chemistry of dibenzylideneacetone-palladium(0) complexes. Novel tris(dibenzylideneacetone)dipalladium(solvent) complexes and their reactions with quinones. *J. Organomet. Chem.* **1974**, *65*, 253–266. [\[CrossRef\]](#)
39. Pino-Chamorro, J.A.; Laricheva, Y.A.; Guillamón, E.; Fernández-Trujillo, M.J.; Bustelo, E.; Gushchin, A.L.; Shmelev, N.Y.; Abramov, P.A.; Sokolov, M.N.; Llusar, R.; et al. Cycloaddition of alkynes to diimino  $\text{Mo}_3\text{S}_4$  cubane-type clusters: A combined experimental and theoretical approach. *New J. Chem.* **2016**, *40*, 7872–7880. [\[CrossRef\]](#)
40. Hernández-Molina, R.; Sokolov, M.; Clegg, W.; Esparza, P.; Mederos, A. Preparation and characterization of triangular clusters  $[\text{M}_3\text{Q}_4(\text{acac})_3(\text{py})_3]^+$  ( $\text{M} = \text{Mo}, \text{W}$ ;  $\text{Q} = \text{S}, \text{Se}$ ). *Inorganica Chim. Acta* **2002**, *331*, 52–58. [\[CrossRef\]](#)