

Supplementary Materials: Synthesis and Characterization of Some New Bis-Pyrazolyl-Thiazoles Incorporating the Thiophene Moiety as Potent Anti-Tumor Agents

Sobhi M. Gomha, Mastoura M. Edrees and Farag M. A. Altalbawy




Figure S1. ¹H NMR spectrum of compound 3a.






Figure S2. ¹³C NMR spectrum of compound 3b.

Figure S3. ¹³C NMR spectrum of compound 3c.Figure S4. ¹H NMR spectrum of compound 3c.




Figure S5. ¹H NMR spectrum of compound 3d.




Figure S6. ¹³C NMR spectrum of compound 3e.

¹H spectrum TE26 in DMSO




Figure S7. ¹H NMR spectrum of compound **3h**.

¹H spectrum TE27 in DMSO





Figure S8. ¹H NMR spectrum of compound **3i**.






Figure S11. ^{13}C NMR spectrum of compound 5a.








Figure S12. ^1H NMR spectrum of compound **5b**.

Figure S13. ¹H NMR spectrum of compound 5f.Figure S14. ¹³C NMR spectrum of compound 5f.

Figure S15. ^1H NMR spectrum of compound 10.Figure S16. ^{13}C NMR spectrum of compound 10.