

**Supplementary Table S3. Haplotypes of women in the RM group and the control group.**

	<i>RM women</i> <i>2n=46</i>		<i>Controls</i> <i>2n=88*</i>		<b>OR</b>	<b>95% C.I.</b>		<b>P</b>
						<b>Lower</b>	<b>Upper</b>	
UTR-1	13	28.3%	28	31.8%	0.90	0.413	1.965	0.792
UTR-2	16	34.8%	23	26.1%	1.60	0.742	3.451	0.231
UTR-3	5	10.9%	8	9.1%	1.28	0.394	4.159	0.681
UTR-4	5	10.9%	14	15.9%	0.68	0.229	2.019	0.487
UTR-5	4	8.7%	4	4.5%	2.10	0.499	8.789	0.312
UTR-7	3	6.5%	7	8.0%	0.85	0.209	3.440	0.817
UTR-8	0	0.0%	0	0.0%	x	x	x	x
UTR-18	0	0.0%	3	3.4%	0.00	0.000		0.999
UTR-N	0	0.0%	1	1.1%	0.00	0.000		1.000
	46	100%	88	100%				

All univariate logistic regression analysis. *P*, *p* value; *OR*, odds ratio; *95% CI*, 95% confidence interval; n.a, not applicable. \*In 2 control subjects the UTR haplotype could not be defined (4%).

The 3'UTR haplotype nomenclature is consistent with publication by Castelli et al.