

Potential associations among alteration of salivary miRNAs, saliva microbiome structure and cognitive impairments in autistic children

Marco Ragusa^{1,2§}, Maria Santagati^{3§}, Federica Mirabella^{1§}, Giovanni Lauretta¹, Matilde Cirigliaro¹, Duilia Brex¹, Cristina Barbagallo¹, Carla Noemi Domini⁴, Mariangela Gulisano⁴, Rita Barone⁴, Laura Trovato³, Salvatore Oliveri³, Gino Mongelli^{3,5}, Ambra Spitale³, Davide Barbagallo¹, Cinzia Di Pietro^{1†}, Stefania Stefani^{3†}, Renata Rizzo^{4†}, Michele Purrello^{1*}.

	White's non parametric t-test	Mann-Whitney-Kruskal-Wallis test	T-test/Anova
	P-values	P-values	P-values
g_Filifactor	9.99E-04	0.029812	0.0016537
g_Rothia	0.011	0.030216	0.012415
g_Actinobacillus	0.016	0.41668	0.016316
f_weeksellaceae	0.016	0.038872	0.011177
g_Ralstonia	0.023	0.046734	0.078283
f_pasteurellaceae	0.029	0.021476	0.122916
g_Aggregatibacter	0.035	0.16022	0.020473
g_Tannerella	0.044	0.038368	0.041844
c_TM7	0.045	0.027238	0.015501
g_Moryella	0.045	0.027239	0.033352

Supplementary Table S1. Statistical analysis of the bacterial abundance at the genus level in ASD and NUC groups by a two-sided White's non-parametric t-test, Mann-Whitney-Kruskal-Wallis test and T-test/Anova, p-value <0.05.