

Supplementary data

Bioactive Phytochemicals from *Salix pseudolasiogyne* Twigs: Anti-Adipogenic Effect of 2'-O-Acetylsalicortin in 3T3-L1 Cells

Hee Jung Kim ^{1,2,†}, Yoon Seo Jang ^{3,†}, Ji Won Ha ³, Moon-Jin Ra ⁴, Sang-Mi Jung ⁴, Jeong-Nam Yu ⁵, Kyunga Kim ⁶,
Ki Hyun Kim ^{3,*} and Sung Hee Um ^{1,2,7,*}

¹ Department of Molecular Cell Biology, Samsung Biomedical Research Institute, School of Medicine, Sungkyunkwan University, Suwon 16419, Korea

² Department of Health Sciences and Technology, Samsung Medical Center, Samsung Advanced Institute for Health Sciences and Technology, Sungkyunkwan University, Seoul 06351, Korea

³ School of Pharmacy, Sungkyunkwan University, Suwon 16419, Korea

⁴ Hongcheon Institute of Medicinal Herb, Hongcheon-gun 25142, Korea

⁵ Nakdonggang National Institute of Biological Resources, Sangju 37242, Korea

⁶ Biomedical Statistics Center, Research Institute for Future Medicine, Samsung Medical Center, Seoul 06351, Korea

⁷ Biomedical Institute Convergence, Sungkyunkwan University, Suwon 16419, Korea

* Correspondence: khkim83@skku.edu (K.H.K.); shum@skku.edu (S.H.U.); Tel.: +82-31-290-7700 (K.H.K.); Tel.: +82-31-299-6123 (S.H.U.)

† These authors contributed equally to this work.

Figure S1. ^1H NMR spectrum of oregonin (**1**) (CD_3OD , 850 MHz)




Figure S2. ^1H NMR spectrum of 2'-*O*-acetylsalicortin (**2**) (CD_3OD , 850 MHz)






Figure S3. HPLC chromatogram of fractions (P1 – P4).

- Column : C18
- Solvent : MeOH Gradient 30~50%
- Sample Wt : 1.0 g



Pure compounds

Compounds 1 and 2