



Formate dehydrogenase improves the resistance to formic acid and acetic acid simultaneously in *Saccharomyces cerevisiae*

Cong Du ^{1,†}, Yimin Li ^{1,†}, Ruijuan Xiang ¹ and Wenjie Yuan ^{1,2,*}

¹ School of Bioengineering, Dalian University of Technology, Dalian 116024, China; 18742515846@163.com (C.D.); yiminli@dlut.edu.cn (Y.L.); xiangruijuan@bluepha.com (R.X.)

² Ningbo Research Institute, Dalian University of Technology, Ningbo 315000, China

* Correspondence: ywj@dlut.edu.cn

† These authors contributed equally to this work.

Table S1. Primers of RT-PCR.

Primer	Sequence	Product size
RCK1-F	CACGGATGACCAAGCTCCTGTTG	145
RCK1-R	CGAAACCAGTTGTGGATGGCAAC	145
DAK2-F	AGAAGTTGCCGAAGTTGCTCAGTC	80
DAK2-R	CAGGAACCGTGCAGTGATCAAGAG	80
CTT1-F	CTGGAGATAAGGCTGCTGAAGTC	132
CTT1-R	TTGCTTGTTCGGGTGTCATTGTTG	132
PDR15-F	TTTCGCATACAAGGGACACCAAGG	84
PDR15-R	CCGCAATGGATACACGCTTCTTC	84
FDH1-F	CGGTAGACAGGGAGTTGAAAGACG	114
FDH1-R	CGGTTACACAGAGCTTCAGGTTAGG	114
PMA1-F	AAGGCTAAGGACGCTTGACCAAG	92
PMA1-R	GGAGATTGACACAACGGCAGTAACC	92
ACT1-F	AAGTGTGATGTCGATGTCCGTAAGG	120
ACT1-R	AGATGGAGCCAAGCGGTGATTTC	120




Figure S1. Clusters analysis and Volcano Plot of DEGs. (a) Cluster analysis of differentially expressed genes; (b) Volcano Plot, red plots were up-regulated, blue plots were down-regulated, blue plots were not significantly different.




Figure S2. Statistics of pathway enrichment.