

SUPPLEMENTARY MATERIAL

The Impact of Isoxazole Derivatives on Pathogenic Biofilm and Cytotoxicity of Fibroblast Cell Line

Urszula Bąchor ¹, Adam Junka ^{2,*}, Małwina Brożyna ² and Marcin Mączyński ¹

¹ Department of Organic Chemistry and Drug Technology, Faculty of Pharmacy, Wrocław Medical University, 50-556 Wrocław, Poland

² Department of Pharmaceutical Microbiology and Parasitology, Unique Application Model Laboratory, Faculty of Pharmacy, Wrocław Medical University, 50-556 Wrocław, Poland

* Correspondence: adam.junka@umw.edu.pl; Tel.: +48-71-784-06-75

Figure S1. ^1H NMR spectrum of compound MAL1.....	2
Figure S2. ^{13}C NMR spectrum of compound MAL1.....	2
Figure S3. ESI-MS spectrum of compound MAL1.....	3
Figure S4. ^1H NMR spectrum of compound MAL2.....	3
Figure S5. ^{13}C NMR spectrum of compound MAL2.....	4
Figure S6. ESI-MS spectrum of compound MAL2	4
Figure S7. ^1H NMR spectrum of compound MAL3.....	5
Figure S8. ^{13}C NMR spectrum of compound MAL3.....	5
Figure S9. ESI-MS spectrum of compound MAL3	6
Figure S10. ^1H NMR spectrum of compound MAL4.....	6
Figure S11. ^{13}C NMR spectrum of compound MAL4.....	7
Figure S12. ESI-MS spectrum of compound MAL4	7
Figure S13. ^1H NMR spectrum of compound MAL5.....	8
Figure S14. ^{13}C NMR spectrum of compound MAL5	8
Figure S15. ESI-MS spectrum of compound MAL5	9
Figure S16. ^1H NMR spectrum of compound PUB9	9
Figure S17. ^{13}C NMR spectrum of compound PUB9	10
Figure S18. ESI-MS spectrum of compound PUB9	10
Figure S19. ^1H NMR spectrum of compound PUB10	11
Figure S20. ^{13}C NMR spectrum of compound PUB10	11
Figure S21. ESI-MS spectrum of compound PUB10	12




Figure S1. ¹H NMR spectrum of compound MAL1




Figure S2. ¹³C NMR spectrum of compound MAL1




Figure S3. ESI-MS spectrum of compound MAL1




Figure S4. ^1H NMR spectrum of compound MAL2




Figure S5. ¹³C NMR spectrum of compound MAL2




Figure S6. ESI-MS spectrum of compound MAL2




Figure S7. ¹H NMR spectrum of compound MAL3




Figure S8. ¹³C NMR spectrum of compound MAL3




Figure S9. ESI-MS spectrum of compound MAL3




Figure S10. ^1H NMR spectrum of compound MAL4




Figure S11. ^{13}C NMR spectrum of compound MAL4




Figure S12. ESI-MS spectrum of compound MAL4




Figure S13. ¹H NMR spectrum of compound MAL5




Figure S14. ¹³C NMR spectrum of compound MAL5




Figure S15. ESI-MS spectrum of compound MAL5




Figure S16. ^1H NMR spectrum of compound PUB9




Figure S17. ^{13}C NMR spectrum of compound PUB9




Figure S18. ESI-MS spectrum of compound PUB9




Figure S19. ¹H NMR spectrum of compound PUB10




Figure S20. ¹³C NMR spectrum of compound PUB10




Figure S21. ESI-MS spectrum of compound PUB10

Table S1. The values of Minimal Inhibitory Concentration (MIC), Minimal Biofilm Eradication Concentration (MBEC) and Antibiofilm Dressing Activity Measurement (A.D.A.M.) performed for octenidine dihydrochloride-based antiseptic agent (method's usability control).

	<i>S.aureus</i>	<i>P.aeruginosa</i>	<i>C.albicans</i>
MIC [mg/L]	1	0,78	1
MBEC [mg/L]	62,5	1000	62,5
A.D.A.M. [%]reduction]	98	95	89