

Serum-resistant ternary DNA-polyplexes for suicide gene therapy of uterine leiomyoma

Anna Egorova ¹, Sofia Shtykalova ¹, Marianna Maretina ¹, Svetlana Freund ¹, Alexander Selutin ², Natalia Shved ¹, Sergei Selkov ², and Anton Kiselev ^{1,*}


¹ Department of Genomic Medicine named after V.S. Baranov, D.O. Ott Research Institute of Obstetrics, Gynecology and Reproductology, Mendelevskaya line 3, 199034 Saint-Petersburg, Russia; egorova_anna@yahoo.com (A.E.); sofia.shtykalova@gmail.com (S.S.); marianna0204@gmail.com (M.M.); svetafreundmax@gmail.com (S.F.); natashved@mail.ru (N.S.);

² Department of Immunology and Intercellular Interactions, D.O. Ott Research Institute of Obstetrics, Gynecology and Reproductology, Mendelevskaya line 3, 199034 Saint-Petersburg, Russia; a_selutin@yahoo.com (A.S.); selkovsa@mail.ru (S.S.)


* Correspondence: ankiselev@yahoo.co.uk; Tel.: +7-812-328-9809

Contents

Figure S1. Evaluation of the cytotoxicity after transfection of PANC-1 cells with DNA/R6p/E6p and DNA/R6p/E6p-cRGD polyplexes.....	3
Figure S2. Typical micrographs of the ternary polyplexes obtained by transmission electronic microscopy.....	5




(a)




(b)


Figure S1. Evaluation of the cytotoxicity after transfection of PANC-1 cells with DNA/R6p/E6p and DNA/R6p/E6p-cRGD polyplexes formed at different P/N/C ratios: **(a)** N/P ratio of 8/1; **(b)** N/P ratio of 12/1. Values are the mean \pm SD of the mean of triplicates.




(a)



(b)



(c)



(d)

Figure S2. Typical micrographs of the ternary polyplexes obtained by transmission electronic microscopy: **(a)** DNA/R6p/E6p at P/N/C ratio 1/8/4; **(b)** DNA/R6p/E6p at P/N/C ratio 1/12/6; **(c)** DNA/R6p/E6p at P/N/C ratio 1/12/12; **(d)** DNA/R6p/E6p-cRGD at P/N/C ratio 1/12/6.