

Synthesis and evaluation of self-assembling properties of 3-(3,5-difluoro-3,5-bis((alkoxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodides


Nadiia Pikun¹, Davis Lacis¹, Arkadij Sobolev¹, Martins Rucins¹, Mara Plotniece², Karlis Pajuste¹, Aiva Plotniece^{1,2*}

¹ Latvian Institute of Organic Synthesis, Aizkraukles 21, Riga LV-1006, Latvia, nadiia@osi.lv (N.P.), davis.lacis@osi.lv (D.L), arkady@osi.lv (A.S.), rucins@osi.lv (M.R.), kpajuste@osi.lv (K.P.)


² Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Riga Stradiņš University, Dzirciema 16, Riga LV-1007, Latvia, mara.plotniece@rsu.lv (M.P.)

* Correspondence: aiva@osi.lv (A.P.)


¹H NMR spectrum of diethyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3a**)




^{19}F NMR spectrum of diethyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3a**)




^1H NMR spectrum of diethyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3a**)




^1H NMR spectrum of dinonyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3b**)




^{19}F NMR spectrum of dinonyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3b**)




^{13}C NMR spectrum of dinonyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3b**)




^1H NMR spectrum of 3-(3,5-bis(ethoxycarbonyl)-3,5-difluoro-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4a**)




^{19}F NMR spectrum of 3-(3,5-bis(ethoxycarbonyl)-3,5-difluoro-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4a**)




^{13}C NMR spectrum of 3-(3,5-bis(ethoxycarbonyl)-3,5-difluoro-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4a**)




^1H NMR spectrum of 3-(3,5-difluoro-3,5-bis((nonyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4b**)




^{19}F NMR spectrum of 3-(3,5-difluoro-3,5-bis((nonyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4b**)




^{13}C NMR spectrum of 3-(3,5-difluoro-3,5-bis((nonyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4b**)




HRMS data of diethyl 2,4-diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3a**)




HRMS data of dinonyl 2,4- diacetyl-2,4-difluoro-3-(pyridin-3-yl)pentanedioate (**3b**)



HRMS data of 3-(3,5-difluoro-3,5-bis((ethyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4a**)



HRMS data of 3-(3,5-difluoro-3,5-bis((nonyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4b**)



Example of DLS results for freshly prepared nanoparticles formed by 3-(3,5-difluoro-3,5-bis((nonyloxy)carbonyl)-2,6-dioxoheptan-4-yl)-1-methylpyridin-1-ium iodide (**4b**)

System

Temperature (°C):	25,1	Duration Used (s):	10
Count Rate (kcps):	147,9	Measurement Position (mm):	3,00
Cell Description:	Disposable micro cuvette (40µl)	Attenuator:	10

Results

	Size (d.nm):	% Intensity:	St Dev (d.n...
Z-Average (d.nm): 371,2	Peak 1: 432,9	100,0	148,2
Pdl: 0,309	Peak 2: 0,000	0,0	0,000
Intercept: 0,447	Peak 3: 0,000	0,0	0,000

Result quality Good

