

Table S1. List of zooplankton taxa identified in bladderwort traps and zooplankton tow samples during the study period. Taxa were categorized as either littoral (which can also mean benthic) or pelagic based on their typical life history and ecology. Calanoida, Cyclopoida, Harpacticoida, and Nauplii were grouped together as Copepoda for analyses.

Taxa	Littoral or pelagic taxon	Reference example
<i>Bosmina</i>	Pelagic	Barnett et al. (2007)
<i>Camptocercus</i>	Littoral	Adamczuk (2014)
<i>Chydorus</i>	Littoral	Barnett et al. (2007)
Calanoida	Pelagic	Barnett et al. (2007)
Cyclopoida	Pelagic	Barnett et al. (2007)
<i>Daphnia</i>	Pelagic	Barnett et al. (2007)
<i>Eurycercus</i>	Littoral	Lemke and Benke (2004)
Harpacticoida	Littoral	Kurashov (1996)
Hydrachnidia	Littoral	Sychra et al. (2010)
Nauplii	Pelagic	Barnett et al. (2007)
<i>Ophryoxus</i>	Littoral	Fryer (1974)
Ostracoda	Littoral	Benzie (1989)
<i>Pleuroxus</i>	Littoral	Frey (1988)

References

1. Adamczuk, M. Niche separation by littoral-benthic Chydoridae (Cladocera, Crustacea) in a deep lake - potential drivers of their distribution and role in littoral-pelagic coupling. *J. Limnol.* **2014**, 73, 490–501.
2. Barnett, A.J.; Finlay, K.; Beisner, B.E. Functional diversity of crustacean zooplankton communities: towards a trait-based classification. *Freshwater Biol.* **2007**, 52, 796–813.
3. Benzie, J.A.H. The distribution and habitat preference of ostracods (Crustacea: Ostracoda) in a coastal sand-dune lake, Loch of Strathbeg, north-east Scotland. *Freshw. Biol.* **1989**, 22, 309–321.
4. Frey, D.G. Separation of *Pleuroxus laevis* Sars, 1861 from two resembling species in North America: *Pleuroxus straminius* Birge, 1879 and *Pleuroxus chiangi* n.sp. (Cladocera, Chydoridae). *Can. J. Zool.* **1988**, 66, 2534–2563.

5. Fryer, G. Evolution and adaptive radiation in the Macrothricidae (Crustacea: Cladocera): a study in comparative functional morphology and ecology. *Phil. Trans. R. Soc. Lond. B* **1974**, 269, 137–274.
6. Kurashov, E.A. Distribution and summer diapause of *Canthocamptus staphylinus* (Jurine) (Copepoda: Harpacticoida) in Lake Ladoga. *Hydrobiologia* **1996**, 320, 191–196.
7. Lemke, A.M.; Benke, A.C. Growth, reproduction, and production dynamics of a littoral microcrustacean, *Eurycercus vernalis* (Chydoridae), from a southeastern wetland, USA. *J. North Am. Benthol. Soc.* **2004**, 23, 806–823.
8. Sychra, J.; Adámek, Z.; Petřivalská, K. Distribution and diversity of littoral macroinvertebrates within extensive reed beds of a lowland pond. *Ann. Limnol.* **2010**, 46, 281–289.