



# Article Discovery of Bioactive Indole-Diketopiperazines from the Marine-Derived Fungus *Penicillium brasilianum* Aided by Genomic Information

Ya-Hui Zhang <sup>1,2,+</sup>, Ce Geng <sup>3,+</sup>, Xing-Wang Zhang <sup>3</sup>, Hua-Jie Zhu <sup>2</sup>, Chang-Lun Shao <sup>1,4</sup>, Fei Cao <sup>2,\*</sup> and Chang-Yun Wang <sup>1,4,5,\*</sup>

- <sup>1</sup> Key Laboratory of Marine Drugs, the Ministry of Education of China, School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, China
- <sup>2</sup> College of Pharmaceutical Sciences, Key Laboratory of Pharmaceutical Quality Control of Hebei Province, Hebei University, Baoding 071002, China
- <sup>3</sup> Shandong Provincial Key Laboratory of Synthetic Biology, CAS Key Laboratory of Biofuels at Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao 266101, China
- <sup>4</sup> Laboratory for Marine Drugs and Bioproducts, Qingdao National Laboratory for Marine Science and Technology, Qingdao 266237, China
- <sup>5</sup> Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao 266003, China
- \* Correspondence: caofei542927001@163.com (F.C.); changyun@ouc.edu.cn (C.-Y.W.)
- + These authors contributed equally to this work.

Received: 2 August 2019; Accepted: 29 August 2019; Published: 1 September 2019

**Abstract:** Identification and analysis of the whole genome of the marine-derived fungus *Penicillium brasilianum* HBU-136 revealed the presence of an interesting biosynthetic gene cluster (BGC) for nonribosomal peptide synthetases (NRPS), highly homologous to the BGCs of indole-diketopiperazine derivatives. With the aid of genomic analysis, eight indole-diketopiperazines (**1**–**8**), including three new compounds, spirotryprostatin G (**1**), and cyclotryprostatins F and G (**2** and **3**), were obtained by large-scale cultivation of the fungal strain HBU-136 using rice medium with 1.0% MgCl<sub>2</sub>. The absolute configurations of **1**–**3** were determined by comparison of their experimental electronic circular dichroism (ECD) with calculated ECD spectra. Selective cytotoxicities were observed for compounds **1** and **4** against HL-60 cell line with the IC<sub>50</sub> values of 6.0 and 7.9 μM, respectively, whereas **2**, **3**, and **5** against MCF-7 cell line with the IC<sub>50</sub> values of 7.6, 10.8, and 5.1 μM, respectively.

**Keywords:** biosynthetic gene cluster (BGC); indole-diketopiperazine; *Penicillium brasilianum*; cytotoxicities

# 1. Introduction

In the past decades, marine-derived fungi have attracted more and more attention to medicinal chemists because of their ability to produce biologically active compounds with diverse structures for drug discovery [1]. Especially, one fungus species could generate structurally versatile natural products because it consists of a large number of biosynthetic gene clusters (BGCs) associated with its secondary metabolism [2,3]. For example, several bioactive compounds have been generated by the same fungus from different sources, including antifungal diketopiperazine-type alkaloids [4], antibacterial dimeric diketopiperazines [5], cytotoxic prenyl asteltoxin derivatives [6], and anti-virus highly oxygenated cyclopiazonic acid-derived alkaloids [7]. However, it has been proven that most of the genes encoding secondary metabolites in fungi are cryptic under certain culture conditions [8]. In recent years, due to the advances in genome sequencing and bioinformatics-based predictions of encoded natural products in BGCs, it has been enabled to obtain structurally-unique secondary metabolites from fungi with the assistance of genomic information [8–10].

During our investigation focused on discovering bioactive natural products from marinederived fungi [11–14], two known cytotoxic diketopiperazine alkaloids, spirocyclic diketopiperazine alkaloid (4) [15] and cyclotryprostatin B (5) [16], were discovered from the marine-derived fungus *Penicillium brasilianum* HBU-136 by using rice medium in the first large-scaled fermentation. After gaining these two known diketopiperazine alkaloid derivatives, we tried to locate their BGC in the genome of the fungus HBU-136. A search of the genome sequence using antiSMASH 4.2 found that the strain harbors multiple BGCs for non-ribosomal peptide synthetases (NRPS), one of which shows high similarity to the BGC of fumitremorgins A–C in the fungus *Aspergillus fumigatus* [17–19]. By gene-to-gene comparisons, it is disclosed that the main difference between the BGC in HBU-136 (*ctp*) against in *Aspergillus fumigatus* (*ftm*) lies at the tailoring genes (Table S2 and Figure S45), which suggested that the BGC in HBU-136 may having the ability to produce new derivatives. Based on this information, we tried more fermentation conditions with LC-MS analysis, and finally found three new indole-diketopiperazines (1–3) from the rice medium with 1.0% MgCl<sub>2</sub> fermentation. Herein, we report the isolation, structural characterization, putative biosynthetic relation and biological activities of these compounds.

# 2. Results and Discussion

### 2.1. Identification and Analysis of the Indole-Diketopiperazine BGC

The whole genome sequence of *Penicillium brasilianum* HBU-136 was sequenced within HiSeq X10 platform (Illumina, CA, USA) (150 fold) and further assembled into 937 contigs (35.5 Mb) with *de novo* assembly software ABySS and SPAdes. For further accurate BGCs mining and comparison, all the genes were predicted by gene finding software GeneMarks ES/ET and Augustus with self-training. Then, 50 secondary metabolite gene clusters were proposed by using antiSMASH 4.2 as a gene cluster mining tool (Table S1). Thereinto, 12 out of the 50 BGCs were described as NRPS, including a BGC highly homologous to the fumitremorgin BGC (8 out of 9 genes show similarity Table S2) [17–19]. This highly similar gene cluster involved an NRPS gene (*ctpNRPS*), three cytochrome P450 genes (*ctpP450-1/2/3*), one oxmethyltransferase gene (*ctpOMT*), two prenyltransferase genes (*ctpPT-1/2*), and one putative oxidase gene (*ctpOx*) (Table S2).

By gene-to-gene comparison, it suggested that the proposed gene clusters were consistent well with the assembly of the dipeptide skeleton for indole-diketopiperazines (Figure 1). The key biosynthetic step of dipeptide skeleton started from two successive amidation reactions catalyzed by the *ctpNRPS* gene and led to the formation of brevianamide F [20,21]. Then *ctpPT-1* gene mediated the isopentane addition and delivered the product directly to *ctpP450-2* gene, who catalyzed the tetrahydropyridine ring formation, or to *ctpOMT* gene to get the methoxyl first before the ring-formation step [20,21]. The formed cyclotryprostatin skeleton could be further transformed into spirotryprostatin-type by *ctpP450-3* gene via a radical formation and migration eliciting semipinacol-type rearrangement mechanism [22]. P450s, the most versatile biocatalysts in nature, have been proved to catalyze such uncommon reactions during skeleton constructing processes in many cases [23]. After the core skeletons constructed, the structural diversity of both spirotryprostatins and cyclotryprostatins mainly come from the tailoring reactions of *ctpP450-3* gene to complete hydroxylation reactions, *ctpP450-1* gene to catalyze the benzene hydroxylation, and *ctpOMT* gene to mediate the methylation reaction.

More intriguingly, the gene *ctpP450-3*, whose corresponding gene in *ftm* is the multi-functional gene *ftmG*, was also inferred to encode a muti-functional P450 according to their sequence identity, suggesting that new polyhydroxylated indole-diketopiperazines could be generated by the fungal strain HBU-136. At this point, we tried eight different media conditions (C1–C8) with LC-MS screening to mine more biosynthetic products for *ctp*. Guided by HPLC-MS, it was found that three previously unrecognized compounds exhibiting the typical indole UV spectrum were produced by the strain HBU-136 when 1.0% MgCl<sub>2</sub> was added to rice medium (C-8). According to previous literature, the addition of MgCl<sub>2</sub> into the medium was reported to impact the secondary metabolism [13,24], which was confirmed once again by our research. Finally, eight indole-diketopiperazines (**1**–

8), including three new compounds, spirotryprostatin L (1) and cyclotryprostatins F–G (2–3), and five known analogs, spirocyclic diketopiperazine alkaloid (4) [15], cyclotryprostatin B (5) [16], 20-hydroxycyclotryprostatin B (6) [25],  $12\beta$ -hydroxy- $13\alpha$ -ethoxyverruculogen TR-2 (7) [26], and an unnamed compound (8) [27], were then obtained (Figure 2).



**Figure 1.** (**A**) The biosynthetic gene cluster of cyclotryprostatin from *Penicillium brasilianum (ctp)*. (**B**) Proposed biosynthetic pathway of the isolated indole-diketopiperazines.



Figure 2. Chemical structures of compounds 1-8.

## 2.2. Structure Elucidation

Spirotryprostatin L (1), a yellow amorphous powder, was obtained with the molecular formula C<sub>22</sub>H<sub>25</sub>N<sub>3</sub>O<sub>7</sub> (12 degrees of unsaturation) by positive HRESIMS. The IR absorption bands at 3447 cm<sup>-1</sup> and 1647 cm<sup>-1</sup> indicated the presences of hydroxyl and carbonyl groups. The characteristic UV absorption maxima at 220, 249, 287, 387 nm suggested a spirotryprostatin skeleton for **1** [15,16]. The NMR signals in **1** (Table 1 and 2) were representative of one keto carbonyl (C-3), two amide carbonyls (C-11 and C-17), one 1,3,4-trisubstituted aromatic unit (C-3a, C-4, C-5, C-6, C-7, and C-7a), and three methyls (one oxygenated) (6-OCH<sub>3</sub>, C-22, and C-21). These structural features suggested a prenylated indole-diketopiperazine nucleus of **1**, structurally similar to the known compound **4** which was previously obtained from the fungus *Aspergillus fumigatus* [15]. A detailed analysis for HMBC

correlations of **1** (from 12-OH to C-12 and C-13) (Figure 3) indicated that an additional hydroxyl group [ $\delta_{\rm H}$  5.14 (1H, brs, 12-OH) and  $\delta_{\rm C}$  90.2 (C-12)] was attached to C-12. Based on the systematic 2D NMR data analyses, the plane structure of **1** was determined.

No.	1	2	3
1		7.86, s	7.97, s
4	7.50, d (8.7)	7.48, d (8.4)	7.58, d (7.6)
5	6.45, dd (8.7, 2.0)	6.80, dd (8.4, 2.0)	7.17, dtd (8.0, 7.6, 0.9)
6			7.17, dtd (8.0, 7.6, 0.9)
7	6.27, d (2.0)	6.86, d (2.0)	7.37, d (7.6)
8	4.70, s	5.02, s	4.79, s
12			4.38, dd (10.7, 6.3)
13	2.35, m	2.40, m	2.50, m
	2.27, m	1.98, m	2.01, m
14	2.15, m	2.10, s	2.12, m
	2.00, m	1.98, m	2.01, m
15	3.74, m	3.80, m	3.76, m
	3.54, m	3.70, m	3.71, m
18	4.83, d (9.5)	6.36, d (9.6)	6.68, d (9.7)
19	4.79, d (9.5)	5.33, d (9.6)	5.57, d (9.7)
21	1.56, s	2.04, s	1.80, s
22	1.79, s	1.77, s	2.07, s
8-OH	4.96, brs		
12-OH	5.14, brs		
6-OCH <sub>3</sub>	3.85, s	3.82, s	
9-OH	8.51, brs		
8-OCH <sub>3</sub>		3.30, s	3.38, s

**Table 1.** <sup>1</sup>Η NMR Data (δ) of **1–3** (500 MHz, CDCl<sub>3</sub>, *J* in Hz).

**Table 2.** <sup>13</sup>C NMR Data (*δ*) of **1–3** (125 MHz, CDCl<sub>3</sub>).

	-		
No.	1	2	3
2	75.0, C	132.7, C	135.2, C
3	200.6, C	104.8, C	105.7, C
3a	112.4, C	122.5, C	128.5, C
4	127.4, CH	118.9, CH	118.2, CH
5	110.8, CH	110.1, CH	120.7, CH
6	169.7, C	156.6, C	122.3, CH
7	94.9, CH	95.3 <i>,</i> CH	111.3, CH
7a	165.4, C	137.1, C	135.9 <i>,</i> C
8	73.9 <i>,</i> CH	74.9, CH	76.8, CH
9	86.1, C	85.7, C	84.9, C
11	167.4, C	166.3, C	167.2, C
12	90.2, C	86.7, C	60.1, CH
13	35.1, CH <sub>2</sub>	36.5, CH <sub>2</sub>	29.8, CH <sub>2</sub>
14	21.3, CH <sub>2</sub>	19.2, CH <sub>2</sub>	22.3, CH <sub>2</sub>
15	45.3, CH <sub>2</sub>	45.4, CH <sub>2</sub>	46.0, CH <sub>2</sub>
17	166.0, C	165.9 <i>,</i> C	166.0, C
18	55.7, CH	49.3, CH	49.2, CH
19	119.4, CH	122.8, CH	123.6, CH
20	142.4, C	138.2, C	138.2, C
21	26.1, CH <sub>3</sub>	18.4, CH <sub>3</sub>	26.2, CH <sub>3</sub>
22	18.8, CH <sub>3</sub>	26.1, CH <sub>3</sub>	18.4, CH <sub>3</sub>
6-OCH <sub>3</sub>	56.0, CH <sub>3</sub>	55.8, OCH3	
8-OCH <sub>3</sub>		57.0, OCH3	56.8, OCH3



**Figure 3.** <sup>1</sup>H-<sup>1</sup>H COSY and key HMBC correlations of **1** and **2**. In order to assign the relative configuration of **1**, its NOESY experiment was carried out.

The NOESY correlations of **1** between H-7 and 8-OH, 8-OH and 9-OH, 9-OH and 12-OH, 12-OH and H-18, and H-8 and H-19 (Figure 4) suggested that H-18, 8-OH, 9-OH, and 12-OH were located on the same face of the molecule, and H-7 had a *cis* relationship with 8-OH/9-OH/12-OH/H-18. The electronic circular dichroism (ECD) spectrum of **1** was tested and calculated to investigate the absolute configuration of **1**. Time-dependent density functional theory (TD-DFT) method was used for ECD calculation of the molecule (2*S*,8*S*,9*R*,12*R*,18*S*)-**1** at the B3LYP/6-311++G(2d,p)//B3LYP/6-311++G(d) level. Boltzmann statistics with a standard deviation of  $\sigma$  0.3 eV was used for the ECD simulation. The calculated ECD spectrum of (2*S*,8*S*,9*R*,12*R*,18*S*)-**1** was in good accordance with the experimental ECD data of **1** (Figure 5), revealing that the absolute configuration of **1** should be 2*S*,8*S*,9*R*,12*R*,18*S*. Additionally, the above conclusion was verified by the fact that the experimental ECD spectrum and the specific optical rotation (OR) value of **1** were similar to those of **4** ([ $\alpha$ ] $_{D^{20}}$  = + 144.6 (*c* 1.00, MeOH) for **1** *vs* [ $\alpha$ ] $_{D^{20}}$  = + 147.2 (*c* 0.1, CHCl<sub>3</sub>) for **4**) [15]. Thus, compound **1** was defined as a 12-hydroxylation derivative of **4**.



Figure 4. The key NOESY correlations of 1.



Figure 5. Experimental ECD of 1 and 4, and calculated ECD of (2S,8S,9R,12R,18S)-1.

Cyclotryprostatin F (2) was also obtained as a yellow amorphous powder, whose molecular formula was deduced as  $C_{23}H_{27}N_3O_6$  from HRESIMS. The NMR data of 2 (Table 1; Table 2) suggested that it also possessed a diketopiperazine skeleton, which was considered similar to the known compound 5 [16] through detailed NMR analyses of them. Combining the upfield shifted C-

constants, NOESY correlations and ECD Cotton effects (Figure 6) of **2** and **5** implied that **2** had the 85,95,12*R*,18*S* absolute configuration, which was also confirmed by TD-DFT ECD calculation of (8*S*,9*S*,12*R*,18*S*)-**2**.

Cyclotryprostatin G (**3**) was also isolated as an analog of **5**, deduced from the similar NMR data of them. Compound **3** was the demethylation derivative of **5** at C-6, which was verified by <sup>1</sup>H-<sup>1</sup>H COSY (H-4/H-5/H-6/H-7) and HMBC (H-6 and C-4, and H-6 and C-7a) experiments for **3**. The stereochemistry of **3** was suggested to be *85,95,125,185* by comparing the NOESY and ECD spectra of **3** to those of **5** (Figure 6).

It is deduced that, the newborn OH-12 in **1** and **3** may be catalyzed by the multi-functional P450 *ctpP450-3* (Figure 1). And the new formed OMe-8 in **23** may be introduced by the methyltransferase encoded gene *ctpOMT* or other genes outside the reported boundary, or they were formed during our isolation processes.



Figure 6. Experimental ECD of 2, 3 and 5, and calculated ECD of (85,95,12R,185)-2.

#### 2.3. Biological Activities Screening

It has been reported that many indole-diketopiperazine derivatives displayed a wide range of biological effects, such as cytotoxic [16,28] and antibacterial activities [29]. In the present study, all of the isolated compounds (1–8) were tested for their cytotoxic and antibacterial activities. Among them, 1 and 4 displayed selective cytotoxicities against HL-60 cell line with the IC<sub>50</sub> values of 6.0 and 7.9  $\mu$ M, respectively. Whereas compounds 2, 3, and 5 exhibited activities against MCF-7 cell line with the IC<sub>50</sub> values of 7.6, 10.8, and 5.1  $\mu$ M, respectively. However, all of the metabolites appeared to be inactive in antibacterial and antifungal assays (MIC > 25  $\mu$ M).

# 3. Materials and Methods

#### 3.1. Instrumentation

Optical rotations were measured on a JASCO P-1020 digital polarimeter (Jasco Corp., Tokyo, Japan). UV spectra were recorded on a Thermo Scientific Multiskan GO microplate spectrophotometer (Thermo Scientific Co., Waltham, MA, USA) in MeOH. ECD spectra were recorded on a JASCO J-815 circular dichroism spectrometer (JASCO Electric Co., Ltd., Tokyo, Japan). IR spectra were determined on a Nicolet-Nexus-470 spectrometer (Thermo Electron Co., Madison, WI, USA) using KBr pellets. 1D and 2D NMR spectroscopic data (TMS as an internal standard) were acquired on Agilent DD2-500 (JEOL, Tokyo, Japan) and Bruker AV-600 NMR spectrometer (Bruker BioSpin GmbH Co., Rheinstetten, Germany). ESIMS and HRESIMS spectra were obtained from a Micromass Q-TOF spectrometer (Waters Corp., Manchester, UK), and a Thermo Scientific LTQ Orbitrap XL spectrometer (Thermo Fisher Scientific, Bremen, Germany). Semipreparative HPLC was performed on a Shimadzu LC-20 AT system using a RP-C18 (Waters, 5  $\mu$ m, 10 × 250 mm) column. 3.2. Genome Sequencing and Bioinformatics Analysis

The fungus *Penicillium brasilianum* HBU-136 was cultured in PDB medium for 30 h at 25 °C, then high-quality genomic DNA was isolated from cultured cells using Fungal DNA Kit (OMEGA Bio-Tek, E.Z.N.A.<sup>®</sup>, (Norcross, GA, USA). All the purified DNA was used to construct 350 bp DNA library with DNA library preparation kit (Illumina, CA, USA), and was sequenced on HiSeq X10 instrument, employing paired-end 150 base reads (Mega genomics, CN). *De novo* assembling of the draft genome was performed by using the software ABySS 2.15 (Vancouver, NA, Canada) and SPAdes 3.11 (St Petersburg, EU, Russia), with multiple k-mers specified as "-k 33,55,77,99,117" and coverage fold of 160X PE clean reads. All the assemble scaffold sequences were tested by quality assessment tool QUAST 4.0, the last draft genome sequence was filtered by over 500 bp and was determined via the largest N50 number. Gene functions were proposed by the online BLAST program (http://blast.ncbi.nlm.nih.gov/) and Conserved Domain Database (CDD) at the NCBI server (https://www.ncbi.nlm.nih.gov/). This Whole Genome Shotgun project of the fungus HBU-136 has been deposited at DDBJ/ENA/GenBank under the accession SSWP00000000. The version described in this paper is version SSWP01000000.

## 3.3. Fungal Material

The fungus *Penicillium brasilianum* HBU-136 was collected from the Bohai Sea (Huanghua, Hebei Province, China, June 2016). The strain with the NCBI GenBank accession number MH377073 was deposited in the College of Pharmaceutical Sciences, Hebei University, China.

## 3.4. Fermentation and Purification

The first fermentation was carried out for the fungus HBU-136 using rice medium (80 mL water and 80 g rice in 1 L Erlenmeyer flasks, 40 flasks) at 28 °C for 45 days. After cultivation, the fermented rice substrate was extracted with the mixture of CH<sub>2</sub>Cl<sub>2</sub>/MeOH (1:1, 500 mL for each flask) for five times and EtOAc for two times successively. The two extract solvents were combined as their TLC profiles were almost the same. The combined extract was concentrated to give a residue (30 g), which was further subjected to silica gel column chromatography (CC), eluting with an EtOAc–petroleum ether stepped gradient elution (0%, 20%, 40%, 60%, 80%, 100%), to offer six fractions Fr.1–Fr.6. Fr.5 was separated by twice Sephadex LH-20 CC [petroleum ether–MeOH–CHCl<sub>3</sub> (v/v, 2:1:1)] to afford four subfractions Fr.5-1–Fr.5-4. Then, Fr.5-3 was fractionated by Sephadex LH-20 CC [MeOH–CHCl<sub>3</sub> (v/v, 1:1)] and silica gel CC (stepped gradient, EtOAc–petroleum ether), and further purified by semipreparative HPLC (MeOH–H<sub>2</sub>O, 50:50, 2 mL/min) to afford 4 (3.1 mg) and 5 (6.7 mg).

The second large-scale fermentation (100 flasks) for the fungus HBU-136 was performed for 45 days in a modified rice medium (including 80 mL water, 80 g rice, and 0.8 g MgCl<sub>2</sub> in 1 L Erlenmeyer flask). The extract was obtained and purified by the same approaches as first fermentation. Compounds **1** (1.8 mg), **4** (5.2 mg), **5** (4.8 mg), **7** (1.5 mg), and **8** (6.2 mg) were given from fraction Fr.5. Compounds **2** (4.0 mg), **3** (3.2 mg), and **6** (4.3 mg) were isolated from fraction Fr.6.

Spirotryprostatin G (1):Yellow amorphous powder;  $[\alpha]_D^{20} = + 44.6$  (*c* 1.00, MeOH); UV (MeOH)  $\lambda_{max}$  (log  $\varepsilon$ ) 204 (4.05), 220 (4.02), 249 (3.96), 287 (3.77), 387 (3.17) nm; CD (0.75 mM, MeOH)  $\lambda_{max}$  ( $\Delta \varepsilon$ ) 201 (+20.7), 219 (-4.3), 239 (-3.4), 252 (-10.9), 327 (+0.16) nm; IR (KBr)  $v_{max}$  3447, 2360 and 1647 cm<sup>-1</sup>; <sup>1</sup>H and <sup>13</sup>C NMR, see Table 1; Table 2; HRESIMS *m*/*z* 444.1757 [M + H]<sup>+</sup> (calcd for C<sub>22</sub>H<sub>26</sub>N<sub>3</sub>O<sub>7</sub>, 444.1765 [M + Na]<sup>+</sup>), 466.1578 [M + Na]<sup>+</sup> (calcd for C<sub>22</sub>H<sub>25</sub>N<sub>3</sub>O<sub>7</sub>Na, 466.1585 [M + Na]<sup>+</sup>).

Cyclotryprostatin F (2):Yellow amorphous powder;  $[\alpha]_D^{20} = +108$  (*c* 1.00, MeOH); UV (MeOH)  $\lambda_{max}$  (log  $\varepsilon$ ) 220 (4.30), 290 (3.70) nm; CD (0.75 mM, MeOH)  $\lambda_{max}$  ( $\Delta \varepsilon$ ) 213 (+13.8), 230 (+8.6), 266 (+5.6), 319 (-0.01) nm; <sup>1</sup>H and <sup>13</sup>C NMR, see Table 1; Table 2; HRESIMS *m*/*z* 440.1837 [M – H]<sup>-</sup> (calcd for C<sub>23</sub>H<sub>26</sub>N<sub>3</sub>O<sub>6</sub>, 440.1827 [M – H]<sup>-</sup>).

Cyclotryprostatin G (**3**):Yellow amorphous powder;  $[\alpha]_D^{20} = +101$  (*c* 0.95, MeOH); UV (MeOH)  $\lambda_{max}$  (log  $\varepsilon$ ) 223 (4.45), 272 (3.93) nm; CD (1.05 mM, MeOH)  $\lambda_{max}$  ( $\Delta \varepsilon$ ) 195 (-8.1), 208 (+11.2), 228 (+5.7), 258 (+3.2), 298 (+0.2) nm; <sup>1</sup>H and <sup>13</sup>C NMR, see Table 1; Table 2; HRESIMS *m*/*z* 418.1737 [M + Na]<sup>+</sup> (calcd for C<sub>22</sub>H<sub>25</sub>N<sub>3</sub>O<sub>4</sub>Na, 418.1737 [M + Na]<sup>+</sup>).

Cytotoxic Assay. The cytotoxic activity of compounds **1–8** were evaluated in vitro by the MTT method [30], with the concentration of 10  $\mu$ M. Three human tumor cell lines were used, including human promyelocytic leukemia cell line (HL-60), human colorectal cancer cell line (HCT-116), and human breast cancer cell line (MCF-7), with cisplatinum (DDP) as a positive control.

Antibacterial Assay. The antibacterial activity of the isolated compounds was determined using the conventional broth dilution assay [31] with ciprofloxacin (CIP) as a positive control. Sixteen pathogenic bacterial strains were used, including *Bacillus megaterium*, *Bacillus subtilis*, *Escherichia coli*, *Bacillus anthraci*, *Bacillus cereus*, *Bacterium paratyphosum B*, *Enterobacter aerogenes*, *Micrococcus lysodeikticus*, *Micrococcus luteus*, *Proteusbacillm vulgaris*, *Shigella dysenteriae*, *Psmdomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella typhi*, *Vibrio anguillarum*, and *Vibrio parahemolyticus*. The concentration of compounds **1–8** was 25 µM.

## 3.6. ECD Spectrum Measurement and Calculation

The tested compounds **1–5** were dissolved in chromatographic methanol at the concentration of 0.5 mg/mL and transferred 300  $\mu$ L to quartz cuvette (1cm for width) for tested. First tested the ECD spectrum of methanol as the blank control, then tested for compounds **1–5**. The wavelength was set at 190–400 nm with the band width of 1 nm, and Nitrogen should be used throughout the experiment at the flow rate of 3 mL/min. The ECD spectrum was calculated by the TDDFT methodology with a larger basis set at the B3LYP/6-311++G(2d,p) level and simulated using SpecDis 1.62 [32] according to Boltzmann distributions.

## 4. Conclusions

In summary, three new indole-diketopiperazine derivatives (1–3) showed selective cytotoxicities have been obtained from the marine-derived fungus *P. brasilianum* HBU-136 with the aid of genomic analysis. ECD quantum chemistry calculations were carried out to assign the absolute configurations of the new compounds 1–3. The present research indicated that it is a powerful approach to discover new natural products from marine-derived fungi by the combination of chemical and bioinformatics analyses.

**Supplementary Materials:** The following are available online at www.mdpi.com/1660-3397/17/9/514/s1, NMR and MS data of **1–8**; Supplementary tables and figures for BGC analyses; tables for biological assay.

**Author Contributions:** Y.-H.Z. contribute to fermentation, extraction, isolation, and manuscript preparation; C.G. and X.-W.Z. contribute to genome sequencing; H.-J.Z. contribute to quantum chemistry calculation; C.-L.S. contribute to bioactivities test; F.C. and C.-Y.W. were the project leaders organizing and guiding the experiments and manuscript writing.

**Funding:** This work was supported by the National Natural Science Foundation of China (Nos. 41606174; 81673350; 41830535), the Marine S&T Fund of Shandong Province for Pilot National Laboratory for Marine Science and Technology (Qingdao, China) (No. 2018SDKJ0406-5), the Fundamental Research Funds for the Central Universities of China (No. 201762017), Shandong Provincial Natural Science Foundation (No. ZR2017BD009), and the High Performance Computer Center of Hebei University, and the Taishan Scholars Program, China.

Conflicts of Interest:. The authors declare no conflict of interest.

## References

- Carroll, A.R.; Copp, B.R.; Davis, R.A.; Keyzers, R.A.; Prinsep, M.R. Marine natural products. *Nat. Prod. Rep.* 2019, 36, 122–173.
- 2. Piel, J.; Hoang, K.; Moore, B.S. Natural metabolic diversity encoded by the enterocin biosynthesis gene cluster. *J. Am. Chem. Soc.* **2000**, *122*, 5415–5416.

- Xu, W.; Dhingra, S.; Chooi, Y.H.; Calvo, A.M.; Lin, H.C.; Tang, Y. The fumagillin biosynthetic gene cluster in *Aspergillus fumigatus* encodes a cryptic terpene cyclase involved in the formation of β-trans-bergamotene. *J. Am. Chem. Soc.* 2013, *135*, 4616–4619.
- 4. Wang, J.F.; He, W.J.; Huang, X.L.; Tian, X.P.; Liao, S.R.; Yang, B.; Wang, F.Z.; Zhou, X.J.; Liu, Y.H. Antifungal new oxepine-containing alkaloids and xanthones from the deep-sea-derived fungus *Aspergillus versicolor* SCSIO 05879. *J. Agric. Food Chem.* **2016**, *64*, 2910–2916.
- Song, F.H.; Liu, X.R.; Guo, H.; Ren, B.; Chen, C.X.; Piggott, A.M.; Yu, K.; Gao, H.; Wang, Q.; Liu, M.; et al. Brevianamides with antitubercular potential from a marine-derived isolate of *Aspergillus versicolor*. Org. Lett. 2012, 14, 4770–4773.
- 6. Wang, M.Z.; Sun, M.W.; Hao, H.L.; Lu, C.H. Avertoxins A–D, Prenyl asteltoxin derivatives from *Aspergillus versicolor* Y10, an endophytic fungus of *Huperzia serrate*. J. Nat. Prod. **2015**, 78, 3067–3070.
- Zhou, M.; Miao, M.M.; Du, G.; Li, X.N.; Shang, S.Z.; Zhao, W.; Liu, Z.H.; Yang, G.Y.; Che, C.T.; Hu, Q.F.; et al. Aspergillines A–E, highly oxygenated hexacyclic indole–tetrahydrofuran–tetramic acid derivatives from *Aspergillus versicolor*. Org. Lett. 2014, 16, 5016–5019.
- 8. Son, S.; Hong, Y.S.; Jang, M.; Heo, K.T.; Lee, B.; Jang, J.P.; Kim, J.W.; Ryoo, I.J.; Kim, W.G.; Ko, S.K.; et al. Genomics-driven discovery of chlorinated cyclic hexapeptides ulleungmycins A and B from a Streptomyces species. *J. Nat. Prod.* **2017**, *80*, 3025–3031.
- Hu, Y.Y.; Wang, M.; Wu, C.Y.; Tan, Y.; Li, J.; Hao, X.M.; Duan, Y.B.; Guan, Y.; Shang, X.Y.; Wang, Y.G.; et al. Identification and proposed relative and absolute configurations of niphimycins C–E from the marine-derived *Streptomyces* sp. IMB7-145 by genomic analysis. *J. Nat. Prod.* 2018, *81*, 178–187.
- Xu, X.K.; Zhou, H.B.; Liu, Y.; Liu, X.T.; Fu, J.; Li, A.Y.; Li, Y.Z.; Shen, Y.M.; Bian, X.Y.; Zhang, Y.M. Heterologous Expression Guides Identification of the Biosynthetic Gene Cluster of Chuangxinmycin, an Indole Alkaloid Antibiotic. *J. Nat. Prod.* 2018, *81*, 1060–1064.
- 11. Zhu, A.; Zhang, X.W.; Zhang, M.; Li, W.; Ma, Z.Y.; Zhu, H.J.; Cao, F. Aspergixanthones I–K, new anti-Vibrio prenylxanthones from the marine-derived fungus *Aspergillus sp.* ZA-01. *Mar. Drugs* **2018**, *16*, 312–319.
- Yang, J.K.; Zhang, B.; Gao, T.; Yang, M.Y.; Zhao, G.Z.; Zhu, H.J.; Cao, F. A pair of enantiomeric 5-oxabicyclic [4.3.0] lactam derivatives and one new polyketide from the marine-derived fungus *Penicillium griseofulvum*. *Nat. Prod. Res.* 2018, *32*, 2366–2369.
- 13. Cao, F.; Meng, Z.H.; Mu, X.; Yue, Y.F.; Zhu, H.J. Absolute configuration of bioactive azaphilones from the marine-derived fungus *Pleosporales* sp. CF09-1. *J. Nat. Prod.* **2019**, *82*, 386–392.
- 14. Zhu, A.; Yang, M.Y.; Zhang, Y.H.; Shao, C.L.; Wang, C.Y.; Hu, L.D.; Cao, F.; Zhu, H.J. Absolute configurations of 14, 15-hydroxylated prenylxanthones from a marine-derived *Aspergillus* sp. fungus by chiroptical methods. *Sci. Rep.* **2018**, *8*, 10621–10630.
- Wang, F.Z.; Fang, Y.C.; Zhu, T.J.; Zhang, M.; Lin, A.Q.; Gu, Q.Q.; Zhu, W.M. Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*. *Tetrahedron* 2008, 64, 7986–7991.
- 16. Cui, C.B.; Kakeya, H.; Osada. H. Novel mammalian cell cycle inhibitors, spirotryprostatins A and B, produced by *Aspergillus fumigatus*, which inhibit mammalian cell cycle at G2/M phase. *Tetrahedron* **1996**, *52*, 12651–12666.
- Kato, N.; Suzuki, H.; Takagi, H.; Asami, Y.; Kakeya, H.; Uramoto, M.; Usui, T.; Takahashi, S.; Sugimoto, Y.; Osada, H. Identification of cytochrome P450s required for fumitremorgin biosynthesis in *Aspergillus fumigatus. ChemBioChem.* 2010, *10*, 920–928.18.
- 18. Maiya, S.; Grundmann, A.; Li, S.M.; Turner, G. The fumitremorgin gene cluster of *Aspergillus fumigatus*: Identification of a gene encoding brevianamide F synthetase. *ChemBioChem.* **2010**, *7*, 1062–1069.
- 19. Grundmann, A.; Kuznetsova, T.; Afiyatullov, S.S.; Li, S.M. FtmPT2, an *N*-Prenyltransferase from *Aspergillus fumigatus*, catalyses the last step in the biosynthesis of fumitremorgin B. *ChemBioChem* **2008**, *9*, 2059–2063.
- Tsunematsu, Y.; Ishiuchi, K.; Hotta, K.; Watanabe, K. Yeast-based genome mining, production and mechanistic studies of the biosynthesis of fungal polyketide and peptide natural products. *Nat. Prod. Rep.* 2013, *30*, 1139–1149.
- 21. Gu, B.; He, S.; Yan, X.; Zhang, L. Tentative biosynthetic pathways of some microbial diketopiperazines. *Appl. Microbiol. Biot.* **2013**, *97*, 8439–8453.
- Tsunematsu, Y.; Ishikawa, N.; Wakana, D.; Goda, Y.; Noguchi, H.; Moriya, H.; Hotta, K.; Watanabe, K. Distinct mechanisms for spiro-carbon formation reveal biosynthetic pathway crosstalk. *Nat. Chem. Biol.* 2013, 9, 818–827.

- 23. Zhang, X.W.; Li, S.Y. Expansion of chemical space for natural products by uncommon P450 reactions. *Nat. Prod. Rep.* **2017**, *34*, 1061–1089.
- 24. Wang, W.J.; Li, D.Y.; Li, Y.C.; Hua, H.M.; Ma, E.L.; Li, Z.L. Caryophyllene sesquiterpenes from the marinederived fungus *Ascotricha* sp. ZJ-M-5 by the one strain–many compounds strategy. *J. Nat. Prod.* **2014**, *77*, 1367–1371.
- 25. Wang, Y.; Li, Z.L.; Bai, J.; Zhang, L.M.; Wu, X.; Zhang, L.; Pei, Y.H.; Jing, Y.K.; Hua, H.M. 2,5-Diketopiperazines from the Marine-Derived Fungus *Aspergillus fumigatus* YK-7. *Chem. Biodivers.* **2012**, *9*, 385–393.
- Yu, L.Y.; Ding, W.J.; Wang, Q.Q.; Ma, Z.J.; Xu, X.W.; Zhao, X.F.; Chen, Z. Induction of cryptic bioactive 2, 5-diketopiperazines in fungus *Penicillium* sp. DT-F29 by microbial co-culture. *Tetrahedron* 2017, 73, 907–914.
- 27. Xue, H.; Lu, C.H.; Liang, L.Y.; Shen, Y.M. Secondary Metabolites of *Aspergillus* sp. CM9a, an Endophytic Fungus of *Cephalotaxus mannii*. *Rec. Nat. Prod.* **2012**, *6*, 28–34.
- 28. Wang, F.Q.; Tong, Q.Y.; Ma, H.R.; Xu, H.F.; Hu, S.; Ma, W.; Xue, Y.B.; Liu, J.J.; Wang, J.P.; Song, H.P.; et al. Indole diketopiperazines from endophytic *Chaetomium* sp. 88194 induce breast cancer cell apoptotic death. *Sci. Rep.* **2015**, *5*, 9294.
- 29. Meng, L.H.; Du, F.Y.; Li, X.M.; Pedpradab, P.; Xu, G.M.; Wang, B.G. Rubrumazines A–C, indolediketopiperazines of the isoechinulin class from *Eurotium rubrum* MA-150, a fungus obtained from marine mangrove-derived rhizospheric soil. *J. Nat. Prod.* **2015**, *78*, 909–913.
- Scudiere, D.A.; Shoemaker, R.H.; Paull, K.D.; Monks, A.; Tierney, S.; Nofziger, T.H.; Currens, M.J.; Seniff, D.; Boyd, M.R. Evaluation of a soluble tetrazolium/formazan assay for cell growth and drug sensitivity in culture using human and other tumor cell lines. *Cancer Res.* 1988, 48, 4827–4833.
- 31. Appendino, G.; Gibbons, S.; Giana, A.; Pagani, A.; Grassi, G.; Stavri, M.; Smith, E.; Rahman, M.M. Antibacterial cannabinoids from Cannabis sativa: A structure–Activity study. *J. Nat. Prod.* **2008**, *71*, 1427–1430.
- 32. Bruhn, T.; Schaumlöffel, A.; Hemberger, Y.; Bringmann, G. SpecDis: Quantifying the comparison of calculated and experimental electronic circular dichroism spectra. *Chirality* **2013**, *25*, 243–249.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).