

SUPPORTING INFORMATION

Induction of Apoptotic Cell Death in Human Leukemia U937 Cells by C18 Hydroxy Unsaturated Fatty Acid Isolated from Red Alga *Tricleocarpa jejuensis*

Shijiao Zha ^{1,†}, Mikinori Ueno ^{2,†}, Yan Liang ¹, Seiji Okada ², Tatsuya Oda ¹ and Fumito Ishibashi ^{1,*}

¹Graduate School of Fisheries and Environmental Sciences, Nagasaki University, 1-14 Bunkyo-machi, Nagasaki 852-8521, Japan; bb53418001@ms.nagasaki-u.ac.jp (S.Z.); bb53419804@ms.nagasaki-u.ac.jp (Y.L.); t-oda@nagasaki-u.ac.jp (T.O.)

²Joint Research Center for Human Retrovirus Infection, Division of Hematopoiesis, Kumamoto University, 2-2-1 Honjo, Chuo-ku, Kumamoto 860-0811, Japan; mikiueno@kumamoto-u.ac.jp (M.U.); okadas@kumamoto-u.ac.jp (S.O.)

*Correspondence: fumito@nagasaki-u.ac.jp; Tel.: +81-95-819-2833

[†] Both authors have contributed equally to this work

Contents

Flow cytometry data of Figure 6


S1~S3

Flow cytometry data of Figure 8

S4


Flow cytometry data of Figure 6

First time




Flow cytometry data of Figure 6

Second time





Flow cytometry data of Figure 6

Third time



Flow cytometry data of Figure 8



Inhibitor: Z-Asp-CH₂-DCB