Supplementary material for

Geocoding error, spatial uncertainty, and implications for exposure assessment

and environmental epidemiology

Ellen J. Kinnee^{1*}, Sheila Tripathy², Leah Schinasi³, Jessie L. C. Shmool⁴, Perry E. Sheffield⁵, Fernando Holguin⁶, and Jane E. Clougherty⁷

- ¹ University Center for Social and Urban Research, University of Pittsburgh, Pittsburgh, Pennsylvania, United States ; ejk40@pitt.edu
- ² Drexel University Dornsife School of Public Health, Department of Environmental and Occupational Health, Philadelphia, Pennsylvania, United States; sheila.tripathy@gmail.com
- ³ Drexel University Dornsife School of Public Health, Department of Environmental and Occupational Health, and Drexel University Urban Health Collaborative (UHC), Philadelphia, Pennsylvania, United States; lhs36@drexel.edu
- ⁴ University of Pittsburgh Graduate School of Public Health, Department of Environmental and Occupational Health, Pittsburgh, Pennsylvania, United States; jlcshmool@gmail.com
- ⁵ Environmental Medicine and Public Health and Pediatrics, Icahn School of Medicine at Mount Sinai, New York, New York United States; perry.sheffield@mssm.edu
- ⁶ Department of Medicine, University of Colorado School of Medicine, Aurora, Colorado, United States
- ⁷ Drexel University Dornsife School of Public Health, Department of Environmental and Occupational Health, Philadelphia, Pennsylvania, United States; jec373@drexel.edu
- * Correspondence: ejk40@pitt.edu; Tel.: +1 412-385-5105

Figure S1: Street segment geocodes using various side offsets, and distance error. The default 20-foot offset (orange) is shown with calculated distance error from the reference (Address point). Alternative side offsets are shown [30 feet (blue), 40 feet (yellow), and 50 feet (purple)].




Figure S2: Buffer-based exposure assignment for NO_2 , using each of the three geocoding methods and the NYCCAS NO_2 surface.




Figure S3: Repeat addresses in the dataset are shown in blue (1-5 obs/address), yellow (5-10 obs/ address), or red (>10 obs/address).




Table S1: Missingness rates: Overall missingness rate for all addresses (n=21,183), and by ZIP Code (*n= 157 ZIP Code areas*)

Method (locator)	Overall missingness rate	Average missingness by ZIP	SD (by ZIP)	Median (by ZIP)	Min (by ZIP)	Max (by ZIP)
Address Point	9.6	9.9	8.4	8.0	0	70.0
Street Segment	6.1	9.9	6.0	6.0	0	42.0
Parcel Centroid	38.1	35.0	23.5	28.0	0	93.0

Table S2: Frequency (*percentage*) with directional error in each of four directional quadrants, by method

Method	North	East	South	West
Street Segment	5,040 (23.8%)	3,853 (18.2%)	5,944 (28.1%)	6,346 (29.9%)
Parcel Centroid	4,968 (23.5%)	5,987 (28.3%)	4,836 (22.8%)	5,392 (25.5%)

Table S3: Change in Census Tract, Block Group, and Block assignment using Street Segment or Parcel Centroid geocodes, compared to Address points (n = 21,183)

Geocoding method	# Tracts changed (%)	<pre># Block Groups changed (%)</pre>	# Blocks changed
			(%)
Street Segment	63 (2.9%)	188 (1.2%)	301 (0.78%)
Parcel Centroid	13 (0.6%)	119 (0.77%)	111 (0.29%)
Total in domain	2,167	15,464	38,794
Mean area of polygon (m ²)	362,888	129,780	20,186

Figure S4: Color-coded spatial uncertainty (distance error), overlaid with above/ below-median number of addresses per ZIP code. There is no consistent pattern in distance error by address density.

