

**Supplementary Table 5.** Changes in symptom prevalence between day 0 and day 7 in 508 patients according to care setting.

Symptoms	Setting	Prevalence, N (%)		p value <sup>a</sup>
		Day 0	Day 7	
Asthenia	Home	348 (93.5)	352 (94.6)	NS
	Hospice	85 (92.4)	81 (88.0)	NS
	Hospital	39 (88.6)	41 (93.2)	NS
	p value <sup>b</sup>	NS	NS	
Poor well-being	Home	302 (81.2)	302 (81.2)	NS
	Hospice	63 (68.5)	66 (71.7)	NS
	Hospital	31 (70.5)	31 (70.5)	NS
	p value <sup>b</sup>	<0.001	<0.01	
Lack of appetite	Home	280 (75.3)	280 (75.3)	NS
	Hospice	68 (73.9)	56 (60.9)	<0.05
	Hospital	26 (59.1)	26 (59.1)	NS
	p value <sup>b</sup>	NS	<0.01	
Drowsiness	Home	264 (71.0)	263 (70.7)	NS
	Hospice	64 (69.6)	63 (68.5)	NS
	Hospital	29 (65.9)	25 (56.8)	NS
	p value <sup>b</sup>	NS	<0.05	
Pain	Home	267 (71.8)	252 (67.7)	NS
	Hospice	45 (48.9)	42 (45.7)	NS
	Hospital	31 (70.5)	25 (56.8)	NS
	p value <sup>b</sup>	<0.001	<0.001	
Depression	Home	224 (60.2)	234 (62.9)	NS
	Hospice	67 (72.8)	63 (68.5)	NS
	Hospital	13 (29.5)	22 (50.0)	<0.05
	p value <sup>b</sup>	<0.001	NS	
Anxiety	Home	209 (56.2)	230 (61.8)	<0.05
	Hospice	56 (60.9)	48 (52.2)	NS
	Hospital	25 (56.8)	23 (52.3)	NS
	p value <sup>b</sup>	NS	NS	
Breathlessness	Home	182 (48.9)	151 (40.6)	<0.001
	Hospice	36 (39.1)	29 (31.5)	NS
	Hospital	15 (34.1)	12 (27.3)	NS
	p value <sup>b</sup>	NS	NS	
Nausea	Home	187 (50.3)	168 (45.2)	<0.05
	Hospice	30 (32.6)	17 (18.5)	<0.01
	Hospital	8 (18.2)	7 (15.9)	NS
	p value <sup>b</sup>	<0.001	<0.001	

NS, not statistically significant; SD, standard deviation. <sup>a</sup>p value for difference between day 7 and day 0; <sup>b</sup>p value for difference between care settings.