



Article

# The Effect of Contextual Variables on Match Performance across Different Playing Positions in Professional Portuguese Soccer Players

Joel Barrera <sup>1,\*</sup>, Hugo Sarmento <sup>1,2,\*</sup>, Filipe Manuel Clemente <sup>3,4</sup>, Adam Field <sup>5</sup> and António J. Figueiredo <sup>1,2</sup>

<sup>1</sup> Faculty of Sport Sciences and Physical Education, University of Coimbra, 3000-248 Coimbra, Portugal; afigueiredo@fcdef.uc.pt

<sup>2</sup> Research Unit for Sport and Physical Activity (CIDAF), 3000-248 Coimbra, Portugal

<sup>3</sup> Escola Superior Desporto e Lazer, Instituto Politécnico de Viana do Castelo, Rua Escola Industrial e Comercial de Nun'Álvares, 4900-347 Viana do Castelo, Portugal; filipe.clemente5@gmail.com

<sup>4</sup> Instituto de Telecomunicações, Delegação da Covilhã, 1049-001 Lisboa, Portugal

<sup>5</sup> School of Human and Health Sciences, University of Huddersfield, Huddersfield HD1 3DH, UK; Adam.field@hud.ac.uk

\* Correspondence: jibarrera@outlook.es (J.B.); hg.sarmiento@gmail.com (H.S.)

**Abstract:** This study investigated the position-specific physical demands of professional Portuguese players. The effects of situational variables on the physical performance demands were also analysed (match location, match half and match result). Match performance observations were collected using Global Navigation Satellite System devices across 11 matches during a competitive season (2019–2020). Data were analysed according to five playing positions: goalkeepers ( $n = 11$ ), central defenders ( $n = 42$ ), wide defenders ( $n = 31$ ), central midfielders ( $n = 34$ ), open attackers ( $n = 28$ ), and centre forwards ( $n = 14$ ). Central midfield players completed the greatest total distance ( $10,787 \pm 1536$  m), while central defenders covered the least distance ( $9272 \pm 455$ ;  $p < 0.001$ ). Open attackers covered the greatest high and very-high-speed distance ( $1504 \pm 363$  m), number of high-speed decelerations per match ( $11 \pm 4$ ) and were the fastest players ( $30.6 \pm 1.5$  km/h), along with center forwards ( $30.6 \pm 2.0$  km/h), versus all other positions ( $p < 0.05$ ). Greater distances were performed in teams that were winning ( $9978 \pm 1963$  m) or drawing ( $10,395 \pm 875$  m) versus losing ( $9415 \pm 2050$ )  $p = 0.036$  and  $p = 0.006$ , respectively. Increases in distance covered at walking speeds were observed during the 2nd half ( $1574 \pm 179$  m) compared with the 1st half ( $1483 \pm 176$ ;  $p < 0.003$ ). A higher number of decelerations across all speeds were performed in the 1st half ( $144 \pm 39$ ) versus the 2nd half ( $135 \pm 37$ ). The distance covered in home matches ( $10,206 \pm 1926$  m) far exceeded away matches ( $9471 \pm 1932$  m;  $p < 0.001$ ). The number of faster accelerations were higher in away ( $7 \pm 5$ ) versus home matches ( $6 \pm 4$ ;  $p < 0.049$ ). The data demonstrate the different physical demands of each playing position and suggest that situational variables influence physical performance. These findings suggest position-specific physical training is required to condition players for the bespoke demands of each playing position.

**Keywords:** football; monitoring; performance; professional team



**Citation:** Barrera, J.; Sarmento, H.; Clemente, F.M.; Field, A.; Figueiredo, A.J. The Effect of Contextual Variables on Match Performance across Different Playing Positions in Professional Portuguese Soccer Players. *Int. J. Environ. Res. Public Health* **2021**, *18*, 5175. <https://doi.org/10.3390/ijerph18105175>

Academic Editor: Dustin Russel Slivka

Received: 9 April 2021

Accepted: 10 May 2021

Published: 13 May 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

The physical performance demands of soccer match-play have been researched extensively in English Premier League soccer players [1–4], and Australian, Italian [5] and Spanish league players [6]. Classic and contemporary match observations suggest professional soccer players cover between 9 and 14 km during match-play [7–9]. Studies primarily of English players also indicate that during competitive encounters, elite soccer players cover between 0.7–3.9 km high-speed distance [8,10] and 0.2–0.6 km sprint distance [11,12], while Spanish players in friendly matches cover smaller distances in these same variables [13]. English players are also reported to perform ~656 accelerations and

~612 decelerations during match-play [14], and Spanish players ~581 accelerations [13]. It is estimated that players spend the majority (80–90%) of matches performing low and medium intensity activities, whereas the decisive moments of a match largely require high-speed and sprint actions [15,16]. However, whilst the match demands have been clearly outlined in professional English players, there remain few match-play studies evaluating the demands of professional Portuguese players. Therefore, elucidating the match demands of professional Portuguese players appears warranted.

During competitive English Premier League matches, there appear to be large differences in the physical running output of different playing positions [17]. The greatest distance seems to be covered by midfield players (11.5 km) [15,18], whilst forwards and defenders cover lesser distances during match-play (10–10.5 km) [15]. A major limitation of the studies discriminating between playing positions is that small samples are used; thus, findings are often difficult to accurately differentiate across varying positional roles. Therefore, while the running demands during matches appear to differ, there are few large-scale studies of professional players elucidating the match demands between different positional roles. Assessing the difference between positional demands may help tailor individual training processes towards improving training drills for the needs of individual playing positions.

Given that soccer is associated with many complex logistical factors, it is plausible that situational variables influence match running profiles. It has also been identified that performance metrics can differ according to specific match factors [19,20]. One such contextual variable shown to influence physical performance is match half, with evidence suggesting that greater running distance is covered during the 1st half versus the 2nd half of matches [21,22]. Match location impacts technical performance [23], although few large-scale studies have assessed whether competing in home or away matches can influence the physical running metrics of soccer players. Another factor that has been investigated for its impact on performance is match status (Lago, 2009), which refers to whether a team wins, draws or loses a competitive match. Spanish La Liga soccer players are reported to cover greater distances when losing versus matches that are being drawn or won [24]. These findings are consistent with a separate study also evaluating the running profiles of Spanish La Liga players, which demonstrates that lesser high-speed running is performed when winning versus losing matches (Lago et al., 2010). Further research is required that develops an understanding of the influence of situational variables on soccer-specific running metrics across different leagues and countries. Considering the different cultures across European countries, it is logical to assume that the contextual variables influencing performance in some leagues may not apply to other countries. Therefore, assessing the influence that situational variables have on physical performance in professional Portuguese players will provide novel interpretations.

The aim of this study was to evaluate the physical performance of professional Portuguese soccer players across different positional roles. The interactive effects between match half, location, and status on physical performance metrics across official Portuguese matches were also assessed.

## 2. Materials and Methods

### 2.1. Experimental Approach to the Problem

A quasi-experimental design was used to evaluate the activity profile of different positional roles in competitive professional soccer matches. Those who played the entire 90 min (including added time) during the 2019–2020 (August to May) season were included for analyses. The matches were played between 11:00 am and 20:30 pm, and all players refrained from strenuous activity for 24 h prior to each match. All matches were played on natural turf. Given the observational nature of the research, no intervention or attempts were made to influence players performance, nor was any feedback provided to players throughout the entirety of data collection.

## 2.2. Participants

A total of 25 professional soccer players (age:  $24.9 \pm 4.2$  years, stature:  $180.9 \pm 6.3$  cm and body mass:  $78.5 \pm 8.5$  kg) were monitored over 11 official matches in the Portuguese LigaPro (second tier). The players were classified into six different positions: goalkeepers (GK), wide defenders (WD), central defenders (CD), center midfielder (CM), open attackers (OA) and center forward (CF). The athletes who did not complete the entire duration of all 11 matches were excluded from analyses ( $n = 38$ ). Each player provided informed consent, as did the leadership of the Coimbra Academic Association club. This study was approved by the scientific council and the ethics committee of the University of Coimbra (CE/FCDEF-UC/00692021). At the beginning of the season each player was evaluated by the club's medical staff and underwent the Portuguese Football Federation's mandatory medical examinations.

## 2.3. Experimental Procedure

Stature was determined using a SECA wall stadiometer (range 10–230 cm, division 1 mm). Body mass was determined through a multifrequency bioimpedance device (In-Body770). Players voided their bowels and bladder before anthropometric characteristics were taken [25]. Assessments were conducted  $\geq 3$  h after the last meal with each player consuming approximately 500 mL of water  $\sim 2$  h before evaluation to standardise hydration. Matches were analysed both across halves (1st and 2nd) and 15-min segments (0–15, 16–30, 31–45, 46–60, 61–75, 76–90 min). Additional ('injury') time for the first and second halves were also separated for analyses. The data were filtered to exclude any activity before kickoff (warm-up), during and reactivation after half-time, and cool-down at the end of the match.

Match data were recorded using portable GNSS (SPI HPU, GPSports, Canberra, Australia) at a sampling frequency of 15 Hz. These devices have reported good inter-unit reliability (coefficient of variation [CV]: 1.9% for TD, 7.6% for speeds of 3.8–5.5 m/s/s and 12.1% for speeds  $\geq 5.5$  m/s/s). Previous models of the 15 Hz units have revealed a degree of error for TD (1.1%) and maximum speed running ( $<1\%$ ) [26,27]. At the end of each match, the data from the GNSS devices were extracted using the manufacturer's software (Team AMS, Canberra, Australia). Match activities were categorised as the following: standing, very low speed walking ( $0\text{--}5.9$  km·h<sup>-1</sup>), low speed walking ( $6\text{--}11.9$  km·h<sup>-1</sup>), low speed jogging ( $12\text{--}13.9$  km·h<sup>-1</sup>), medium speed running ( $14\text{--}17.9$  km·h<sup>-1</sup>), high speed running ( $18\text{--}23.9$  km·h<sup>-1</sup>) and sprinting ( $24$  km·h<sup>-1</sup>). The different speed thresholds from the 11 official matches are summarised in Table 1.

## 2.4. Statistical Analysis

Data were not normally distributed, and thus, non-parametric tests were used for analyses. Kruskal–Wallis was used to compare performance between the different positions and according to the match result. A Mann–Whitney U was performed for comparisons of match location and match half. Data were analysed using Statistical Package for the Social Sciences (IBM SPSS Statistics 24 for Windows; SPSS Inc., Chicago, IL, USA). Significance was accepted at  $p \leq 0.05$  prior to analyses.

**Table 1.** Speed thresholds for the reported match variables.

| Variables                 | Abbreviation       |                                      | Rank          |
|---------------------------|--------------------|--------------------------------------|---------------|
| Total distance            | TD m               |                                      | 0–≤24 km/h    |
| Total distance by parts   | TDP m              |                                      | 0–≤24 km/h    |
| Zone 1 distance           | ZD1 m              | standing, walking very low intensity | 0–5.9 km/h    |
| Zone 2 distance           | ZD2 m              | low intensity walking                | 6–11.9 km/h   |
| Zone 3 distance           | ZD3 m              | jog at low intensity                 | 12–13.9 km/h  |
| Zone 4 distance           | ZD4 m              | running at medium intensity          | 14–17.9 km/h  |
| Zone 5 distance           | ZD5 m              | running high speed                   | 18–23.9 km/h  |
| Zone 6 distance           | ZD6 m              | Sprint                               | ≤24 km/h      |
| Sum distances (1,2,3)     | ΣD m (1,2,3)       |                                      | 0–13.9 km/h   |
| Sum distances (4,5,6)     | ΣD m (4,5,6)       |                                      | 14–≤24 km/h   |
| Acceleration 1            | Acc1 m/s/s         |                                      | 1.0–1.9 m/s/s |
| Acceleration 2            | Acc2 m/s/s         |                                      | 2.0–2.9 m/s/s |
| Acceleration 3            | Acc3 m/s/s         |                                      | 3.0–4.0 m/s/s |
| Sum Accelerations (1,2,3) | ΣAcc (1,2,3) m/s/s |                                      | 1.0–4.0 m/s/s |
| Deceleration 1            | Des1 m/s/s         |                                      | 1.0–1.9 m/s/s |
| Deceleration 2            | Des2 m/s/s         |                                      | 2.0–2.9 m/s/s |
| Deceleration 3            | Des3 m/s/s         |                                      | 3.0–4.0 m/s/s |
| Sum Deceleration (1,2,3)  | ΣDes (1,2,3) m/s/s |                                      | 1.0–4.0 m/s/s |
| Maximum speed             | MS km/h            |                                      | ≤24 km/h      |

### 3. Results

Detailed results according to playing position are presented in Table 2 and between-group comparisons in Table 3. The lowest TD was performed by CD, while the highest TD was covered by CM ( $10.787 \pm 1.536$  m;  $p < 0.001$ ). The greatest high-speed distance in the sum of ZD4, 5 and 6 was performed by OA versus all other positions ( $p < 0.001$ ). The WD, MC and OA demonstrated significantly higher values for the sum of accelerations and decelerations compared with CD and CF ( $p < 0.001$ ).

It was identified that TD was higher for the tied and won matches ( $10.395 \pm 87$  m, and  $9.978 \pm 1.962$  m, respectively) versus lost matches ( $9.414 \pm 2.050$  m). Significant differences were found for TD between drawing and losing ( $p = 0.036$ ), and winning and losing ( $p = 0.006$ ) (Table 4).

No significant differences were observed for average TD covered between the 1st ( $4942 \pm 945$  m) and 2nd ( $4868 \pm 980$ ) halves ( $p = 0.432$ ). A decrease in the total number of decelerations in the 2nd half versus the 1st half was also identified ( $p = 0.036$ ) (Table 5). Greater TD ( $p = 0.0002$ ), TDP ( $p = 0.0008$ ) and total displacements at low intensity ( $p = 0.00003$ ) were observed in the home games in versus away matches. A higher number of Acc3 ( $p = 0.049$ ) was observed in the away matches. (Table 6).

The results revealed that the highest number of sprints were performed by OA ( $n = 5.39 \pm 2.6$ ), while the CD ( $n = 2.47 \pm 1.5$ ) showed the lowest values ( $p = 0.000$ ). Furthermore, the highest number of sprints based in accelerations (SBA) were performed in Q1 ( $n = 5.04 \pm 2.9$ ), while the lowest frequencies were identified during Q4 ( $n = 3.77 \pm 2.5$ ) (Table 7).

**Table 2.** Distances covered across each speed threshold according to each playing position (Data are reported as mean ± SD).

| Variables          | GK n = 11 |          | WD n = 31 |         | CD n = 42 |         | CM n = 34 |          | OA n = 28 |          | CF n = 14 |         | Differences Between Positions |        |
|--------------------|-----------|----------|-----------|---------|-----------|---------|-----------|----------|-----------|----------|-----------|---------|-------------------------------|--------|
|                    | Mean      | SD       | Mean      | SD      | Mean      | SD      | Mean      | SD       | Mean      | SD       | Mean      | SD      | H de Kruskal-Wallis           | Sig    |
| TD m               | 4237.9    | ± 2342.1 | 10,582.4  | ± 504.4 | 9272.5    | ± 455.7 | 10,787.9  | ± 1536.8 | 10,640.6  | ± 1011.4 | 9336.9    | ± 673.6 | 90.451                        | 0.0001 |
| TDP m              | 2139.2    | ± 1243.9 | 5284.0    | ± 292.7 | 4636.2    | ± 263.9 | 5544.5    | ± 366.8  | 5320.3    | ± 535.8  | 4668.4    | ± 439.8 | 89.757                        | 0.0001 |
| ZD1 m              | 1320.5    | ± 139.8  | 1401.8    | ± 96.2  | 1672.0    | ± 129.1 | 1382.4    | ± 122.6  | 1679.6    | ± 98.6   | 1596.8    | ± 107.2 | 102.348                       | 0.0001 |
| ZD2 m              | 533.0     | ± 581.7  | 2010.8    | ± 149.3 | 1806.1    | ± 164.4 | 2175.2    | ± 209.0  | 1605.3    | ± 163.4  | 1764.3    | ± 206.4 | 95.502                        | 0.0001 |
| ZD3 m              | 92.9      | ± 158.9  | 533.6     | ± 77.8  | 433.8     | ± 84.0  | 692.0     | ± 144.0  | 529.5     | ± 100.7  | 375.2     | ± 63.7  | 92.521                        | 0.0001 |
| ZD4 m              | 126.4     | ± 260.6  | 707.3     | ± 115.2 | 475.6     | ± 102.7 | 853.9     | ± 194.3  | 741.8     | ± 172.2  | 479.4     | ± 56.9  | 94.934                        | 0.0001 |
| ZD5 m              | 55.7      | ± 162.4  | 472.2     | ± 110.0 | 211.8     | ± 58.3  | 380.9     | ± 95.1   | 561.4     | ± 143.2  | 338.4     | ± 71.2  | 107.543                       | 0.0001 |
| ZD6 m              | 8.5       | ± 28.3   | 155.1     | ± 76.5  | 33.9      | ± 23.9  | 56.9      | ± 41.1   | 199.6     | ± 89.9   | 110.9     | ± 64.9  | 102.121                       | 0.0001 |
| ΣD m (1,2,3)       | 1947.8    | ± 800.4  | 3947.8    | ± 222.4 | 3913.4    | ± 217.2 | 4251.2    | ± 246.7  | 3816.1    | ± 248.0  | 3737.8    | ± 310.3 | 63.108                        | 0.0001 |
| ΣD m (4,5,6)       | 191.4     | ± 451.0  | 1336.2    | ± 227.7 | 722.8     | ± 149.1 | 1293.3    | ± 251.1  | 1504.3    | ± 362.8  | 930.5     | ± 161.5 | 105.854                       | 0.0001 |
| Acc1 m/s/s         | 34.5      | ± 35.0   | 117.5     | ± 19.4  | 107.2     | ± 14.8  | 135.0     | ± 23.4   | 103.6     | ± 16.4   | 87.1      | ± 12.1  | 71.114                        | 0.0001 |
| Acc2 m/s/s         | 9.6       | ± 10.6   | 43.1      | ± 8.1   | 26.8      | ± 6.9   | 36.8      | ± 12.5   | 48.3      | ± 13.0   | 33.2      | ± 6.8   | 76.501                        | 0.0001 |
| Acc3 m/s/s         | 1.7       | ± 1.6    | 8.8       | ± 3.9   | 3.3       | ± 2.1   | 4.9       | ± 2.7    | 11.3      | ± 3.7    | 10.8      | ± 4.7   | 92.085                        | 0.0001 |
| ΣAcc (1,2,3) m/s/s | 45.8      | ± 46.6   | 169.3     | ± 23.0  | 137.4     | ± 16.9  | 176.7     | ± 33.7   | 163.3     | ± 28.1   | 131.1     | ± 16.3  | 69.553                        | 0.0001 |
| Des1 m/s/s         | 31.7      | ± 28.0   | 99.3      | ± 15.5  | 95.5      | ± 12.7  | 116.1     | ± 17.8   | 83.1      | ± 12.5   | 74.9      | ± 10.7  | 81.279                        | 0.0001 |
| Des2 m/s/s         | 8.2       | ± 9.1    | 36.5      | ± 7.5   | 26.4      | ± 5.8   | 38.9      | ± 11.5   | 39.4      | ± 9.0    | 28.0      | ± 6.3   | 69.645                        | 0.0001 |
| Des3 m/s/s         | 2.5       | ± 4.6    | 17.9      | ± 5.2   | 10.4      | ± 4.2   | 12.9      | ± 5.2    | 25.6      | ± 7.7    | 19.8      | ± 4.1   | 90.250                        | 0.0001 |
| ΣDes (1,2,3) m/s/s | 42.4      | ± 41.0   | 153.7     | ± 15.6  | 132.2     | ± 15.8  | 167.9     | ± 30.3   | 148.1     | ± 23.7   | 122.7     | ± 15.7  | 68.209                        | 0.0001 |
| MS km/h            | 20.2      | ± 3.1    | 29.7      | ± 1.9   | 27.4      | ± 2.2   | 28.0      | ± 2.1    | 30.6      | ± 1.5    | 30.6      | ± 2.0   | 70.830                        | 0.0001 |

Note. Goalkeeper = GK Wide defenders = WD, central defenders = CD, center midfielder = CM, open attackers = OA) and center forward = CF.

**Table 3.** Comparisons between playing positions for each of the speed thresholds.

| Variables          | WD >               | CD >         | CM >                     | OA >                    | CF >        |
|--------------------|--------------------|--------------|--------------------------|-------------------------|-------------|
| TD m               | DC **/CD **        |              | DL **/DC **/CD **        | DC **/CD **             |             |
| TDP m              | DC **/CD **        |              | DL **/DC **/CD **        | DC **/CD **             |             |
| ZD1 m              |                    | DL **/MC **  |                          | DL **/MC **/CD **       | DL **       |
| ZD2 m              | DC **/EXT **/CD ** | EXT **       | DL **/DC **/EXT **/CD ** |                         | EXT **      |
| ZD3 m              | DC **/CD **        | CD *         | DL **/DC **/EXT **/CD ** | DC **/CD **             |             |
| ZD4 m              | DC **/CD **        |              | DL **/DC **/CD **        | DC **/CD **             |             |
| ZD5 m              | DC **/MC **/CD **  |              | DC **                    | DL **/DC **/MC **/CD ** | DC **       |
| ZD6 m              | DC **/MC **/CD **  |              | DC **                    | DL **/DC **/MC **/CD ** | DC **/MC ** |
| ΣD m (1,2,3)       | EXT **/CD **       |              | DL **/DC **/EXT **/CD ** |                         |             |
| ΣD m (4,5,6)       | DC **/CD **        |              | DC **/CD **              | DL **/DC **/MC **/CD ** | DC **       |
| Acc1 m/s/s         | DC **/EXT **/CD ** | CD **        | DL **/DC **/EXT **/CD ** | CD **                   |             |
| Acc2 m/s/s         | DC **/MC **/CD **  |              | DC **                    | DC **/MC **/CD **       | DC **       |
| Acc3 m/s/s         | DC **/MC **        |              | DC **                    | DL **/DC **/MC **       | DC **/MC ** |
| ΣAcc (1,2,3) m/s/s | DC **/CD **        |              | DC **/CD **              | DC **/CD **             |             |
| Des1 m/s/s         | EXT **/CD **       | EXT **/CD ** | DL **/DC **/EXT **/CD ** | CD *                    |             |
| Des2 m/s/s         | DC **/CD **        |              | DC **/CD **              | DC **/CD **             |             |
| Des3 m/s/s         | DC **/MC **        |              | DC **                    | DL **/DC **/MC **/CD ** | DC **/MC ** |
| ΣDes (1,2,3) m/s/s | DC **/CD **        |              | DC **/EXT **/CD **       | DC **/CD **             |             |
| MS km/h            | DC **/MC **        |              |                          | DL **/DC **/MC **       | DC **/MC ** |

Note. GK comparisons are omitted from analyses. Wide defenders = WD, central defenders = CD, center midfielder = CM, open attackers = OA) and center forward = CF. \*  $p \leq 0.05$ ; \*\*  $p \leq 0.001$

**Table 4.** Distances covered across each speed threshold when the teams drew, lost, or won (Data reported as mean ± SD).

| Variables          | Draw n = 14 |         | Lose n = 76 |          | Win n = 70 |          | Difference between Groups |       | Difference between Pairs |     |          |
|--------------------|-------------|---------|-------------|----------|------------|----------|---------------------------|-------|--------------------------|-----|----------|
|                    | Mean        | SD      | Mean        | SD       | Mean       | SD       | H de Kruskal-Wallis       | Sig   | D >                      | L > | W >      |
| TD m               | 10,395.1    | ± 875.0 | 9414.9      | ± 2050.2 | 9978.5     | ± 1962.5 | 9.342                     | 0.009 | $p^*$                    |     | $p^{**}$ |
| TDP m              | 5197.6      | ± 452.4 | 4707.5      | ± 1048.2 | 5062.4     | ± 899.2  | 9.170                     | 0.010 |                          |     | $p^{**}$ |
| ZD1 m              | 1518.4      | ± 186.3 | 1496.6      | ± 185.7  | 1565.6     | ± 175.2  | 3.996                     | 0.136 |                          |     | $p^*$    |
| ZD2 m              | 1905.3      | ± 270.2 | 1706.6      | ± 489.2  | 1875.5     | ± 428.5  | 10.052                    | 0.007 |                          |     | $p^{**}$ |
| ZD3 m              | 533.1       | ± 95.8  | 475.1       | ± 196.8  | 511.7      | ± 177.6  | 3.518                     | 0.172 |                          |     |          |
| ZD4 m              | 675.3       | ± 183.1 | 597.3       | ± 251.9  | 642.3      | ± 259.0  | 1.361                     | 0.506 |                          |     |          |
| ZD5 m              | 433.5       | ± 165.0 | 339.4       | ± 183.0  | 367.0      | ± 178.3  | 3.209                     | 0.201 |                          |     |          |
| ZD6 m              | 128.9       | ± 104.7 | 89.4        | ± 83.8   | 97.1       | ± 89.9   | 2.443                     | 0.295 |                          |     |          |
| ΣD m (1,2,3)       | 3958.3      | ± 179.6 | 3679.8      | ± 680.1  | 3954.4     | ± 570.5  | 20.241                    | 0.000 |                          |     | $p^{**}$ |
| ΣD m (4,5,6)       | 1239.2      | ± 423.4 | 1027.6      | ± 458.0  | 1108.0     | ± 462.7  | 2.312                     | 0.315 |                          |     |          |
| Acc1 m/s/s         | 116.6       | ± 22.2  | 103.0       | ± 34.0   | 111.1      | ± 28.7   | 4.960                     | 0.084 |                          |     | $p^*$    |
| Acc2 m/s/s         | 36.9        | ± 11.2  | 34.6        | ± 13.7   | 35.6       | ± 15.4   | 0.151                     | 0.927 |                          |     |          |
| Acc3 m/s/s         | 7.9         | ± 4.9   | 6.6         | ± 4.8    | 6.4        | ± 4.4    | 1.019                     | 0.601 |                          |     |          |
| ΣAcc (1,2,3) m/s/s | 161.4       | ± 27.2  | 144.2       | ± 45.1   | 153.1      | ± 41.0   | 2.851                     | 0.240 |                          |     |          |
| Des1 m/s/s         | 94.2        | ± 17.4  | 89.3        | ± 29.0   | 95.0       | ± 23.4   | 2.407                     | 0.300 |                          |     |          |
| Des2 m/s/s         | 33.7        | ± 6.8   | 31.3        | ± 12.2   | 32.9       | ± 12.4   | 0.665                     | 0.717 |                          |     |          |
| Des3 m/s/s         | 18.1        | ± 7.9   | 14.6        | ± 8.1    | 15.5       | ± 8.6    | 2.141                     | 0.343 |                          |     |          |
| ΣDes (1,2,3) m/s/s | 146.0       | ± 21.5  | 135.2       | ± 42.2   | 143.4      | ± 35.3   | 2.586                     | 0.274 |                          |     |          |
| MS km/h            | 29.8        | ± 1.8   | 28.1        | ± 3.4    | 28.3       | ± 3.2    | 3.944                     | 0.139 | $p^*$                    |     |          |

\*  $p \leq 0.05$ ; \*\*  $p \leq 0.001$ .

**Table 5.** Distances covered across each speed threshold according to first and second half (data are reported as mean ± SD).

| Variables          | 1st Part <i>n</i> = 80 |         | 2nd Part <i>n</i> = 80 |         | Difference between 1st and 2nd part |       |       |       |
|--------------------|------------------------|---------|------------------------|---------|-------------------------------------|-------|-------|-------|
|                    | Mean                   | SD      | Mean                   | SD      | U de Mann-Whitney                   | Sig   | 1st > | 2nd > |
| TDP m              | 4942.7                 | ± 945.2 | 4868.6                 | ± 980.5 | 2969.5                              | 0.432 |       |       |
| ZD1 m              | 1483.3                 | ± 176.6 | 1574.1                 | ± 179.3 | 2320.5                              | 0.003 |       | 1st * |
| ZD2 m              | 1829.3                 | ± 437.2 | 1766.5                 | ± 471.0 | 2814.5                              | 0.188 |       |       |
| ZD3 m              | 500.6                  | ± 183.9 | 491.8                  | ± 181.2 | 3124.0                              | 0.795 |       |       |
| ZD4 m              | 639.9                  | ± 254.0 | 607.7                  | ± 246.5 | 2950.0                              | 0.394 |       |       |
| ZD5 m              | 380.1                  | ± 186.1 | 339.3                  | ± 173.2 | 2776.0                              | 0.148 |       |       |
| ZD6 m              | 106.5                  | ± 91.9  | 86.0                   | ± 84.3  | 2785.5                              | 0.157 |       |       |
| ΣD m (1,2,3)       | 3814.7                 | ± 585.8 | 3834.0                 | ± 650.9 | 2987.0                              | 0.467 |       |       |
| ΣD m (4,5,6)       | 1128.0                 | ± 471.1 | 1034.6                 | ± 444.0 | 2777.0                              | 0.149 |       |       |
| Acc1 m/s/s         | 110.1                  | ± 31.8  | 105.4                  | ± 30.4  | 2819.0                              | 0.193 |       |       |
| Acc2 m/s/s         | 36.1                   | ± 15.2  | 34.4                   | ± 13.3  | 3039.5                              | 0.584 |       |       |
| Acc3 m/s/s         | 7.0                    | ± 5.1   | 6.3                    | ± 4.2   | 3072.5                              | 0.663 |       |       |
| ΣAcc (1,2,3) m/s/s | 153.1                  | ± 43.1  | 146.1                  | ± 41.2  | 2742.0                              | 0.118 |       |       |
| Des1 m/s/s         | 94.5                   | ± 26.5  | 90.0                   | ± 25.1  | 2892.0                              | 0.293 |       |       |
| Des2 m/s/s         | 33.6                   | ± 12.5  | 30.7                   | ± 11.2  | 2721.0                              | 0.102 |       |       |
| Des3 m/s/s         | 16.3                   | ± 9.0   | 14.3                   | ± 7.5   | 2803.0                              | 0.175 |       |       |
| ΣDes (1,2,3) m/s/s | 144.4                  | ± 38.7  | 135.1                  | ± 36.7  | 2585.5                              | 0.036 | 2nd * |       |
| MS km/h            | 28.5                   | ± 3.6   | 28.1                   | ± 2.9   | 2761.5                              | 0.134 |       |       |

\*  $p \leq 0.05$

**Table 6.** Distances covered across each speed threshold when the teams competed home or away (Data reported as mean ± SD).

| Variables          | Local <i>n</i> = 60 |          | Visit <i>n</i> = 100 |          | Difference between Groups |          |      |     |
|--------------------|---------------------|----------|----------------------|----------|---------------------------|----------|------|-----|
|                    | Mean                | SD       | Mean                 | SD       | U de Mann-Whitney         | Sig      | L >  | V > |
| TD m               | 10,208.0            | ± 1926.2 | 9470.8               | ± 1932.4 | 1948.0                    | 0.0002   | V ** |     |
| TDP m              | 5104.0              | ± 974.1  | 4786.6               | ± 937.4  | 2050.5                    | 0.0008   | V ** |     |
| ZD1 m              | 1549.8              | ± 190.3  | 1516.1               | ± 178.5  | 2719.0                    | 0.3220   |      |     |
| ZD2 m              | 1892.5              | ± 476.4  | 1741.1               | ± 432.6  | 2109.5                    | 0.0017   | V ** |     |
| ZD3 m              | 530.5               | ± 193.3  | 475.6                | ± 172.7  | 2315.0                    | 0.0158   | V ** |     |
| ZD4 m              | 654.9               | ± 266.3  | 605.1                | ± 239.2  | 2648.0                    | 0.2147   |      |     |
| ZD5 m              | 370.5               | ± 189.4  | 353.2                | ± 175.4  | 2866.0                    | 0.6367   |      |     |
| ZD6 m              | 102.6               | ± 97.5   | 92.4                 | ± 82.9   | 2896.0                    | 0.7138   |      |     |
| ΣD m (1,2,3)       | 3974.3              | ± 625.1  | 3734.3               | ± 597.8  | 1666.0                    | 0.000003 | V ** |     |
| ΣD m (4,5,6)       | 1129.7              | ± 483.0  | 1052.3               | ± 443.4  | 2691.0                    | 0.2761   |      |     |
| Acc1 m/s/s         | 111.7               | ± 32.1   | 105.3                | ± 30.4   | 2518.5                    | 0.0896   |      |     |
| Acc2 m/s/s         | 34.4                | ± 15.5   | 35.7                 | ± 13.5   | 2806.0                    | 0.4940   |      |     |
| Acc3 m/s/s         | 5.6                 | ± 4.0    | 7.3                  | ± 4.9    | 2444.0                    | 0.0493   |      | L * |
| ΣAcc (1,2,3) m/s/s | 151.8               | ± 44.4   | 148.3                | ± 41.0   | 2826.5                    | 0.5408   |      |     |
| Des1 m/s/s         | 96.0                | ± 26.2   | 90.0                 | ± 25.4   | 2464.0                    | 0.0588   |      |     |
| Des2 m/s/s         | 34.1                | ± 12.9   | 31.0                 | ± 11.2   | 2596.5                    | 0.1547   |      |     |
| Des3 m/s/s         | 14.2                | ± 8.4    | 16.0                 | ± 8.2    | 2603.5                    | 0.1618   |      |     |
| ΣDes (1,2,3) m/s/s | 144.3               | ± 39.4   | 137.0                | ± 36.9   | 2497.0                    | 0.0762   |      |     |
| MS km/h            | 28.2                | ± 3.2    | 28.4                 | ± 3.3    | 2906.0                    | 0.7404   |      |     |

\*  $p \leq 0.05$ ; \*\*  $p \leq 0.001$ .

**Table 7.** Summary of each 15-min period of matches, playing positions, match result, first and second half and match location (Data reported as mean  $\pm$  SD).

| Variables | <i>n</i>          | Mean | SD             | H de Kruskal-Wallis | Sig   | U de Mann-Whitney      |
|-----------|-------------------|------|----------------|---------------------|-------|------------------------|
| Quartile  | 00-15 = Q1 >      | 378  | 5.04 $\pm$ 2.9 | 82.234              | 0.000 | Q4 **/Q5 */Q7 **/Q8 ** |
|           | 16-30 = Q2 >      | 360  | 4.68 $\pm$ 3.0 |                     |       | Q7 **/Q8 **            |
|           | 31-45 = Q3 >      | 335  | 4.72 $\pm$ 2.8 |                     |       | Q4 */Q7 **/Q8 **       |
|           | 46-60 = Q4 >      | 264  | 3.77 $\pm$ 2.5 |                     |       | Q7 **/Q8 **            |
|           | 61-75 = Q5 >      | 287  | 3.99 $\pm$ 2.5 |                     |       | Q7 **/Q8 **            |
|           | 76-90 = Q6 >      | 319  | 4.09 $\pm$ 2.2 |                     |       | Q7 **/Q8 **            |
|           | More of 45 = Q7 > | 28   | 1.27 $\pm$ 0.5 |                     |       |                        |
|           | More of 90 = Q8 > | 106  | 2.12 $\pm$ 1.1 |                     |       | Q7 **                  |
| Positions | GK >              | 41   | 1.71 $\pm$ 1.0 | 107.057             | 0.000 | GK **/CD **            |
|           | WD >              | 526  | 4.78 $\pm$ 2.7 |                     |       | GK **                  |
|           | CD >              | 301  | 2.47 $\pm$ 1.5 |                     |       | GK **/CD **/           |
|           | CM >              | 444  | 4.11 $\pm$ 2.6 |                     |       | GK **/CD **/CM **/CF * |
|           | OA >              | 544  | 5.39 $\pm$ 2.6 |                     |       | GK **CD **             |
|           | CF >              | 221  | 4.42 $\pm$ 2.8 |                     |       |                        |
| Result    | D >               | 400  | 4.35 $\pm$ 2.9 | 1.091               | 0.579 | No difference          |
|           | L >               | 761  | 3.90 $\pm$ 2.6 |                     |       |                        |
|           | W >               | 916  | 4.02 $\pm$ 2.6 |                     |       |                        |
| Parts     | 1st >             | 1104 | 4.52 $\pm$ 2.9 |                     |       | 27,322.0/0.001/2da **  |
|           | 2nd >             | 973  | 3.59 $\pm$ 2.3 |                     |       |                        |
| Place     | Local             | 702  | 3.84 $\pm$ 2.6 |                     |       | 465,352.0/0.433/No Dif |
|           | Visit             | 1375 | 4.14 $\pm$ 2.7 |                     |       |                        |

\*  $p \leq 0.05$ ; \*\*  $p \leq 0.001$ .

#### 4. Discussion

The aim of this study was to analyse performance in professional Portuguese soccer players across different playing positions and contextual variables. Different positions require different physical demands, which are influenced by the location of the match and the result. Overall, players covered a mean TD of  $9839 \pm 1929$  m, with midfielders covering the greatest distance and central defenders performing the least TD. Wingers covered the most high-speed distance, number of decelerations (3.0–4.0 m/s/s) and were the fastest players, along with center forwards. Players performed greater distances when their team drew or won, and reductions for some running metrics were observed in the 2nd half of matches. The number of faster accelerations (3.0–4.0 m/s/s) was higher in away versus home games. These data may be considered by coaches and physical practitioners as a guide for the training prescription for their teams.

The TD covered was similar with values reported by studies conducted in soccer players across other European countries [1,28]. Specifically, the mean TD covered was similar in the present study for CM ( $10,788 \pm 1537$  m) and OA ( $10,641 \pm 1537$  1011 m), compared to the 2006–2007 ( $10,679 \pm 956$ ) and 2012–13 ( $10,881 \pm 885$  m) seasons in the English Premier League [1]. However, lower sprint distances across all playing positions were reported in the present study ( $\leq 199.6 \pm 89.9$  m) versus the English Premier League players ( $232 \pm 114$  (2006–2007 season) and  $350 \pm 139$  m (2012–2013 season)). Therefore, this study presents novel evidence that soccer players competing in the second Portuguese league cover similar distances during competitive match-play with English Premier League players, but lesser sprint distances. The discrepancy in sprint distance may be reflective of the players across both studies competing in different countries (tactical modifications) and tiers (high-standard athletes). This is largely supported by evidence that suggests high-speed and sprint performance is superior in higher level soccer players during match-play [11]. These data indicate that practitioners may direct their focus on the development of sprint capacity to increase physical performance of lower division players during match-play,

considering the specific demands of each position for the training prescription, optimizing the preparation process.

High-speed running performance is thought to be capable of discriminating between competitive levels and is a better indicator of physical performance versus TD alone [1]. However, a paucity of published information is available on the differences in high-speed running across playing positions in the Portuguese soccer leagues. In relation to high-speed running performance across different playing positions in the current investigation, it appears that CD perform considerably less high-speed activity versus all other positional roles (excluding the GK). This observation largely agrees with a previous study assessing English Championship and Premier League players [3]. The study found that English championship ( $540 \pm 129$  m) and Premier League players ( $482 \pm 116$  m) cover considerably less high-speed running distance compared with all other outfield positions, showing significant performance differences between positions. In relation to the WD and OA players, the results are consistent with players of wider positions in the English Premier League high-speed running values [29]. These results suggest that training programmes for soccer players must be individualised and specific to the demands of the playing position to enhance physical performance during match-play.

The number of accelerations and decelerations present a similar pattern as identified in previous studies, with a greater quantity of actions reported in the 1st half versus the 2nd half [30], but it differs in that the position that performs the greatest number of accelerations in the present study is the CM, while in preparation games in Spanish soccer it is the CD [13], context that can influence these returns. Concerning the playing roles, the highest number of Acc1 (1.0–1.9 m/s/s) was performed by CM, who presented substantially higher values versus all positions. With reference to Acc2 (2.0–2.9 m/s/s), the highest quantity of actions was performed by the OA, who had differences with all positions except the WD, who also presented differences with CD, CF, and CM. Pertaining to Acc3 (3.0–4.0 m/s/s), the OA and CF achieved the highest performance and were considerably different compared to other positions (CM, CD, and WD). This suggests that the quantity of accelerations performed at different speeds varies depending on the demands of the position. These data can, therefore, be used as a guide to inform acceleration-based training paradigms that incorporate the different acceleration loads of each playing position.

Decelerations are also an important component to consider in soccer, since players must frequently decelerate at high speeds, placing lower-limb musculature under a high eccentric loading demand [31]. The number of decelerations (1.0–4.0 m/s/s) performed by CM players was higher than CD, CF, and OA. These results may reflect the greater CM involvement in the game, both in offensive and defensive tasks. However, the OA players performed significantly more Dec3 (3.0–4.0 m/s/s) than other positions. This is possibly linked with the fact that OAs tend to run faster ( $30.6 \pm 1.5$ ) than other players, which necessitates a higher frequency of rapid decelerations. These demanding actions are associated with increased fatigue, which decreases locomotive efficiency at the end of each half in competitive football and thus increases the likelihood of injury incidence [32]. In this way, both accelerations and decelerations are another indicator that shows the variability of effort between the different positions; therefore, it is recommended that the functional specificity and movement drills incorporated within training programmes expose players to the quantity of accelerations and decelerations reported in the current data.

Lower performance was observed in the games played away from home, which may be influenced by tactical approaches and defensive strategies adopted to limit the opposition rather than providing an attacking threat [33]. There may also be additional logistical issues involved with travelling that limits players preparation and reduces overall performance output and recovery [34]. Thus, this specific aspect should be considered when scheduling the weekly planning and periodisation of training sessions [35]. Greater very-low intensity (DZ1) activity was performed in the 2nd half versus the 1st half. This may be a subconscious self-pacing approach to conserve energy during the second half, to avoid significant reductions in high-speed capacity [36] or avoid fatigue-induced injury

during the latter stages of match-play [37]. It is also possible that the reductions in intensity during the 2nd half are linked to a fatigue-induced inability to perform the actions rather than pacing strategies employed to maintain energy for the final moments of a match [38]. It is integral that physical capacity is maintained for the entire 90 min as approximately 57% of goals are scored in the 2nd half [39,40]. Therefore, it is crucial that players are conditioned to be able to compete for an entire 90 min and that substitutions are utilised for the players incapable of performing the required running output across an entire match [41].

The SBA estimated in 6 blocks of 15 min showed a gradual decrease as the game progressed in the 1st half, while during the 2nd half the trend was the opposite and the lowest performance was presented at the beginning and increased during the game. This is perhaps associated with declines in body temperature that occur due to passive recovery in the changing rooms and a short and ineffective reactivation at the beginning of the 2nd half [42]. This information should be used to better manage reactivation during matches and to create recovery contexts during training sessions that reflect what happens in competition. This may allow optimal adaptation to passive recovery and reduce insufficient performance during the start of each half. Adequate half-time re-warm-up and passive heat maintenance strategies may also enhance performance in the initial stages of the 2nd half [43]. Thus, adherence to such protocols may gain soccer teams the competitive advantage during the early stages of the second half of matches.

## 5. Conclusions

The quantification of the demands of second-division Portuguese matches through GNSS devices presents novel perspectives. The current data suggest that the physical demands of each playing position are impacted by match location and result. The demands of each playing position are reported throughout the article, with midfield players covering the most distance and largely performing the greatest quantity of actions throughout match-play. Players completed greater distances when their teams had a positive result (i.e., won or draw), and running performance largely decreased in the 2nd half of matches, an observation that is consistent with the literature. Based on the current findings, training drills should be position-specific, where appropriate, to prepare players for the physical capacity and running output of the various positional roles during matches. Future studies are required to evaluate match running profiles across the major European Leagues (German Bundesliga, French Ligue 1 and the Spanish La Liga) to compare with the current data and the wealth of observations available on English Premier League players.

**Author Contributions:** J.B., H.S. and A.J.F. led the project, established the protocol, and wrote and revised the original manuscript. F.M.C. and A.F. wrote and revised the original manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Institutional Review Board (or Ethics Committee) of University of Coimbra (code: CE/FCDEF-UC/0069202).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Barnes, C.; Archer, D.T.; Hogg, B.; Bush, M.; Bradley, P.S. The Evolution of Physical and Technical Performance Parameters in the English Premier League. *Int. J. Sports Med.* **2014**, *35*, 1095–1100. [[CrossRef](#)]
2. Di Salvo, V.; Gregson, W.; Atkinson, G.; Tordoff, P.; Drust, B. Analysis of High Intensity Activity in Premier League Soccer. *Int. J. Sports Med.* **2009**, *30*, 205–212. [[CrossRef](#)]
3. Di Salvo, V.; Pigozzi, F.; González-Haro, C.; Laughlin, M.S.; de Witt, J.K. Match Performance Comparison in Top English Soccer Leagues. *Int. J. Sports Med.* **2013**, *34*, 526–532. [[CrossRef](#)]
4. Malone, J.J.; di Michele, R.; Morgans, R.; Burgess, D.; Morton, J.P.; Drust, B. Seasonal Training-Load Quantification in Elite English Premier League Soccer Players. *Int. J. Sports Physiol. Perform.* **2015**, *10*, 489–497. [[CrossRef](#)]

5. Hands, D.E.; de Jonge, X.Y. Current time-motion analyses of professional football matches in top-level domestic leagues: A systematic review. *Int. J. Perform. Anal. Sport* **2020**, *20*, 747–765. [[CrossRef](#)]
6. Oliva-Lozano, J.M.; Rojas-Valverde, D.; Gómez-Carmona, C.D.; Fortes, V.; Pino-Ortega, J. Impact of contextual variables on the representative external load profile of Spanish professional soccer match-play: A full season study. *Eur. J. Sport Sci.* **2020**, 1–10. [[CrossRef](#)]
7. Bangsbo, J.; Nørregaard, L.; Thorsø, F. Activity profile of competition soccer. *Can. J. Sport Sci.* **1991**, *16*, 110–116.
8. Rampinini, E.; Impellizzeri, F.M.; Castagna, C.; Coutts, A.J.; Wisløff, U. Technical performance during soccer matches of the Italian Serie A league: Effect of fatigue and competitive level. *J. Sci. Med. Sport* **2009**, *12*, 227–233. [[CrossRef](#)]
9. Carling, C.; Dupont, G. Are declines in physical performance associated with a reduction in skill-related performance during professional soccer match-play? *J. Sports Sci.* **2011**, *29*, 63. [[CrossRef](#)]
10. Bradley, P.S.; Sheldon, W.; Wooster, B.; Olsen, P.D.; Boanas, P.; Krstrup, P. High-intensity running in English FA Premier League soccer matches. *J. Sports Sci.* **2009**, *27*, 159–168. [[CrossRef](#)]
11. Mohr, M.; Krstrup, P.; Bangsbo, J. Match performance of high-standard soccer players with special reference to development of fatigue. *J. Sports Sci.* **2003**, *21*, 519–528. [[CrossRef](#)]
12. Morgans, R.; Orme, P.; Anderson, L.; Drust, B.; Morton, J.P. An Intensive Winter Fixture Schedule Induces a Transient Fall in Salivary IgA in English Premier League Soccer Players. *Res. Sports Med.* **2014**, *22*, 346–354. [[CrossRef](#)]
13. Mallo, J.; Mena, E.; Nevado, F.; Paredes, V. Physical Demands of Top-Class Soccer Friendly Matches in Relation to a Playing Position Using Global Positioning System Technology. *J. Hum. Kinet.* **2015**, *47*, 179–188. [[CrossRef](#)]
14. Russell, M.; Sparkes, W.; Northeast, J.; Cook, C.J.; Love, T.D.; Bracken, R.M.; Kilduff, L.P. Changes in Acceleration and Deceleration Capacity Throughout Professional Soccer Match-Play. *J. Strength Cond. Res.* **2016**, *30*, 2839–2844. [[CrossRef](#)]
15. Bangsbo, J. The physiology of soccer—With special reference to intense intermittent exercise. *Acta Physiol. Scand. Suppl.* **1994**, *619*, 1–155.
16. Stølen, T.; Chamari, K.; Castagna, C.; Wisløff, U. Physiology of Soccer. *Sports Med.* **2005**, *35*, 501–536. [[CrossRef](#)]
17. Bloomfield, J.; Polman, R.; O'Donoghue, P. Physical Demands of Different Positions in FA Premier League Soccer. *J. Sports Sci. Med.* **2007**, *6*, 63–70.
18. Rienzi, E.; Drust, B.; Reilly, T.; Carter, J.E.; Martin, A. Investigation of anthropometric and work-rate profiles of elite South American international soccer players. *J. Sports Med. Phys. Fit.* **2000**, *40*, 162.
19. Carling, C.; Bloomfield, J.; Nelsen, L.; Reilly, T. The role of motion analysis in elite soccer: Contemporary performance measurement techniques and work rate data. *Sports Med.* **2012**, *38*, 389. [[CrossRef](#)]
20. Lago, C. The influence of match location, quality of opposition, and match status on possession strategies in professional association football. *J. Sports Sci.* **2009**, *27*, 1463–1469. [[CrossRef](#)]
21. Rampinini, E.; Coutts, A.J.; Castagna, C.; Sassi, R.; Impellizzeri, F.M. Variation in top level soccer match performance. *Int. J. Sports Med.* **2007**, *28*, 1018–1024. [[CrossRef](#)]
22. Barros, R.M.; Misuta, M.S.; Menezes, R.P.; Figueroa, P.J.; Moura, F.A.; Cunha, S.A.; Anido, R.; Leite, N.J. Analysis of the Distances Covered by First Division Brazilian Soccer Players Obtained with an Automatic Tracking Method. *J. Sports Sci. Med.* **2007**, *6*, 233. [[PubMed](#)]
23. González-Rodenas, J.; Aranda-Malavés, R.; Tudela-Desantes, A.; Moreno, F.C.; Casal, C.A.; Aranda, R. Effect of Match Location, Team Ranking, Match Status and Tactical Dimensions on the Offensive Performance in Spanish 'La Liga' Soccer Matches. *Front. Psychol.* **2019**, *10*, 2089. [[CrossRef](#)] [[PubMed](#)]
24. Castellano, J.; Blanco-Villaseñor, A.; Álvarez, D. Contextual Variables and Time-Motion Analysis in Soccer. *Int. J. Sports Med.* **2011**, *32*, 415–421. [[CrossRef](#)]
25. Saunders, M.; Blevins, J.; Broeder, C. Effects of hydration changes on bioelectrical impedance in endurance trained individuals. *Med. Sci. Sports Exerc.* **1998**, *30*, 885–892. [[PubMed](#)]
26. Johnston, R.J.; Watsford, M.L.; Kelly, S.J.; Pine, M.J.; Spurrs, R.W. Validity and Interunit Reliability of 10 Hz and 15 Hz GPS Units for Assessing Athlete Movement Demands. *J. Strength Cond. Res.* **2014**, *28*, 1649–1655. [[CrossRef](#)]
27. Mara, J.K.; Thompson, K.G.; Pumpa, K.L. Physical and Physiological Characteristics of Various-Sided Games in Elite Women's Soccer. *Int. J. Sports Physiol. Perform.* **2016**, *11*, 953–958. [[CrossRef](#)]
28. Smpokos, E.; Mourikis, C.; Linardakis, M. Seasonal physical performance of a professional team's football players in a national league and European matches. *J. Hum. Sport Exerc.* **2018**, *13*, 720–730. [[CrossRef](#)]
29. Bush, M.; Barnes, C.; Archer, D.T.; Hogg, B.; Bradley, P.S. Evolution of match performance parameters for various playing positions in the English Premier League. *Hum. Mov. Sci.* **2015**, *39*, 1–11. [[CrossRef](#)]
30. Dalen, T.; Lorås, H.; Hjelde, G.H.; Kjøsnes, T.N.; Wisløff, U. Accelerations—A new approach to quantify physical performance decline in male elite soccer? *Eur. J. Sport Sci.* **2019**, *19*, 1015–1023. [[CrossRef](#)]
31. Mair, S.D.; Seaber, A.V.; Glisson, R.R.; Garrett, W.E. The Role of Fatigue in Susceptibility to Acute Muscle Strain Injury. *Am. J. Sports Med.* **1996**, *24*, 137–143. [[CrossRef](#)]
32. Barrett, S.; Midgley, A.; Reeves, M.; Joel, T.; Franklin, E.; Heyworth, R.; Garrett, A.; Lovell, R. The within-match patterns of locomotor efficiency during professional soccer match play: Implications for injury risk? *J. Sci. Med. Sport* **2016**, *19*, 810–815. [[CrossRef](#)]

33. Staufenbiel, K.; Lobinger, B.; Strauss, B. Home advantage in soccer—A matter of expectations, goal setting and tactical decisions of coaches? *J. Sports Sci.* **2015**, *33*, 1932–1941. [[CrossRef](#)] [[PubMed](#)]
34. Fowler, P.; Duffield, R.; Vaile, J. Effects of Domestic Air Travel on Technical and Tactical Performance and Recovery in Soccer. *Int. J. Sports Physiol. Perform.* **2014**, *9*, 378–386. [[CrossRef](#)]
35. Kelly, V.G.; Coutts, A.J. Planning and monitoring training loads during the competition phase in team sports. *Strength Cond. J.* **2007**, *29*, 32. [[CrossRef](#)]
36. Di Salvo, V.; Baron, R.; Tschan, H.; Montero, F.J.C.; Bachl, N.; Pigozzi, F. Performance Characteristics According to Playing Position in Elite Soccer. *Int. J. Sports Med.* **2007**, *28*, 222–227. [[CrossRef](#)]
37. Small, K.; McNaughton, L.; Greig, M.; Lovell, R. The effects of multidirectional soccer-specific fatigue on markers of hamstring injury risk. *J. Sci. Med. Sport* **2010**, *13*, 120–125. [[CrossRef](#)]
38. Field, A.; Naughton, R.J.; Haines, M.; Lui, S.; Corr, L.D.; Russell, M.; Page, R.M.; Harper, L.D. The demands of the extra-time period of soccer: A systematic review. *J. Sport Health Sci.* **2020**. [[CrossRef](#)]
39. Simiyu, W.W.N. Analysis of goals scored in the 2010 world cup soccer tournament held in South Africa. *J. Phys. Educ. Sport* **2013**, *13*, 6–13.
40. Michailidis, Y.; Michailidis, C.; Primpa, E. Analysis of goals scored in European Championship 2012. *J. Hum. Sport Exerc.* **2013**, *8*, 367–375. [[CrossRef](#)]
41. Hills, S.P.; Barwood, M.J.; Radcliffe, J.N.; Cooke, C.B.; Kilduff, L.P.; Cook, C.J.; Russell, M. Profiling the Responses of Soccer Substitutes: A Review of Current Literature. *Sports Med.* **2018**, *48*, 2255–2269. [[CrossRef](#)]
42. Mohr, M.; Krstrup, P.; Nybo, L.; Nielsen, J.J.; Bangsbo, J. Muscle temperature and sprint performance during soccer matches—Beneficial effect of re-warm-up at half-time. *Scand. J. Med. Sci. Sports* **2004**, *14*, 156–162. [[CrossRef](#)]
43. Silva, L.M.; Neiva, H.P.; Marques, M.C.; Izquierdo, M.; Marinho, D.A. Effects of warm-up, post-warm-up, and re-warm-up strategies on explosive efforts in team sports: A systematic review. *Sports Med.* **2018**, *48*, 2285–2299. [[CrossRef](#)] [[PubMed](#)]