

Supplementary Materials

Copper–Calcium Hydroxide and Permanent Electrophoretic Current for Treatment of Apical Periodontitis

Agron Meto ¹, Etleva Droboniku ², Elisabetta Blasi ³, Bruna Colombari ³, Emiljano Tragaj ¹, Gabriele Cervino ⁴, Luca Fiorillo ^{4,5,*}, and Aida Meto ^{2,3,*}

¹ Department of Therapy, Faculty of Dentistry, University of Aldent, 1000 Tirana, Albania; agronmeto@yahoo.com (A.M.); emiljanotragaj@yahoo.com (E.T.)


² Department of Dental Therapy, Faculty of Dental Medicine, University of Medicine, 1005 Tirana, Albania; etleva.droboniku@umed.edu.al

³ Department of Surgical, Medical, Dental and Morphological Sciences with Interest in Transplant, Oncological and Regenerative Medicine, University of Modena and Reggio Emilia, 41125 Modena, Italy; elisabetta.blasi@unimore.it (E.B.); bruna.colombari@unimore.it (B.C.)

⁴ Department of Biomedical and Dental Sciences, Morphological and Functional Images, University of Messina, 98100 Messina, Italy; gcervino@unime.it

⁵ Multidisciplinary Department of Medical-Surgical and Odontostomatological Specialties, University of Campania “Luigi Vanvitelli”, 80121 Naples, Italy

* Correspondence: lfiorillo@unime.it (L.F.); aidameto@yahoo.com (A.M.); Tel.: +35-569-331-7677 (A.M. & L.F.)



Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).

Figure S1. An example of how focal size measurement was made using the ruler, before (a) and after (b) treatment in follow-up.