

Article

Stimulation, reduction and compensation growth, and variable phenological responses to spring and/or summer-autumn warming in *Corylus taxa* and *Cornus sanguinea* L.

 Kristine Vander Mijnsbrugge ^{1,*}, Jessa May Malanguis ², Stefaan Moreels ¹, Arion Turcsán ³ and Eduardo Notivol Paino ⁴
¹ Department of Forest Ecology and Management, Research Institute for Nature and Forest, 9500 Geraardsbergen, Belgium; stefaan.moreels@inbo.be

² Natural Sciences and Mathematics, Visayas Tacloban College, University of the Philippines, Tacloban City 6500, Philippines; malanguisjessamay@gmail.com

³ Bavarian State Institute of Forestry, Department Soil and Climate, Hans-Carl-von-Carlowitz-Platz 1, D-85354 Freising, Germany; raup25@gmail.com

⁴ Department of Environment, Agricultural and Forest Systems, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Avda. Montañana 930, 50059 Zaragoza, Spain; enotivol@cita-aragon.es

* Correspondence: kristine.vandermijnsbrugge@inbo.be

Supplementary Tables

Table S1. Estimates and p-values for the modeled height and diameter of the plants at the end of 2017, for the *Corylus taxa* and for *Cornus sanguinea*. For the *Corylus taxa*, the standard taxon/region is the Belgian *Corylus avellana* to which the Belgian intermediate forms (Be_Cx), the Belgian *Corylus maxima* (Be_Cmax) and the Spanish *Corylus avellana* (Sp_Cave) are compared. For *Cornus sanguinea*, the Belgian provenance is the standard provenance to which the Spanish provenance (Sp) is compared.

<i>Corylus taxa</i>					<i>Cornus sanguinea</i>				
	Height		Diameter			Height		Diameter	
Variable	Estimate	p-Value	Estimate	p-Value	Variable	Estimate	p-Value	Estimate	p-Value
(Intercept)	40.30	<0.001***	3.85	<0.001***	(Intercept)	81.54	<0.001***	4.55	<0.001***
Be_Cx	4.66	0.344	0.19	0.463	Sp	-14.91	<0.001***	-0.69	<0.001***
Be_Cmax	0.51	0.947	0.42	0.299					
Sp_Cave	-4.10	0.371	-0.04	0.864					

 *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.