

Hyaluronan-Cyclodextrin Conjugates as Doxorubicin Delivery Systems

Noemi Bognanni¹, Maurizio Viale², Luana La Piana¹, Simone Strano¹, Rosaria Gangemi², Cinzia Lombardo³, Maria Teresa Cambria³ and Graziella Vecchio^{1,*}

¹ Dipartimento di Scienze Chimiche, Università degli Studi di Catania, Viale A. Doria 6, 95125 Catania, Italy

² UOC Bioterapie, IRCCS Ospedale Policlinico San Martino, Largo R. Benzi 10, 16132 Genova, Italy

³ Dipartimento di Scienze Biomediche e Biotecnologiche, Sezione di Biochimica Medica, Università degli Studi di Catania, Via S. Sofia 97, 95125 Catania, Italy

* Correspondence: gr.vecchio@unict.it




Figure S1. ¹H NMR spectrum of HAL β CyD (D_2O , 500 MHz)




Figure S2. ¹H NMR spectrum of HAH β CyD (D_2O , 500 MHz)




Figure S3. UV-Vis spectra of saturated solution of DOXO in the presence of an increasing concentration of the polymer: HAL β CyD (Left) and HAH β CyD (right) at pH 7.4, phosphate buffer




Figure S4. Graphs represent the mean dose-response curves of SK-N-SH (top) and SK-N-SH-PMA (bottom) cells treated with DOXO or DOXO/HAL β CyD 8/1, DOXO/HAH β CyD 8/1, DOXO/HAL β CyD 16/1, and DOXO/HAH β CyD 16/1 complexes.