



**Supplementary table.** Presence of co-morbidities in 231 currently adult patients diagnosed with celiac disease in childhood with and without daily life restrictions caused by a gluten-free diet (GFD).

	Restrictions due to GFD		<i>p</i> -value
	Yes, n=107 %	No, n=124 %	
Celiac disease-associated diseases			
<i>Thyroid disease</i>	10	11	0.831
<i>Type 1 diabetes</i>	10	8	0.694
Other conditions			
<i>Allergy<sup>1</sup></i>	44	40	0.591
<i>Osteoporosis or fractures</i>	27	24	0.613
<i>Skin disease</i>	18	14	0.458
<i>Depression</i>	14	12	0.651
<i>Asthma</i>	10	13	0.470
<i>Miscarriages<sup>2</sup></i>	7	11	0.369
<i>Other intestinal disease<sup>3</sup></i>	5	8	0.325
<i>Hypertension</i>	6	4	0.540
<i>Rheumatic disease</i>	6	3	0.307
<i>Eating disorder</i>	5	3	0.746
<i>Cancer<sup>4</sup></i>	2	2	1.000

<sup>1</sup>E.g., allergies to animals, food, pollen, and medicine; <sup>2</sup>only in women; <sup>3</sup>E.g., Crohn's disease, ulcerative colitis, and gastric ulcer; <sup>4</sup>E.g., Hodgkin's lymphoma, non-Hodgkin's lymphoma, breast cancer, and brain cancer.