

Table S1. Characteristics of the sample.

		Control (N = 41)		GD (N = 297)							GD (N = 297)	
		n	%	n	%	p			Mean	SD		
Sex	Women	4	9.8%	19	6.4%	0.423	Age of onset of GD		29.10	12.42		
	Men	37	90.2%	278	93.6%		Duration of GD		5.23	6.02		
Education	Primary	7	17.1%	157	52.9%	<0.001 *	Number DSM-5 criteria for GD		7.13	1.80		
	Secondary	15	36.6%	112	37.7%				n	%		
Marital	University	19	46.3%	28	9.4%	0.014 *	Debts due to GD		No	122	41.1%	
	Single	17	41.5%	158	53.2%				Yes	175	58.9%	
	Married	23	56.1%	103	34.7%		Gambling		Non-stratetigic	148	49.8%	
	Divorced	1	2.4%	36	12.1%				Strategic	92	31.0%	
Social index	High	0	0.0%	8	2.7%	0.005 *			Mixed	57	19.2%	
	Mean-high	2	4.9%	19	6.4%		Substances use		n	%		
	Mean	3	7.3%	24	8.1%		Tobacco	Yes	138	46.5%		
	Mean-low	28	68.3%	113	38.0%		Alcohol	Yes	43	14.5%		
	Low	8	19.5%	133	44.8%		Other drugs	Yes	34	11.4%		
		Mean	SD	Mean	SD	p						
Age (years-old)		49.27	15.23	39.58	14.16	<0.001 *						
BMI (kg/m²)		24.79	1.98	26.48	5.04	0.035 *						

Note. GD: gambling disorder. BMI: body mass index. SD: standard deviation. DSM-5: Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition. * Bold: significant comparison.

Table S2. Partial correlation matrix.

	Control (N = 41)				Gambling disorder (N = 297)			
	Ghrelin	LEAP2	Leptin	Adeno.	Ghrelin	LEAP2	Leptin	Adeno.
¹ Psychopathology: SCL-90R								
Somatization	-0.088	0.220	0.128	-0.101	0.064	0.092	0.042	0.018
Obsessive/compulsive	0.119	0.047	-0.143	-0.079	0.043	0.141	0.084	0.046
Interpersonal sensitivity	0.173	0.266 [†]	0.043	-0.009	0.050	0.026	-0.028	0.028
Depressive	0.135	0.134	-0.116	0.048	0.073	0.056	0.040	-0.037
Anxiety	0.061	0.271 [†]	0.032	0.073	0.018	0.122	0.046	-0.019
Hostility	-0.065	0.058	0.020	0.064	0.070	0.055	-0.013	-0.006
Phobic anxiety	0.126	0.317 [†]	0.237 [†]	-0.119	0.023	0.057	0.119	0.082
Paranoid Ideation	0.199	0.304 [†]	0.097	-0.030	0.011	0.104	-0.021	0.021
Psychotic	0.168	0.192	0.063	0.079	0.028	0.074	0.023	0.004
GSI score	0.093	0.226	-0.016	0.001	0.054	0.092	0.033	0.010
PST score	0.151	0.179	-0.031	-0.069	0.042	0.063	0.018	0.004
PSDI score	-0.075	0.370 [†]	0.093	0.155	0.057	0.080	0.016	-0.013
¹ Impulsivity: UPPS-P								
Lack premeditation	0.012	-0.064	0.089	-0.290 [†]	0.040	-0.018	-0.028	0.024
Lack perseverance	0.029	-0.184	-0.098	-0.150	0.009	0.013	0.033	0.066
Sensation seeking	-0.079	-0.234	-0.123	-0.004	-0.013	-0.009	-0.101	0.058
Positive urgency	-0.055	-0.120	0.006	-0.183	0.028	0.096	-0.074	0.004
Negative urgency	-0.091	0.132	0.126	-0.159	0.050	0.073	-0.055	0.019
Total	-0.068	-0.146	-0.010	-0.207	0.027	0.053	-0.076	0.050
¹ Gambling severity								
DSM-5 criteria	-0.073	0.141	0.073	-0.126	0.029	0.013	0.081	-0.041
² Neuropsychological data								
IGT: block 1	0.150	0.070	0.003	-0.002	0.009	0.091	-0.037	0.054
IGT: block 2	0.005	0.141	-0.058	0.095	-0.013	-0.002	-0.052	0.034
IGT: block 3	0.020	0.008	-0.263 [†]	-0.075	0.020	0.062	0.011	0.047
IGT: block 4	0.005	0.075	-0.345 [†]	-0.005	-0.025	0.043	-0.081	0.017
IGT: block 5	-0.258 [†]	-0.096	-0.224	-0.071	-0.181	0.039	0.029	-0.107
IGT: total	-0.044	0.034	-0.280 [†]	-0.035	-0.074	0.071	-0.033	0.001
IGT: learning	-0.191	-0.077	-0.331 [†]	-0.076	-0.124	0.014	0.009	-0.090
IGT: risk	-0.130	-0.003	-0.330 [†]	-0.040	-0.127	0.048	-0.026	-0.057
WCST: trials	-0.317 [†]	0.127	-0.006	0.008	0.005	-0.063	0.042	-0.054
WCST: errors	-0.215	0.164	0.100	0.025	0.002	-0.065	-0.023	-0.090
WCST: errors perseverant	-0.253 [†]	0.189	0.089	0.028	0.025	-0.066	-0.011	-0.100
WCST: conceptual	-0.154	-0.082	-0.272 [†]	-0.109	-0.002	0.036	0.077	0.060
WCST: categories completed	0.164	-0.053	-0.126	0.002	-0.038	0.019	0.054	0.040
WCST: trials complete 1-cat.	0.005	0.065	-0.124	0.061	0.046	-0.014	-0.023	-0.091
TMT: A	-0.103	0.385 [†]	-0.062	0.061	-0.008	0.047	-0.030	0.025
TMT: B	-0.133	0.368 [†]	0.295 [†]	0.034	-0.066	0.036	-0.036	0.108
TMT: Diff	-0.118	0.282 [†]	0.389 [†]	0.015	-0.066	0.029	-0.032	0.110
Stroop: words	0.082	-0.195	0.110	0.103	-0.066	-0.027	0.053	-0.075
Stroop: colors	-0.050	-0.160	-0.035	-0.099	-0.031	-0.085	0.004	-0.085
Stroop: words-colors	-0.052	-0.340 [†]	0.067	0.024	-0.027	-0.032	-0.023	-0.008
Stroop: estimated	-0.008	-0.192	0.033	-0.032	-0.050	-0.068	0.027	-0.091
Stroop: interference	-0.065	-0.316 [†]	0.066	0.057	0.000	0.006	-0.050	0.055
WMS-III: direct	0.041	-0.271 [†]	-0.232	-0.289 [†]	-0.026	-0.134	0.021	-0.040
WMS-III: direct-span	0.026	-0.151	-0.103	-0.190	-0.036	-0.128	0.009	-0.067
WMS-III: inverse	0.028	-0.061	-0.111	-0.207	-0.075	-0.064	0.041	0.002
WMS-III: inverse-span	0.052	-0.172	-0.060	-0.195	-0.063	-0.070	0.049	0.031
WMS-III: total	0.040	-0.195	-0.200	-0.289 [†]	-0.056	-0.110	0.035	-0.021
WAIS: vocabulary	-0.237	-0.003	0.008	0.046	0.050	-0.014	0.012	0.054

Note. ¹ Partial correlation adjusted by sex, age, and BMI. ² Partial correlation adjusted by sex, age, BMI and education level. LEAP2: liver enriched antimicrobial peptide 2. SCL-90R: Symptom Checklist-90-Revised. GSI: Global Severity Index. PST: Positive Symptom Total. PSDI: Positive Symptom Distress Index. UPPS-P: Impulsive Behavior Scale. IGT: Iowa Gambling Test. WCST: Wisconsin Card Sorting Test. TMT: Trail Making Test. WMS-III: Wechsler Memory Scale Third Edition. WAIS: Wechsler Adult Intelligence Scale. [†] Bold: effect size into the range mild-moderate ($|R| > 0.24$) to high-large ($|R| > 0.37$).