Supplementary Materials: Long-Term Occurrence of Deoxynivalenol in Feed and Feed Raw Materials with a Special Focus on South Korea

Juhee Park, Hansub Chang, Dongho Kim, Soohyun Chung and Chan Lee

Table S1. Description of compound feeds for cattle.

| Feed Type | Period for feeding | |
|------------------------|---|--|
| Early breeding calves | Birth to 3 months | |
| Middle breeding calves | 3 to 6 months | |
| Late breeding calves | 6 to 12 months | |
| Sire or breeding bull | After 13 months | |
| Gestating beef | 13 months to farrowing | |
| Lactating beef | Farrowing to 3 months after birth | |
| Early beef calves | Before 3 months | |
| Middle beef calves | After 3 months to 250 kg of body weight | |
| Early beef cattle | 250 kg to 400 kg of body weight | |
| Middle beef cattle | 400 kg to 500 kg of body weight | |
| Late beef cattle | Over 500 kg of body weight | |

Table S2. Description of compound feeds for swine.

| Feed Type | Period for feeding | |
|------------------------|---|--|
| Sucking piglets | Before lactating | |
| Weanling piglets | Over 5 kg of body weight or after lactating to 20 kg of body weight | |
| Early growing pigs | 20 kg to 50 kg of body weight | |
| Late growing pigs | 50 kg to 80 kg of body weight | |
| Growing pigs | 50 kg or 80 kg of body weight to before the 15th for shipment | |
| Growing-finishing pigs | Before the 15th for shipment to shipment | |
| Sire, boar | Over 25 kg of body weight | |
| Gilts | 25 kg of body weight to before pregnant | |
| Gestating sows | Gestating period | |
| Lactating sows | Lactating period | |

Table S3. Description of compound feeds for poultry.

| Feed Type | Period for feeding |
|-----------------------|--|
| Early layer chicks | Before 6 to 10 weeks |
| Middle layer chicks | 6 to 12 weeks or 2 weeks before laying |
| Late layer chicks | 12 weeks to 2 weeks before laying |
| Early broiler chicks | 3 weeks or before 6 weeks |
| Middle broiler chicks | 3 weeks or 6 weeks to before laying |
| Before laying | 2 weeks before laying to laying commencement |
| Early laying | Laying commencement to 40 weeks |
| Middle laying | 40 weeks to 65 weeks |
| Late laying | After 65 weeks |
| Breeding broiler | Layer breeder or broiler breeder |
| Early broiler | Before 3 weeks |
| Middle broiler | 3 weeks to before the 7-10th before shipment |
| Finishing broiler | After the 7-10th before shipment to shipment |

Table S4. Description of compound feeds for dairy cows.

| Feed Type | Period for feeding | |
|-------------------------------|--|--|
| Early dairy calves | Birth to 3 months | |
| Middle dairy calves | 3 months to 6 months | |
| Late dairy calves | 6 months to before pregnancy | |
| Gestating dairy cows | Pregnancy to 2 months after farrowing | |
| Dairy sires | After 13 months | |
| Dairy cows in early lactation | After farrowing to lactation 3 months (31 to 40 kg of milk production) | |
| Dairy cows in mid lactation | Lactation 3 months to 6 months (21–30 kg of milk production) | |
| Dairy cows in late lactation | Lactation 6 months to dry period (11–20 kg of milk production) | |
| Dairy cow on dry | Dry period (Less than 10 kg of milk production) | |
| High yielding dairy cows | Over 40 kg of milk production | |

Table S5. Description of feed ingredients.

| Class | Feed type |
|---------------------------|-------------------------|
| Caraina a | Grains |
| Grains | Grain products |
| | Corn gluten feed |
| | Soybean hull |
| | Wheat shorts |
| Grain by-products (Bran) | Cotton seeds hull |
| | Wheat bran |
| | Corn bran |
| | Other grain by-products |
| | Soybean meal |
| | Wheat gluten |
| | Corn gluten meal |
| Moal (Vogatable proteins) | Corn germ meal |
| Meal (Vegetable proteins) | Distillers dried grains |
| | Coffee meal |
| | Palm oil meal |
| | Other meal |
| Fibrous feed | Fibrous feed |
| Food by-products | Food by-products |
| Beans | Beans |
| Seed nuts | Seed nuts |
| Mixed formulation | Mixed formulation |