

Supplementary Materials: Removal of Microcystin-LR by a Novel Native Effective Bacterial Community Designated as YFMCD4 Isolated from Lake Taihu

Fei Yang ^{1,2,3,*}, Jian Guo ¹, Feiyu Huang ¹, Isaac Yaw Massey ¹, Ruixue Huang ^{1,*}, Yunhui Li ², Cong Wen ¹, Ping Ding ¹, Weiming Zeng ⁴ and Geyu Liang ²

¹ Department of Occupational and Environmental Health, Xiangya School of Public Health, Central South University, 110 Xiangya Road, Changsha 410078, China; guojianph@csu.edu.cn (J.G.); huangfeiyu@csu.edu.cn (F.H.); robert@csu.edu.cn (I.Y.M.); wencong941017@csu.edu.cn (C.W.); yixp176911007@csu.edu.cn (P.D.)

² Key Laboratory of Environmental Medicine Engineering, Ministry of Education, School of Public Health Southeast University, Nanjing 210009, China; yhli@seu.edu.cn (Y.L.); gyliang@seu.edu.cn (G.L.)

³ Key laboratory of Hunan Province for Water Environment and Agriculture Product Safety, Central South University, Changsha 410083, China;

⁴ Key Laboratory of Biometallurgy, Ministry of Education, School of Minerals Processing and Bioengineering, Central South University, Changsha 410083, China; hyacinth_hai@csu.edu.cn (W.Z.)

* Correspondence: phfyang@csu.edu.cn (F.Y.); zhengshl@csu.edu.cn (R.H.); Tel./Fax: +86-731-84805460 (F.Y. & R.H.)




Figure S1. The degradation activities of MC-LR by YFMCD4-1 and YFMCD4-2.