

Donor-Type Nickel–Dithiolene Complexes Fused with Bulky Cycloalkane Substituents and Their Application in Molecular Conductors

Kazuya Kubo ^{1,*}, Mamoru Sadahiro ¹, Sonomi Arata ¹, Norihisa Hoshino ², Tomofumi Kadoya ¹, Tomoyuki Akutagawa ², Reizo Kato ³ and Jun-ichi Yamada ¹

¹ Graduate School of Science, University of Hyogo, 3-2-1, Kouto, Kamigori-cho, Akou-gun, Hyogo 678-1297, Japan; g1127sm@gmail.com (M.S.); rinrin520sono48@gmail.com (S.A.); kadoya.t@sci.u-hyogo.ac.jp (T.K.); yamada@sci.u-hyogo.ac.jp (J.-i.Y.)

² Institute of Multidisciplinary Research for Advanced Materials, Tohoku University, 2-1-1 Katahira, Aoba-ku, Sendai, Miyagi 980-8577, Japan; norihisa.hoshino.b5@tohoku.ac.jp (N.H.); akutagawa@tohoku.ac.jp (T.A.)

³ RIKEN, 2-1 Hirosawa, Wako-shi, Saitama 351-0023, Japan; reizo@riken.jp

* Correspondence: kubo@sci.u-hyogo.ac.jp; Tel.: +81-791-58-0163

Supporting Information




Figure S1. Molecular structure of **10** along vertical and parallel directions with atom labels.




Figure S2. Crystal structure of 10 along (a) the *a*, (b) *b*, and (c) *c* axes. Hydrogen atoms are omitted for clarity. Dashed lines indicate sulfur-sulfur distances shorter than the sum of the van der Waals radius (< 3.70 Å).




Figure S3. Molecular arrangement of the [Ni(dddt)₂] moiety of the donor molecules in Crystal **9** viewed along the end-on direction.