

Supplementary Materials

Mold Fungal Resistance of Loose-Fill Thermal Insulation Materials Based on Processed Wheat Straw, Corn Stalk and Reed

Ramunas Tupciauskas ^{1,*}, Zigmunds Orlovskis ^{2,3}, Karlis Trevors Blums ^{2,3}, Janis Liepins ², Andris Berzins ^{1,4}, Gunars Pavlovics ¹ and Martins Andzs ¹

¹ Laboratory of Biorefinery, Latvian State Institute of Wood Chemistry, Dzerbenes 27, 1006 Riga, Latvia; andris.berzins@kki.lv (A.B.); pavlovichs@inbox.lv (G.P.); martins.andzs@kki.lv (M.A.)

² Faculty of Biology, University of Latvia, Jelgavas 1, 1004 Riga, Latvia; zigmunds.orlovskis@lu.lv (Z.O.); ktblums@gmail.com (K.T.B.); janis.liepins@lu.lv (J.L.)

³ Latvian Biomedical Research and Study Centre, Ratsupites 1 k-1, 1067 Riga, Latvia

⁴ Faculty of Forest and Environmental Sciences, Latvia University of Life Sciences and Technologies, Akademijas 11, 3001 Jelgava, Latvia

* Correspondence: ramunas.tupciauskas@kki.lv




Figure S1. Microscopical surface view (25 \times , scale bar 1 mm) of WS-raw samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S2. Microscopical surface view (25 \times , scale bar 1 mm) of WS-TMP samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S3. Microscopical surface view (25 \times , scale bar 1 mm) of WS-SE samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S4. Microscopical surface view (25 \times , scale bar 1 mm) of WS-SE* samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S5. Microscopical surface view (25 \times , scale bar 1 mm) of Corn-raw samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S6. Microscopical surface view (25 \times , scale bar 1 mm) of Corn-TMP samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H_2O).




Figure S7. Microscopical surface view (25 \times , scale bar 1 mm) of Corn-SE samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H_2O).




Figure S8. Microscopical surface view (25 \times , scale bar 1 mm) of Corn-SE* samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S9. Microscopical surface view (25 \times , scale bar 1 mm) of Reed-raw samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S10. Microscopical surface view (25 \times , scale bar 1 mm) of Reed-TMP samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S11. Microscopical surface view (25 \times , scale bar 1 mm) of Reed-SE samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).




Figure S12. Microscopical surface view (25 \times , scale bar 1 mm) of Reed-SE* samples after 4 weeks incubation colonized with: (a) *T. viride*, (b) *C. globulosum*, (c) *P. variotii*, (d) *P. pinophilum*, (e) *A. niger*, (f) control (H₂O).