



Figure S1. Precipitation in the location of the experiments for the cropping periods of the 2019/20 (i.e., Year 1) and 2020/21 (i.e., Year 2) seasons.



Figure S2. Symptoms of phytotoxicity after applying *cinnamon* extract at 3 mL/L.



Figure S3. Tomato fruits of the varieties used. Left: “cv. Caniles” (GH3); right: “Valenciano type” (GH1 + GH2).

Table S1. List of fungicides analysed in the formulations.

Azoxystrobin	Fenbuconazole	Mepanipyrim	Thiophanate-methyl
Boscalid (Nicobifen)	Fenhexamid	Metiram	Thiram
Bromuconazole I	Fenpropidin	Metrafenona	Triazoxide
Bromuconazole II	Fenpropimorph	Pencycuron	Trifloxystrobin
Bupirimate	Ferbam	Prochloraz	Triflumizol
Carbendazim (Azole)	Flutriafol	Propamocarb	Triticonazole
Carboxin	Fuberidazole	Propineb	Zinc thiazole
Difenoconazol	Imazalil	Pyraclostrobin	Zineb
Dimethomorph(E)	Iprovalicarb	Spiroxamine	Ziram
Dimoxystrobin	Mancozeb	Tebuconazole	Zoxamida
Dodine	Maneb	Thiabendazole	