

Table S2: Difference in percentage [%] of partially matured (CD83+CD1a+HLA-DR+) dendritic cells after stimulation with bacterial lysates between individual patients.

Difference in percentage [%] of partially matured dendritic cells in lysate-stimulated cultures.															
	χ^2 ANOVA = 18.53 p<0.18379														
	Pa 1	Pa 2	Pa 3	Pa 4	Pa 5	Pa 6	Pa 7	Pa 8	Pa 9	Pa 10	Pa 11	Pa 12	Pa 13	Pa 14	Pa 15
Pa 1	-	NS	NS	NS	NS	NS	NS								
Pa 2	NS	-	NS	NS	NS	NS	NS	NS							
Pa 3	NS	NS	-	NS	NS	NS	NS	NS	NS						
Pa 4	NS	NS	NS	-	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Pa 5	NS	NS	NS	NS	-	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS
Pa 6	NS	NS	NS	NS	NS	-	NS	NS	NS	NS	NS	NS	NS	NS	NS
Pa 7	NS	NS	NS	NS	NS	NS	-	NS	NS	NS	NS	NS	NS	NS	NS
Pa 8	NS	NS	NS	NS	NS	NS	NS	-	NS	NS	NS	NS	NS	NS	NS
Pa 9	NS	NS	NS	NS	NS	NS	NS	NS	-	NS	NS	NS	NS	NS	NS
Pa 10	NS	NS	NS	NS	NS	NS	NS	NS	NS	-	NS	NS	NS	NS	NS
Pa 11	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	-	NS	NS	NS	NS
Pa 12	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	-	NS	NS	NS
Pa 13	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	-	NS	NS
Pa 14	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	-	NS
Pa 15	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	-
No.	Pa 1	Pa 2	Pa 3	Pa 4	Pa 5	Pa 6	Pa 7	Pa 8	Pa 9	Pa 10	Pa 11	Pa 12	Pa 13	Pa 14	Pa 15
median	0.00	0.21	0.00	0.00	0.1	0.09	0.00	0.07	0.00	0.00	0.00	0.00	0.00	0.1	0.00
IQR	0.06	0.46	0.00	0.13	0.17	0.15	0.13	0.28	0.1	0.06	0.23	0.14	0.00	0.2	0.00