

Agrobacterium tumefaciens-mediated genetic transformation of the ect-endomycorrhizal fungus *Terfezia boudieri*

Lakkakula Satish ^{1,2}, Madhu Kamle ^{1,4}, Guy Keren ¹, Galit Yehezkel ¹, Ze'ev Barak ³, Varda Kagan-Zur ³, Ariel Kushmaro ² and Yaron Sitrit ^{1*}

¹ The Albert Katz International School for Desert Studies, The Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Beer Sheva - 84105, Israel. lsatish@post.bgu.ac.il (LS); guyguy123@gmail.com (GK); gality@bgu.ac.il (GY); sitrit@bgu.ac.il (YS)

² Avram and Stella Goldstein-Goren Department of Biotechnology Engineering and The Ilse Katz Center for Meso and Nanoscale Science and Technology, Ben-Gurion University of the Negev, Beer Sheva - 84105, Israel; Israel. lsatish@post.bgu.ac.il (LS); arielkus@bgu.ac.il (AK)

³ Department of Life Sciences, Ben-Gurion University of the Negev, Beer Sheva - 84105, Israel. barakz@post.bgu.ac.il (ZB); zur@bgu.ac.il (VK)

⁴ Department of Forestry, North Eastern Regional Institute of Science & Technology, Nirjuli, Arunachal Pradesh - 791109, India (current address). madhu.kamle18@gmail.com (MK)

* Correspondence: sitrit@bgu.ac.il

Supplementary figures




Figure S1. Flow chart of the transformation protocol for the desert truffle *T. boudieri*.




Figure S2. *T. boudieri* sensitivity to hygromycin and geneticin. Mycelial growth was determined after 2 weeks on Fontana medium supplemented with different concentrations of hygromycin (a) and geneticin (b). Values are mean $\pm$ SE ($n=3$), letters denote significant differences $p=0.05$ using Duncan's multiple range test.