



Supplementary Materials

Potential of ARIMA-MLP, ARIMA-SVM, DT, and CatBoost for PM_{2.5} and PM₁₀ prediction in Bangladesh

Shihab Ahmad Shahriar ¹, Imrul Kayes ¹, Kamrul Hasan ¹, Mahadi Hasan ¹, Rashik Islam ², Norrimi Rosaida Awang³, Zulhazman Humzah³, Aweng Eh Rak³ and Mohammed Abdus Salam ^{1,*}

- Department of Environmental Science and Disaster Management, Noakhali Science and Technology University, Noakhali 3814, Bangladesh; shihab0212@gmail.com (S.A.S); kamrulh9560@gmail.com (K.H.); ikayes1@lakeheadu.ca (I.K.); mahadihasan.nstu10@gmail.com (M.H.)
- ² Department of Computer Science and Media, Beuth University of Applied Sciences, Berlin, Germany; s7688@beuth-hochschule.de
- ³ Faculty of Earth Science, University Malaysia Kelantan, Jeli Campus, 17600 Jeli, Kelantan, Malaysia; norrimi.a@umk.edu.my, zulhazman@umk.edu.my, aweng@umk.edu.my

List of figures

Figure S1. Monthly and daily pattern of particulate matters across the air monitoring stations from 2013-2019.

Figure S2. Season-wise correlation among the meteorological variables and atmospheric pollutants in Dhaka, Narayanganj, and Gazipur.

Figure S3. ACF and PACF plot of daily PM2.5 concentration at the air monitoring stations

Figure S4. Residual plot of the models

Figure S5. Diagnostic test of residuals from the models (Normal Q-Q plot)

Figure S6. Variable importance for predicting PM2.5 across A. Dhaka, B. Narayanganj, and C. Gazipur

^{*}Correspondence: s_salam1978@yahoo.com/masalam.esdm@nstu.edu.bd; Tel.: +880-191-763-5348








Figure S1. Monthly and daily pattern of particulate matters across the air monitoring stations from 2013-2019








Figure S2. Season-wise correlation among the meteorological variables and atmospheric pollutants in Dhaka, Narayanganj, and Gazipur.










Figure S3. ACF and PACF plot of daily PM2.5 concentration at the air monitoring stations













Figure S4. Residual plot of the models






Normal Q-Q Plot of residuals (Dhaka)



Normal Q-Q Plot of residuals (Narayanganj)



Normal Q-Q Plot of residuals (Gazipur)




Figure S5. Diagnostic test of residuals from the models (Normal Q-Q plot)








Figure S6. Variable importance for predicting PM2.5 across A. Dhaka, B. Narayanganj, and C. Gazipur