

Terminal Value Problem for Differential Equations with Hilfer–Katugampola Fractional Derivative

Mouffak Benchohra ^{1,2}, Soufyane Bouriah ³  and Juan J. Nieto ^{4,*} 

¹ Laboratory of Mathematics, Djillali Liabes University of Sidi Bel-Abbes, P.O. Box 89, Sidi Bel Abbes 22000, Algeria; benchohra@yahoo.com

² Department of Mathematics, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

³ Department of Mathematics, Faculty of Exact Sciences and Informatics, Hassiba Benbouali University, P.O. Box 151, Chlef 02000, Algeria; s.bouriah@univ-chlef.dz or Bouriahsoufiane@yahoo.fr

⁴ Departamento de Estatística, Análise Matemática e Optimización, Instituto de Matemáticas, Universidade de Santiago de Compostela, 15705 Santiago de Compostela, Spain

* Correspondence: juan jose.nieto.roig@usc.es

Received: 11 April 2019; Accepted: 6 May 2019; Published: 15 May 2019



Abstract: We present in this work the existence results and uniqueness of solutions for a class of boundary value problems of terminal type for fractional differential equations with the Hilfer–Katugampola fractional derivative. The reasoning is mainly based upon different types of classical fixed point theory such as the Banach contraction principle and Krasnoselskii’s fixed point theorem. We illustrate our main findings, with a particular case example included to show the applicability of our outcomes.

Keywords: Hilfer–Katugampola fractional derivative; terminal value problem; existence; uniqueness; fixed point

MSC: AMS (MOS) Subject Classifications: 26A33

1. Introduction

Recently, by means of different tools from nonlinear analysis, many classes of differential equations with Caputo fractional derivative have extensively been studied in books [1–5] and in some papers, for example, [6–11]. In order to solve fractional differential equations, we mention the works [12,13] where the authors propose and prove the equivalence between an initial value problem and the Volterra integral equation.

We consider a new fractional derivative which interpolates the Hilfer, Hilfer–Hadamard, Riemann–Liouville, Hadamard, Caputo, Caputo–Hadamard, generalized and Caputo-type fractional derivatives, as well as the Weyl and Liouville fractional derivatives for particular cases of integration extremes. for more details, see [14–21] and the references therein.

It is well known [22] that the comparison principle for initial value problems of ordinary differential equations is a very useful tool in the study of qualitative and quantitative theory. Recently, attempts have been made to study the corresponding comparison principle for terminal value problems (TVP) [23].

Motivated by the works above, we establish in this paper existence and uniqueness results to the terminal value problem of the following Hilfer–Katugampola type fractional differential equation:

$$\left({}^{\rho}D_{a+}^{\alpha,\beta}y\right)(t) = f\left(t, y(t), \left({}^{\rho}D_{a+}^{\alpha,\beta}y\right)(t)\right), \text{ for each } t \in (a, T], \quad a > 0 \quad (1)$$

$$y(T) = c \in \mathbb{R} \quad (2)$$

where ${}^{\rho}D_{a+}^{\alpha,\beta}$ is the Hilfer–Katugampola fractional derivative (to be defined below) of order $\alpha \in (0, 1)$ and type $\beta \in [0, 1]$ and $f : (a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a given function. To our knowledge, no papers on terminal value problem for implicit fractional differential equations exist in the literature, in particular for those involving the Hilfer–Katugampola fractional derivative.

This paper is organized as follows. In Section 2, some notations are introduced and we recall some concepts of preliminaries about Hilfer–Katugampola fractional derivative. In Section 3, two results for Equations (1) and (2) are presented: The first one is based on the Banach contraction principle, the second one on Krasnoselskii’s fixed point theorem. Finally, in Section 4, we give an example to show the applicability of our main results.

2. Preliminaries

In this part, we present notations and definitions that we will use throughout this paper. Let $0 < a < T, J = [a, T]$. By $C(J, \mathbb{R})$ we denote the Banach space of all continuous functions from J into \mathbb{R} with the norm:

$$\|y\|_{\infty} = \sup\{|y(t)| : t \in J\}$$

We consider the weighted spaces of continuous functions:

$$C_{\gamma,\rho}(J) = \left\{ y : (a, T] \rightarrow \mathbb{R} : \left(\frac{t^{\rho} - a^{\rho}}{\rho} \right)^{\gamma} y(t) \in C(J, \mathbb{R}) \right\}, \quad 0 \leq \gamma < 1$$

and:

$$\begin{aligned} C_{\gamma,\rho}^n(J) &= \left\{ y \in C^{n-1}(J) : y^{(n)} \in C_{\gamma,\rho}(J) \right\}, \quad n \in \mathbb{N}, \\ C_{\gamma,\rho}^0(J) &= C_{\gamma,\rho}(J) \end{aligned}$$

with the norms:

$$\|y\|_{C_{\gamma,\rho}} = \sup_{t \in J} \left| \left(\frac{t^{\rho} - a^{\rho}}{\rho} \right)^{\gamma} y(t) \right|$$

and:

$$\|y\|_{C_{\gamma,\rho}^n} = \sum_{k=0}^{n-1} \|y^{(k)}\|_{\infty} + \|y^{(n)}\|_{C_{\gamma,\rho}}$$

Consider the space $X_c^p(a, b)$, ($c \in \mathbb{R}$, $1 \leq p \leq \infty$) of those complex-valued Lebesgue measurable functions f on $[a, b]$ for which $\|f\|_{X_c^p} < \infty$, where the norm is defined by:

$$\|f\|_{X_c^p} = \left(\int_a^b |t^c f(t)|^p \frac{dt}{t} \right)^{\frac{1}{p}}, \quad (1 \leq p < \infty, c \in \mathbb{R})$$

In particular, when $c = \frac{1}{p}$, the space $X_c^p(a, b)$ coincides with the $L_p(a, b)$ space: $X_{\frac{1}{p}}^p(a, b) = L_p(a, b)$.

Definition 1 ([16]). (Katugampola fractional integral).

Let $\alpha \in \mathbb{R}_+, c \in \mathbb{R}$ and $g \in X_c^p(a, b)$. The Katugampola fractional integral of order α is defined by:

$$({}^{\rho}I_{a+}^{\alpha} g)(t) = \int_a^t s^{\rho-1} \left(\frac{t^{\rho} - s^{\rho}}{\rho} \right)^{\alpha-1} \frac{g(s)}{\Gamma(\alpha)} ds, \quad t > a, \rho > 0$$

where $\Gamma(\cdot)$ is the Euler gamma function defined by: $\Gamma(\alpha) = \int_0^{\infty} t^{\alpha-1} e^{-t} dt$, $\alpha > 0$.

Definition 2 ([16]). (Katugampola fractional derivative).

Let $\alpha \in \mathbb{R}_+ \setminus \mathbb{N}$ and $\rho > 0$. The Katugampola fractional derivative ${}^\rho D_{a^+}^\alpha$ of order α is defined by:

$$\begin{aligned} ({}^\rho D_{a^+}^\alpha g)(t) &= \delta_\rho^n ({}^\rho I_{a^+}^{n-\alpha} g)(t) \\ &= \left(t^{1-\rho} \frac{d}{dt} \right)^n \int_a^t s^{\rho-1} \left(\frac{t^\rho - s^\rho}{\rho} \right)^{n-\alpha-1} \frac{g(s)}{\Gamma(n-\alpha)} ds, \quad t > a, \rho > 0 \end{aligned}$$

where $n = [\alpha] + 1$ and $\delta_\rho^n = \left(t^{1-\rho} \frac{d}{dt} \right)^n$.

Lemma 1 ([24]). Let $\alpha > 0$, and $0 \leq \gamma < 1$. Then, ${}^\rho I_{a^+}^\alpha$ is bounded from $C_{\gamma,\rho}(J)$ into $C_{\gamma,\rho}(J)$.

Lemma 2 ([24]). Let $0 < a < T < \infty, \alpha > 0, 0 \leq \gamma < 1$ and $y \in C_{\gamma,\rho}(J)$. If $\alpha > \gamma$, then ${}^\rho I_{a^+}^\alpha y$ is continuous on J and

$$({}^\rho I_{a^+}^\alpha y)(a) = \lim_{t \rightarrow a^+} ({}^\rho I_{a^+}^\alpha y)(t) = 0$$

Lemma 3 ([12]). Let $x > a$. Then, for $\alpha \geq 0$ and $\beta > 0$, we have:

$$\begin{aligned} \left[{}^\rho I_{a^+}^\alpha \left(\frac{s^\rho - a^\rho}{\rho} \right)^{\beta-1} \right](t) &= \frac{\Gamma(\beta)}{\Gamma(\alpha + \beta)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\alpha+\beta-1} \\ \left[{}^\rho D_{a^+}^\alpha \left(\frac{s^\rho - a^\rho}{\rho} \right)^{\alpha-1} \right](t) &= 0, \quad 0 < \alpha < 1 \end{aligned}$$

Lemma 4 ([24]). Let $\alpha > 0, 0 \leq \gamma < 1$ and $g \in C_\gamma[a, b]$. Then:

$$({}^\rho D_{a^+}^\alpha {}^\rho I_{a^+}^\alpha g)(t) = g(t), \quad \text{for all } t \in (a, b]$$

Lemma 5 ([24]). Let $0 < \alpha < 1, 0 \leq \gamma < 1$. If $g \in C_{\gamma,\rho}[a, b]$ and ${}^\rho I_{a^+}^{1-\alpha} g \in C_{\gamma,\rho}^1[a, b]$, then:

$$({}^\rho I_{a^+}^\alpha {}^\rho D_{a^+}^\alpha g)(t) = g(t) - \frac{({}^\rho I_{a^+}^{1-\alpha} g)(a)}{\Gamma(\alpha)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\alpha-1}, \quad \text{for all } t \in (a, b]$$

Definition 3 ([24]). Let order α and type β satisfy $n - 1 < \alpha < n$ and $0 \leq \beta \leq 1$, with $n \in \mathbb{N}$. The Hilfer–Katugampola fractional derivative to t , with $\rho > 0$ of a function $g \in C_{1-\gamma,\rho}[a, b]$, is defined by:

$$\begin{aligned} ({}^\rho D_{a^+}^{\alpha,\beta} g)(t) &= \left({}^\rho I_{a^+}^{\beta(n-\alpha)} \left(t^{\rho-1} \frac{d}{dt} \right)^n {}^\rho I_{a^+}^{(1-\beta)(n-\alpha)} g \right)(t) \\ &= \left({}^\rho I_{a^+}^{\beta(n-\alpha)} \delta_\rho^n {}^\rho I_{a^+}^{(1-\beta)(n-\alpha)} g \right)(t) \end{aligned}$$

In this paper we consider the case $n = 1$ only, because $0 < \alpha < 1$.

Property 1 ([24]). The operator ${}^\rho D_{a^+}^{\alpha,\beta}$ can be written as:

$${}^\rho D_{a^+}^{\alpha,\beta} = {}^\rho I_{a^+}^{\beta(1-\alpha)} \delta_\rho {}^\rho I_{a^+}^{1-\gamma} = {}^\rho I_{a^+}^{\beta(1-\alpha)} {}^\rho D_{a^+}^\gamma, \quad \gamma = \alpha + \beta - \alpha\beta$$

Property 2. The fractional derivative ${}^\rho D_{a^+}^{\alpha,\beta}$ is an interpolator of the following fractional derivatives: Hilfer ($\rho \rightarrow 1$) [14], Hilfer–Hadamard ($\rho \rightarrow 0^+$) [25], generalized ($\beta = 0$) [16], Caputo-type ($\beta = 1$), Riemann–Liouville ($\beta = 0, \rho \rightarrow 1$) [17], Hadamard ($\beta = 0, \rho \rightarrow 0^+$) [17], Caputo ($\beta = 1, \rho \rightarrow 1$) [17], Caputo–Hadamard ($\beta = 1, \rho \rightarrow 0^+$) [21], Liouville ($\beta = 0, \rho \rightarrow 1, a = 0$) [17] and Weyl ($\beta = 0, \rho \rightarrow 1, a = -\infty$) [15].

Definition 4. We consider the following parameters α, β, γ satisfying:

$$\gamma = \alpha + \beta - \alpha\beta, \quad 0 < \alpha, \beta, \gamma < 1.$$

Thus, we define the spaces:

$$C_{1-\gamma, \rho}^{\alpha, \beta}(J) = \{y \in C_{1-\gamma, \rho}(J), {}^{\rho}D_{a+}^{\alpha, \beta}y \in C_{1-\gamma, \rho}(J)\}$$

and:

$$C_{1-\gamma, \rho}^{\gamma}(J) = \{y \in C_{1-\gamma, \rho}(J), {}^{\rho}D_{a+}^{\gamma}y \in C_{1-\gamma, \rho}(J)\}$$

Since ${}^{\rho}D_{a+}^{\alpha, \beta}y = {}^{\rho}I_{a+}^{\gamma(1-\alpha)} {}^{\rho}D_{a+}^{\gamma}y$, it follows from Lemma 1 that:

$$C_{1-\gamma, \rho}^{\gamma}(J) \subset C_{1-\gamma, \rho}^{\alpha, \beta}(J) \subset C_{1-\gamma, \rho}(J)$$

Lemma 6 ([24]). Let $0 < \alpha < 1, 0 \leq \beta \leq 1$ and $\gamma = \alpha + \beta - \alpha\beta$. If $y \in C_{1-\gamma, \rho}^{\gamma}(J)$, then:

$${}^{\rho}I_{a+}^{\gamma} {}^{\rho}D_{a+}^{\gamma}y = {}^{\rho}I_{a+}^{\alpha} {}^{\rho}D_{a+}^{\alpha, \beta}y$$

and:

$${}^{\rho}D_{a+}^{\gamma} {}^{\rho}I_{a+}^{\alpha}y = {}^{\rho}D_{a+}^{\beta(1-\alpha)}y$$

Theorem 1 ([26]). ($PC_{1-\gamma}$ type Arzela–Ascoli Theorem). Let $A \subset PC_{1-\gamma}(J, \mathbb{R})$. A is relatively compact (i.e., \overline{A} is compact) if:

1. A is uniformly bounded, i.e., there exists $M > 0$ such that:

$$|f(x)| < M \text{ for every } f \in A \text{ and } x \in (t_k, t_{k+1}], k = 1, \dots, m$$

2. A is equicontinuous on $(t_k, t_{k+1}]$, i.e., for every $\epsilon > 0$, there exists $\delta > 0$ such that for each $x, \bar{x} \in (t_k, t_{k+1}]$, $|x - \bar{x}| \leq \delta$ implies $|f(x) - f(\bar{x})| \leq \epsilon$ for every $f \in A$.

Theorem 2 ([27]). (Banach's fixed point theorem). Let C be a non-empty closed subset of a Banach space E , then any contraction mapping T of C into itself has a unique fixed point.

Theorem 3 ([27]). (Krasnoselskii's fixed point theorem). Let M be a closed, convex and nonempty subset of a Banach space X , and A, B be the operators such that:

1. $Ax + By \in M$ for all $x, y \in M$
2. A is compact and continuous
3. B is a contraction mapping

Then there exists $z \in M$ such that $z = Az + Bz$.

3. Existence of Solutions

We consider the following linear fractional differential equation:

$$({}^{\rho}D_{a+}^{\alpha, \beta}y)(t) = \varphi(t), \quad t \in (a, T] \quad (3)$$

where $\varphi(\cdot) \in C_{1-\gamma, \rho}(J)$, with the terminal condition:

$$y(T) = c, \quad c \in \mathbb{R} \quad (4)$$

The following theorem shows that Equations (3) and (4) have a unique solution given by:

$$\begin{aligned} y(t) &= \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \varphi(s) ds \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \\ &+ \frac{1}{\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \varphi(s) ds \end{aligned} \quad (5)$$

Theorem 4. Let $\gamma = \alpha + \beta - \alpha\beta$, where $0 < \alpha < 1$ and $0 \leq \beta \leq 1$. If $\varphi : (a, T] \rightarrow \mathbb{R}$ is a function such that $\varphi(\cdot) \in C_{1-\gamma, \rho}(J)$, then y satisfies Equations (3) and (4) if and only if it satisfies Equation (5).

Proof. (\Rightarrow) Let $y \in C_{1-\gamma, \rho}^\gamma(J)$ be a solution of Equations (3) and (4). We prove that y is also a solution of Equation (5). From the definition of $C_{1-\gamma, \rho}^\gamma(J)$, Lemma 1, and using Definition 2, we have:

$${}^\rho I_{a+}^{1-\gamma} y \in C(J) \text{ and } {}^\rho D_{a+}^\gamma y = \delta_\rho {}^\rho I_{a+}^{1-\gamma} y \in C_{1-\gamma, \rho}(J) \quad (6)$$

By the Definition of the space $C_{1-\gamma, \rho}^\gamma(J)$, it follows that:

$${}^\rho I_{a+}^{1-\gamma} y \in C_{1-\gamma, \rho}^1(J)$$

Using Lemma 5, with $\alpha = \gamma$, we obtain:

$$({}^\rho I_{a+}^\gamma {}^\rho D_{a+}^\gamma y)(t) = y(t) - \frac{({}^\rho I_{a+}^{1-\gamma} y)(a)}{\Gamma(\gamma)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \quad (7)$$

where $t \in (a, T]$. By hypothesis, $y \in C_{1-\gamma, \rho}^\gamma(J)$, using Lemma 6 with Equation (3), we have:

$$({}^\rho I_{a+}^\gamma {}^\rho D_{a+}^\gamma y)(t) = ({}^\rho I_{a+}^\alpha {}^\rho D_{a+}^{\alpha, \beta} y)(t) = ({}^\rho I_{a+}^\alpha \varphi)(t) \quad (8)$$

Comparing Equations (7) and (8), we see that:

$$y(t) = \frac{({}^\rho I_{a+}^{1-\gamma} y)(a)}{\Gamma(\gamma)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} + ({}^\rho I_{a+}^\alpha \varphi)(t) \quad (9)$$

Using Equation (4) we obtain:

$$\begin{aligned} y(t) &= \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \varphi(s) ds \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \\ &+ \frac{1}{\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \varphi(s) ds \end{aligned}$$

with $t \in (a, b]$, that is $y(\cdot)$ satisfies Equation (5).

(\Leftarrow) Let $y \in C_{1-\gamma, \rho}^\gamma(J)$, satisfying Equation (5). We show that y also satisfies Equations (3) and (4). Apply operator ${}^\rho D_{a+}^\gamma$ on both sides of Equation (5). Then, from Lemmas 3 and 6 we get:

$$({}^\rho D_{a+}^\gamma y)(t) = ({}^\rho D_{a+}^{\beta(1-\alpha)} \varphi)(t) \quad (10)$$

By Equation (6) we have ${}^\rho D_{a+}^\gamma y \in C_{1-\gamma, \rho}(J)$; then, Equation (10) implies:

$$({}^\rho D_{a+}^\gamma y)(t) = \left(\delta_\rho {}^\rho I_{a+}^{1-\beta(1-\alpha)} \varphi \right)(t) = ({}^\rho D_{a+}^{\beta(1-\alpha)} \varphi)(t) \in C_{1-\gamma, \rho}(J) \quad (11)$$

As $\varphi(\cdot) \in C_{1-\gamma,\rho}(J)$ and from Lemma 1, it follows:

$$\left({}^\rho I_{a+}^{1-\beta(1-\alpha)} \varphi\right) \in C_{1-\gamma,\rho}(J) \quad (12)$$

From Equations (11) and (12) and by the Definition of the space $C_{1-\gamma,\rho}^n(J)$, we obtain:

$$\left({}^\rho I_{a+}^{1-\beta(1-\alpha)} \varphi\right) \in C_{1-\gamma,\rho}^1(J)$$

Applying operator ${}^\rho I_{a+}^{\beta(1-\alpha)}$ on both sides of Equation (11) and using Lemmas 2 and 5, we have:

$$\begin{aligned} \left({}^\rho I_{a+}^{\beta(1-\alpha)} {}^\rho D_{a+}^\gamma y\right)(t) &= \varphi(t) + \frac{\left({}^\rho I_{a+}^{1-\beta(1-\alpha)} \varphi(t)\right)(a)}{\Gamma(\beta(1-\alpha))} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\beta(1-\alpha)-1} \\ &= \left({}^\rho D_{a+}^{\alpha,\beta} y\right)(t) = \varphi(t) \end{aligned}$$

that is, Equation (3) holds. Clearly, if $y \in C_{1-\gamma,\rho}^\gamma(J)$ satisfies Equation (5), then it also satisfies Equation (4). \square

As a consequence of Theorem 4, we have Theorem 5.

Theorem 5. Let $\gamma = \alpha + \beta - \alpha\beta$ where $0 < \alpha < 1$ and $0 \leq \beta \leq 1$; let $f : (a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(\cdot, y(\cdot), u(\cdot)) \in C_{1-\gamma,\rho}(J)$ for any $y, u \in C_{1-\gamma,\rho}(J)$.

If $y \in C_{1-\gamma,\rho}^\gamma(J)$, then y satisfies Equations (1) and (2) if and only if y is the fixed point of the operator $N : C_{1-\gamma,\rho}(J) \rightarrow C_{1-\gamma,\rho}(J)$ defined by:

$$Ny(t) = M \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\gamma-1} + \frac{1}{\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho}\right)^{\alpha-1} s^{\rho-1} g(s) ds, \quad t \in (a, T] \quad (13)$$

where:

$$M := \left(\frac{T^\rho - a^\rho}{\rho}\right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho}\right)^{\alpha-1} s^{\rho-1} g(s) ds \right]$$

and $g : (0, T] \rightarrow \mathbb{R}$ be a function satisfying the functional equation:

$$g(t) = f(t, y(t), g(t))$$

Clearly, $g \in C_{1-\gamma,\rho}(J)$. In addition, by Lemma 1, $Ny \in C_{1-\gamma,\rho}(J)$.

Suppose that the function $f : (a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies the conditions:

(H1) The function $f : (a, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is such that:

$$f(\cdot, u(\cdot), v(\cdot)) \in C_{1-\gamma,\rho}^{\beta(1-\alpha)} \text{ for any } u, v \in C_{1-\gamma,\rho}(J)$$

(H2) There exist constants $K > 0$ and $0 < L < 1$ such that:

$$|f(t, u, v) - f(t, \bar{u}, \bar{v})| \leq K|u - \bar{u}| + L|v - \bar{v}|$$

for any $u, v, \bar{u}, \bar{v} \in \mathbb{R}$ and $t \in (a, T]$.

Now, we state and prove our existence result for Equations (1) and (2) based on Banach's fixed point.

Theorem 6. Assume (H1) and (H2) hold. If:

$$\frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1 - L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha < \frac{1}{2} \quad (14)$$

then the Equations (1) and (2) has unique solution in $C_{1-\gamma,\rho}^\gamma(J) \subset C_{1-\gamma,\rho}^{\alpha,\beta}(J)$.

Proof. The proof will be given in two steps:

Step 1: We show that the operator N defined in Equation (13) has a unique fixed point y^* in $C_{1-\gamma,\rho}(J)$. Let $y, u \in C_{1-\gamma,\rho}(J)$ and $t \in (a, T]$, then, we have:

$$\begin{aligned} & |Ny(t) - Nu(t)| \\ & \leq \frac{1}{\Gamma(\alpha)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |g(s) - h(s)| ds \\ & + \frac{1}{\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |g(s) - h(s)| ds \end{aligned}$$

where $g, h \in C_{1-\gamma,\rho}(J)$ such that:

$$\begin{aligned} g(t) &= f(t, y(t), g(t)) \\ h(t) &= f(t, u(t), h(t)) \end{aligned}$$

By (H2), we have:

$$\begin{aligned} |g(t) - h(t)| &= |f(t, y(t), g(t)) - f(t, u(t), h(t))| \\ &\leq K|y(t) - u(t)| + L|g(t) - h(t)| \end{aligned}$$

Then:

$$|g(t) - h(t)| \leq \frac{K}{1-L} |y(t) - u(t)|$$

Hence, for each $t \in (a, T]$:

$$\begin{aligned} & |Ny(t) - Nu(t)| \\ & \leq \frac{K}{(1-L)\Gamma(\alpha)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |y(s) - u(s)| ds \\ & + \frac{K}{(1-L)\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |y(s) - u(s)| ds \\ & \leq \frac{K}{(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \|y - u\|_{C_{1-\gamma,\rho}} \left(\rho I_{a^+}^\alpha \left(\frac{s^\rho - a^\rho}{\rho} \right)^{\gamma-1} \right) (T) \\ & + \frac{K}{(1-L)} \left(I_{a^+}^\alpha \left(\frac{s^\rho - a^\rho}{\rho} \right)^{\gamma-1} \right) (t) \|y - u\|_{C_{1-\gamma,\rho}} \end{aligned}$$

By Lemma 3, we have:

$$\begin{aligned} |Ny(t) - Nu(t)| &\leq \left[\frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1 - L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \right. \\ &\quad \left. + \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1 - L)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\alpha+\gamma-1} \right] \|y - u\|_{C_{1-\gamma,\rho}}, \end{aligned}$$

hence:

$$\begin{aligned}
\left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} (Ny(t) - Nu(t)) \right| &\leq \left[\frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \right. \\
&\quad \left. + \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{t^\rho - a^\rho}{\rho} \right)^\alpha \right] \|y - u\|_{C_{1-\gamma,\rho}} \\
&\leq \frac{2K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \|y - u\|_{C_{1-\gamma,\rho}},
\end{aligned}$$

which implies that:

$$\|Ny - Nu\|_{C_{1-\gamma,\rho}} \leq \frac{2K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \|y - u\|_{C_{1-\gamma,\rho}}.$$

By Equation (14), the operator N is a contraction. Hence, by Banach's contraction principle, N has a unique fixed point $y^* \in C_{1-\gamma,\rho}(J)$.

Step 2: We show that such a fixed point $y^* \in C_{1-\gamma,\rho}(J)$ is actually in $C_{1-\gamma,\rho}^\gamma(J)$.

Since y^* is the unique fixed point of operator N in $C_{1-\gamma,\rho}(J)$, then, for each $t \in (a, T]$, we have:

$$\begin{aligned}
y^*(t) &= \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} f(s, y^*(s), g(s)) ds \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \\
&\quad + {}^\rho I_{a+}^\alpha f(s, y^*(s), g(s))
\end{aligned}$$

Applying ${}^\rho D_{a+}^\gamma$ to both sides and by Lemmas 3 and 6, we have:

$$\begin{aligned}
{}^\rho D_{a+}^\gamma y^*(t) &= ({}^\rho D_{a+}^\gamma {}^\rho I_{a+}^\alpha f(s, y^*(s), g(s))) (t) \\
&= ({}^\rho D_{a+}^{\beta(1-\alpha)} f(s, y^*(s), g(s))) (t)
\end{aligned}$$

Since $\gamma \geq \alpha$, by (H1), the right hand side is in $C_{1-\gamma,\rho}(J)$ and thus ${}^\rho D_{a+}^\gamma y^* \in C_{1-\gamma,\rho}(J)$, which implies that $y^* \in C_{1-\gamma,\rho}^\gamma(J)$. As a consequence of Steps 1 and 2 together with Theorem 5, we can conclude that Equations (1) and (2) have a unique solution in $C_{1-\gamma,\rho}^\gamma(J)$. \square

We present now the second result, which is based on Krasnoselskii fixed point theorem.

Theorem 7. Assume (H1) and (H2) hold. If:

$$\frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha < 1 \quad (15)$$

then Equations (1) and (2) have at least one solution.

Proof. Consider the set:

$$B_{\eta^*} = \{y \in C_{1-\gamma,\rho}(J) : \|y\|_{C_{1-\gamma,\rho}} \leq \eta^*\}$$

where:

$$\eta^* \geq \frac{\left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{\Gamma(\gamma)f^*}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \right]}{1 - \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha}$$

and $f^* = \sup_{t \in J} |f(t, 0, 0)|$.

We define the operators P and Q on B_{η^*} by:

$$Py(t) = \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} g(s) ds \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \quad (16)$$

$$Qy(t) = \frac{1}{\Gamma(\alpha)} \int_a^t \left(\frac{t^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} g(s) ds \quad (17)$$

Then the fractional integral Equation (13) can be written as the operator equation:

$$Ny(t) = Py(t) + Qy(t), \quad y \in C_{1-\gamma, \rho}(J)$$

The proof will be given in several steps:

Step 1: We prove that $Py + Qu \in B_{\eta^*}$ for any $y, u \in B_{\eta^*}$. For operator P , multiplying both sides of Equation (16) by $\left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma}$, we have:

$$\left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} Py(t) = \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[c - \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} g(s) ds \right]$$

then:

$$\left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} Py(t) \right| \leq \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{1}{\Gamma(\alpha)} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |g(s)| ds \right] \quad (18)$$

By (H3), we have for each $t \in (a, T]$:

$$\begin{aligned} |g(t)| &= |f(t, y(t), g(t)) - f(t, 0, 0) + f(t, 0, 0)| \\ &\leq |f(t, y(t), g(t)) - f(t, 0, 0)| + |f(t, 0, 0)| \\ &\leq K|y(t)| + L|g(t)| + f^* \end{aligned}$$

Multiplying both sides of the above inequality by $\left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma}$, we get:

$$\begin{aligned} \left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} g(t) \right| &\leq \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} f^* + K \left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} y(t) \right| \\ &\quad + L \left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} g(t) \right| \\ &\leq \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} f^* + K\eta^* + L \left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} g(t) \right| \end{aligned}$$

Then, for each $t \in (a, T]$, we have:

$$\left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} g(t) \right| \leq \frac{\left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} f^* + K\eta^*}{1-L} := M \quad (19)$$

Thus, Equation (18) and Lemma 3, imply:

$$\left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} Py(t) \right| \leq \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{M\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{\alpha + \gamma - 1} \right]$$

This gives:

$$\|Py\|_{C_{1-\gamma, \rho}} \leq \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{M\Gamma(\gamma)}{\Gamma(\alpha + \gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{\alpha + \gamma - 1} \right] \quad (20)$$

Using Equation (19) and Lemma 3, we have:

$$|Q(u)(t)| \leq \left[\frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha+\gamma)} \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\alpha+\gamma-1}$$

Therefore:

$$\begin{aligned} \left| \left(\frac{t^\rho - a^\rho}{\rho} \right)^{1-\gamma} Qu(t) \right| &\leq \left[\frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha+\gamma)} \right] \left(\frac{t^\rho - a^\rho}{\rho} \right)^\alpha, \\ &\leq \frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma+\alpha} + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \end{aligned}$$

Thus:

$$\|Qu\|_{C_{1-\gamma,\rho}} \leq \frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma+\alpha} + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \quad (21)$$

Linking Equations (20) and (21), for every $y, u \in B_{\eta^*}$ we obtain:

$$\begin{aligned} \|Py + Qu\|_{C_{1-\gamma,\rho}} &\leq \max \left\{ \|Py\|_{C_{1-\gamma,\rho}}, \|Qu\|_{C_{1-\gamma,\rho}} \right\} \\ &\leq \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{M\Gamma(\gamma)}{\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{\alpha+\gamma-1} \right] \\ &= \frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma+\alpha} \\ &\quad + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha+\gamma)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha + \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} |c| \end{aligned}$$

Since:

$$\eta^* \geq \frac{\left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left[|c| + \frac{\Gamma(\gamma)f^*}{\Gamma(\alpha+\gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \right]}{1 - \frac{K\Gamma(\gamma)}{\Gamma(\alpha+\gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha}$$

we have:

$$\|Py + Qu\|_{PC_{1-\gamma,\rho}} \leq \eta^*$$

which infers that $Py + Qu \in B_{\eta^*}$.

Step 2: P is a contraction.

Let $y, u \in C_{1-\gamma,\rho}(J)$ and $t \in (a, T]$; then, we have:

$$\begin{aligned} &|Py(t) - Pu(t)| \\ &\leq \frac{1}{\Gamma(\alpha)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho} \right)^{\gamma-1} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} |g(s) - h(s)| ds \end{aligned}$$

where $g, h \in C_{1-\gamma,\rho}(J)$ such that:

$$\begin{aligned} g(t) &= f(t, y(t), g(t)) \\ h(t) &= f(t, u(t), h(t)) \end{aligned}$$

By (H2), we have:

$$\begin{aligned} |g(t) - h(t)| &= |f(t, y(t), g(t)) - f(t, u(t), h(t))| \\ &\leq K|y(t) - u(t)| + L|g(t) - h(t)| \end{aligned}$$

Then,

$$|g(t) - h(t)| \leq \frac{K}{1-L}|y(t) - u(t)|$$

Therefore, for each $t \in (a, T]$:

$$\begin{aligned} &|Py(t) - Pu(t)| \\ &\leq \frac{K}{(1-L)\Gamma(\alpha)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\gamma-1} \int_a^T \left(\frac{T^\rho - s^\rho}{\rho}\right)^{\alpha-1} s^{\rho-1} |y(s) - u(s)| ds \\ &\leq \frac{K}{(1-L)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^{1-\gamma} \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\gamma-1} \|y - u\|_{C_{1-\gamma,\rho}} \left(\rho I_{a^+}^\alpha \left(\frac{s^\rho - a^\rho}{\rho}\right)^{\gamma-1}\right)(T). \end{aligned}$$

By Lemma 3, we have:

$$|Py(t) - Pu(t)| \leq \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^\alpha \left(\frac{t^\rho - a^\rho}{\rho}\right)^{\gamma-1} \|y - u\|_{C_{1-\gamma,\rho}},$$

hence:

$$\left| \left(\frac{t^\rho - a^\rho}{\rho}\right)^{1-\gamma} (Py(t) - Pu(t)) \right| \leq \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^\alpha \|y - u\|_{C_{1-\gamma,\rho}},$$

which implies that:

$$\|Py - Pu\|_{C_{1-\gamma,\rho}} \leq \frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1-L)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^\alpha \|y - u\|_{C_{1-\gamma,\rho}}.$$

By Equation (15) the operator P is a contraction.

Step 3: Q is compact and continuous.

The continuity of Q follows from the continuity of f . Next we prove that Q is uniformly bounded on B_{η^*} .

Let any $u \in B_{\eta^*}$. Then by Equation (21) we have:

$$\|Qu\|_{PC_{1-\gamma,\rho}} \leq \frac{\Gamma(\gamma)f^*}{(1-L)\Gamma(\alpha + \gamma)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^{1-\gamma+\alpha} + \frac{K\Gamma(\gamma)\eta^*}{(1-L)\Gamma(\alpha + \gamma)} \left(\frac{T^\rho - a^\rho}{\rho}\right)^\alpha$$

This means that Q is uniformly bounded on B_{η^*} . Next, we show that QB_{η^*} is equicontinuous.

Let any $u \in B_{\eta^*}$ and $0 < a < \tau_1 < \tau_2 \leq T$. Then:

$$\begin{aligned} &\left| \left(\frac{\tau_2^\rho - a^\rho}{\rho}\right)^{1-\gamma} Q(y)(\tau_2) - \left(\frac{\tau_1^\rho - a^\rho}{\rho}\right)^{1-\gamma} Q(y)(\tau_1) \right| \leq \\ &\frac{\left(\frac{\tau_2^\rho - a^\rho}{\rho}\right)^{1-\gamma}}{\Gamma(\alpha)} \int_{\tau_1}^{\tau_2} \left(\frac{\tau_2^\rho - s^\rho}{\rho}\right)^{\alpha-1} s^{\rho-1} |g(s)| ds \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{\Gamma(\alpha)} \int_a^{\tau_1} \left| \left[\left(\frac{\tau_2^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{\tau_2^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \right. \right. \\
& \left. \left. - \left(\frac{\tau_1^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{\tau_1^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \right] \right| |g(s)| ds \\
& \leq \frac{M\Gamma(\gamma) \left(\frac{\tau_2^\rho - a^\rho}{\rho} \right)^{1-\gamma}}{\Gamma(\alpha + \gamma)} \left(\frac{\tau_2^\rho - \tau_1^\rho}{\rho} \right)^{\alpha+\gamma-1} \\
& + \frac{M}{\Gamma(\alpha)} \int_a^{\tau_1} \left| \left[\left(\frac{\tau_2^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{\tau_2^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \right. \right. \\
& \left. \left. - \left(\frac{\tau_1^\rho - a^\rho}{\rho} \right)^{1-\gamma} \left(\frac{\tau_1^\rho - s^\rho}{\rho} \right)^{\alpha-1} s^{\rho-1} \right] \right| \left(\frac{s^\rho - a^\rho}{\rho} \right)^{\gamma-1} ds
\end{aligned}$$

Note that:

$$\left| \left(\frac{\tau_2^\rho - a^\rho}{\rho} \right)^{1-\gamma} Q(y)(\tau_2) - \left(\frac{\tau_1^\rho - a^\rho}{\rho} \right)^{1-\gamma} Q(y)(\tau_1) \right| \rightarrow 0 \quad \text{as } \tau_2 \rightarrow \tau_1$$

This shows that Q is equicontinuous on J . Therefore, Q is relatively compact on B_{η^*} . By $C_{1-\gamma}$, type Arzela–Ascoli Theorem Q is compact on B_{η^*} .

As a consequence of Krasnoselskii's fixed point theorem, we conclude that N has at least a fixed point $y^* \in C_{1-\gamma,\rho}(J)$ and by the same way of the proof of Theorem 6, we can easily show that $y^* \in C_{1-\gamma,\rho}^\gamma(J)$. Using Lemma 5, we conclude that Equations (1) and (2) have at least one solution in the space $C_{1-\gamma,\rho}^\gamma(J)$. \square

4. An Example

Consider the following terminal value problem:

$${}^{\frac{1}{2}}D_{1+}^{\frac{1}{2},0} y(t) = \frac{2 + |y(t)| + \left| {}^{\frac{1}{2}}D_{0+}^{\frac{1}{2},0} y(t) \right|}{108e^{-t+3} \left(1 + |y(t)| + \left| {}^{\frac{1}{2}}D_{0+}^{\frac{1}{2},0} y(t) \right| \right)} + \frac{\ln(\sqrt{t} + 1)}{3\sqrt{\sqrt{t} - 1}}, \quad t \in (1, 2] \quad (22)$$

$$y(2) = c \in \mathbb{R} \quad (23)$$

Set:

$$f(t, u, v) = \frac{2 + u + v}{108e^{-t+3}(1 + u + v)} + \frac{\ln(\sqrt{t} + 1)}{3\sqrt{t}}, \quad t \in (1, 2], \quad u, v \in [0, +\infty)$$

We have:

$$C_{1-\gamma,\rho}^{\beta(1-\alpha)}([1, 2]) = C_{\frac{1}{2}, \frac{1}{2}}^0([1, 2]) = \left\{ h : (1, 2] \rightarrow \mathbb{R} : \sqrt{2} \left(\sqrt{t} - 1 \right)^{\frac{1}{2}} h \in C([1, 2]) \right\}$$

with $\gamma = \alpha = \rho = \frac{1}{2}$ and $\beta = 0$. Clearly, the function $f \in C_{\frac{1}{2}, \frac{1}{2}}([1, 2])$.

Hence condition (H1) is satisfied.

For each $u, \bar{u}, v, \bar{v} \in \mathbb{R}$ and $t \in (1, 2]$:

$$\begin{aligned} |f(t, u, v) - f(t, \bar{u}, \bar{v})| &\leq \frac{1}{108e^{-t+3}} (|u - \bar{u}| + |v - \bar{v}|) \\ &\leq \frac{1}{108e} (|u - \bar{u}| + |v - \bar{v}|) \end{aligned}$$

Therefore, (H2) is verified with $K = L = \frac{1}{108e}$.

The condition:

$$\frac{K\Gamma(\gamma)}{\Gamma(\alpha + \gamma)(1 - L)} \left(\frac{T^\rho - a^\rho}{\rho} \right)^\alpha \approx 0.0055 < 1$$

is satisfied with $T = 2$ and $a = 1$. It follows from Theorem 7 that Equations (22) and (23) have a solution in the space $C^{\frac{1}{2}, \frac{1}{2}}([1, 2])$.

5. Conclusions

We have provided sufficient conditions ensuring the existence and uniqueness of solutions to a class of terminal value problem for differential equations with the Hilfer–Katugampola type fractional derivative. The arguments are based on the classical Banach contraction principle, and the Krasnoselskii’s fixed point theorem. An example is included to show the applicability of our results.

Author Contributions: All authors contributed equally and significantly in writing this article. All authors read and approved the final manuscript. Conceptualization, M.B. and S.B.; Formal analysis, M.B., S.B. and J.J.N.; Investigation, M.B., S.B. and J.J.N.; Writing—original draft, M.B., S.B. and J.J.N.; Writing—review & editing, M.B., S.B. and J.J.N.; Funding, J.J.N.

Funding: This research was partially funded by the Agencia Estatal de Investigacion (AEI) of Spain under grant MTM2016-75140-P.

Acknowledgments: The research of J.J. Nieto was partially supported by the AEI of Spain under Grant MTM2016-75140-P and co-financed by the European Community fund FEDER. This paper was completed while the second author was visiting the University of Santiago de Compostela. He is grateful for the warm hospitality.

Conflicts of Interest: The authors declare that they have no competing interests concerning the publication of this article.

References

1. Abbas, S.; Benchohra, M.; Graef, J.R.; Henderson, J. *Implicit Fractional Differential and Integral Equations: Existence and Stability*; Walter de Gruyter: London, UK, 2018.
2. Abbas, S.; Benchohra, M.; N’Guérékata, G.M. *Topics in Fractional Differential Equations*; Springer-Verlag: New York, NY, USA, 2012.
3. Abbas, S.; Benchohra, M.; N’Guérékata, G.M. *Advanced Fractional Differential and Integral Equations*; Nova Science Publishers: New York, NY, USA, 2014.
4. Ahmad, B.; Alsaedi, A.; Ntouyas, S.K.; Tariboon, J. *Hadamard-type Fractional Differential Equations, Inclusions and Inequalities*; Springer: Cham, Switzerland, 2017.
5. Zhou, Y.; Wang, J.-R.; Zhang, L. *Basic Theory of Fractional Differential Equations*, 2nd ed.; World Scientific Publishing Co. Pte. Ltd.: Hackensack, NJ, USA, 2017.
6. Ahmad, B.; Ntouyas, S.K. Fractional differential inclusions with fractional separated boundary conditions. *Fract. Calc. Appl. Anal.* **2012**, *15*, 362–382. [\[CrossRef\]](#)
7. Benchohra, M.; Bouriah, S.; Darwish, M.A. Nonlinear boundary value problem for implicit differential equations of fractional order in Banach spaces. *Fixed Point Theory* **2017**, *18*, 457–470. [\[CrossRef\]](#)
8. Benchohra, M.; Bouriah, S.; Graef, J.R. Nonlinear implicit differential equations of fractional order at resonance. *Electron. J. Differ. Equ.* **2016**, *2016*, 1–10.
9. Abbas, S.; Benchohra, M.; Bouriah, S.; Nieto, J.J. Periodic solutions for nonlinear fractional differential systems. *Differ. Equ. Appl.* **2018**, *10*, 299–316. [\[CrossRef\]](#)

10. Benchohra, M.; Bouriah, S.; Henderson, J. Existence and stability results for nonlinear implicit neutral fractional differential equations with finite delay and impulses. *Commun. Appl. Nonlinear Anal.* **2015**, *22*, 46–67.
11. Benchohra, M.; Lazreg, J.E. Nonlinear fractional implicit differential equations. *Commun. Appl. Anal.* **2013**, *17*, 471–482.
12. Almeida, R.; Malinowska, A.B.; Odziejewicz, T. Fractional differential equations with dependence on the Caputo–Katugampola derivative. *J. Comput. Nonlinear Dyn.* **2016**, *11*, 061017. [[CrossRef](#)]
13. Kilbas, A.A.; Trujillo, J.J. Hadamard-type integrals as G-transforms. *Integral Transforms Spec. Funct.* **2003**, *14*, 413–427. [[CrossRef](#)]
14. Hilfer, R. *Applications of Fractional Calculus in Physics*; World Scientific: Singapore, 2000.
15. Hilfer, R. Threefold Introduction to Fractional Derivatives. In *Anomalous Transport: Foundations and Applications*; Wiley-VCH: Weinheim, Germany, 2008; p. 17.
16. Katugampola, U. A new approach to a generalized fractional integral. *Appl. Math. Comput.* **2011**, *218*, 860–865. [[CrossRef](#)]
17. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. Theory and Applications of Fractional Differential Equations. In *North-Holland Mathematics Studies*; Elsevier Science B.V.: Amsterdam, The Netherlands, 2006; Volume 204.
18. Podlubny, I. *Fractional Differential Equations*; Academic Press: San Diego, CA, USA, 1999.
19. Baleanu, D.; Diethelm, K.; Scalas, E.; Trujillo, J.J. *Fractional Calculus Models and Numerical Methods*; World Scientific Publishing: New York, NY, USA, 2012.
20. Baleanu, D.; Güvenc, Z.B.; Machado, J.A.T. *New Trends in Nanotechnology and Fractional Calculus Applications*; Springer: New York, NY, USA, 2010.
21. Gambo, Y.Y.; Jarad, F.; Baleanu, D.; Abdeljawad, T. On Caputo modification of the Hadamard fractional derivatives. *Adv. Differ. Equ.* **2014**, *2014*, 10. [[CrossRef](#)]
22. Lakshmikantham, V.; Leela, S. *Differential and Integral Inequalities*; Academic Press: New York, NY, USA, 1969; Volume I.
23. Hallam, T.G. A comparison principle for terminal value problems in ordinary differential equations. *Trans. Am. Math. Soc.* **1972**, *169*, 49–57. [[CrossRef](#)]
24. Oliveira, D.S.; de Oliveira, E.C. Hilfer–Katugampola fractional derivative. *Comput. Appl. Math.* **2018**, *37*, 3672–3690. [[CrossRef](#)]
25. Kassim, M.D.; Tatar, N.E. Well-posedness and stability for a differential problem with Hilfer–Hadamard fractional derivative. *Abstr. Appl. Anal.* **2013**, *2013*, 605029. [[CrossRef](#)]
26. Wei, W.; Xiang, X.; Peng, Y. Nonlinear impulsive integro-differential equations of mixed type and optimal controls. *Optimization* **2006**, *55*, 141–156. [[CrossRef](#)]
27. Granas, A.; Dugundji, J. *Fixed Point Theory*; Springer-Verlag: New York, NY, USA, 2003.

