

Table S1. Postoperative pathological characteristics and lymph node metastasis of final-EC patients.

Characteristic	Patients (<i>n</i>)	final-EC (<i>n</i> = 190)
FIGO stage (2009)	190	
IA		170 (89.5%)
IB		7 (3.7%)
II		6 (3.2%)
IIIA		3 (1.6%)
IIIC1		1 (0.5%)
IIIC2		3 (1.6%)
Grade	183	
1		172 (94.0%)
2		9 (4.9%)
3		2 (1.1%)
Myometrial invasion	190	
No invasion		48 (25.3%)
< 50% invasion		131 (68.9%)
≥ 50% invasion		11 (5.8%)
Histology	190	
Endometrioid		184 (96.8%)
Serous		1 (0.5%)
Mucinous		1 (0.5%)
Secretory		4 (2.1%)
LVSI	190	
Absent		180 (94.7%)
Present		10 (5.3%)
Peritoneal washes	71	
(-)		68 (95.8%)
(+)		3 (4.2%)
Cervical infiltration	190	
(-)		183 (96.3%)
(+)		7 (3.7%)
Parametrium involvement	190	
(-)		190 (100%)
(+)		0 (0%)
MELF	190	
(-)		182 (95.8%)
(+)		8 (4.2%)
Para-aortic lymph node	16	
(-)		13 (81.2%)
(+)		3 (18.8%)
Pelvic lymph node	37	
(-)		34 (91.9%)
(+)		3 (8.1%)

Data shown were number (%).

Abbreviations: final-EC, endometrial cancer diagnosed by final histopathology; LVSI, lymph-vascular space invasion; MELF, microcystic, elongated and fragmented.