

Table S1 Oligonucleotides and PCR conditions used for the molecular identification and/or characterization of the intestinal protist parasites investigated in Tengchong City, southwest China.

Enteric pathogen	Locus	Primers name	Primer (5'-3')	Amplicon size (bp)	Cycling conditions	Reference
<i>Giardia duodenalis</i>	<i>tpi</i>	AL3543	AAATIATGCCTGCTCGTCG	605	Denaturation: 94 °C for 45 s	[49]
		AL3546	CAAACCTTITCCGCAAACC		Annealing: 50 °C for 45 s	
		AL3544	CCCTTCATCGGIGGTAACCT	530	Elongation: 72 °C for 1 min	
		AL3545	GTGCCACCAACICCCGTGCC		No. cycles: 35	
<i>Cryptosporidium</i> spp.	<i>ssu-rRNA</i>	Primer F1	TTCTAGAGCTAATACATGCG	1,325	Denaturation: 94 °C for 1 min	[48]
		Primer R1	CCCATTCCCTCGAACAGGA		Annealing: 55 (1 st) or 58 (2 nd) °C for 1 min	
		Primer F2	GGAAGGGTTGTATTATTAGATAAAG	840	Elongation: 72 °C for 1 min	
		Primer R2	CTCATAGGTGCTGAAGGAGTA		No. cycles: 35	
<i>E. histolytica</i>	<i>ssu-rRNA</i>	E1	TGCTGTGATTAAAACGCT	1,947	Denaturation: 94 °C for 1 min	[50]
		E2	TTAACTATTCATCTCGG		Annealing: 47 (1 st) or 58 (2 nd) °C for 1 min	
		Eh-L	ACATTGAAAGACTTATGTAAGTA	427	Elongation: 72 °C for 1 min	
		Eh-R	CAGATCTAGAACAAATGCTTCT		No. cycles: 35	
<i>Blastocystis</i> sp.	<i>ssu-rRNA</i>	Forward	GGAGGTAGTGACAATAATC	1,100	Denaturation: 94 °C for 1 min	[47]
		Reverse	ACTAGGAATTCCCTCGTTCATG		Annealing: 54 °C for 1 min	
					Elongation: 72 °C for 1 min	
						No. cycles: 35

ssu rRNA, small subunit ribosomal RNA; *tpi*, Triose phosphate isomerase