


**Supplementary Table S1.** Susceptibility of planktonic yeast cells of *S. brasiliensis* and *S. schenckii*. Concentrations in µg/ml.

	<b>MIC<sub>50</sub> <sup>LR</sup></b>	<b>MIC<sub>50</sub></b>	<b>MIC<sub>90</sub></b>	<b>MFC</b>
<b><i>S. brasiliensis</i> ATCC 4823</b>				
Itraconazole	0.025	< 0.03 - 0.03	0.125 - 0.25	> 16
Amphotericin B	0.069	0.06 - 0.125	0.125	< 0.5
Miltefosine	0.313	0.5	2 - 4	2 - 1
<b><i>S. brasiliensis</i> ATCC 4824</b>				
Itraconazole	0.098	0.03 - 0.125	0.125 - 0.25	> 16
Amphotericin B	0.394	0.06	0.06 - 0.125	< 0.5
Miltefosine	0.959	0.25 - 0.5	0.5 - 1	1
<b><i>S. brasiliensis</i> ss245</b>				
Itraconazole	0.546	0.125 - 0.25	1 - 2	> 16
Amphotericin B	2.891	0.125	0.25	< 0.5
Miltefosine	3.051	0.25 - 1	0.5 - 2	1
<b><i>S. schenckii</i> ATCC 32286</b>				
Itraconazole	0.079	0.06 - 0.25	0.25 - 2	> 16
Amphotericin B	0.125	0.125	0.25	1 - < 0.5
Miltefosine	0.667	1	2	2

<sup>LR</sup>: MIC<sub>50</sub> calculated by Linear Regression.



**Supplementary Figure S1.** Spotting inhibitory concentrations from the MIC assay shows a potential fungicidal profile for miltefosine.