

Bacteraemia caused by probiotic strains of *Lactocaseibacillus rhamnosus* – case studies highlighting the need for careful thought before using microbes for health benefits

Agnieszka Mikucka¹, Aleksander Deptuła², Tomasz Bogiel^{1*}, Agnieszka Chmielarczyk³, Elżbieta Nurczyńska⁴, Eugenia Gospodarek-Komkowska¹

¹ Department of Microbiology Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Torun, Poland

² Department of Propaedeutics of Medicine and Infection Prevention Ludwik Rydygier Collegium Medicum in Bydgoszcz, Nicolaus Copernicus University in Torun, Poland

³ Chair of Microbiology, Jagiellonian University Medical College in Cracow, Poland

⁴ Anesthesiology and Intensive Care Unit, University Hospital No 1 in Bydgoszcz, Poland

* **Correspondence:** 9 Maria Skłodowska-Curie Street, 85-094 Bydgoszcz, e-mail: t.bogiel@cm.umk.pl (T.B.), telephone number: +48 52 585 44 80




Figure S1. PFGE typing profiles of *L. rhamnosus* obtained with SgsI restriction enzyme digestion. Analysis was done in Gel Compar software (coefficient Dice, UPGMA algorithm, UPGMA, Tol. 1%, Opt 0.5%, Min area 0.0%), 3708-7914 – the strains isolated from the patients, Lacidofil – probiotic strain *L. rhamnosus* derived from Lacidofil (LALLEMAND SAS); GG – *L. rhamnosus* strain GG isolated from Dicoflor (Bayer)