



Supplementary Materials

Table S1. *Giardia duodenalis* reference strains used in this study.

Accession Number	Isolate	Assemblage / Genotype	Host	Country	Content	Year	Reference
EU769206	Swecat171	A	cat	Sweden	Europe	2011	[1]
EU621373	JC002	A	deer	Poland	Europe	2008	[2]
EU769205	Swecat078	A	cat	Sweden	Europe	2008	[1]
GU396696	H3-001	A	human	Poland	Europe	2016	[3]
HM165227	Sweh040	A	human	Sweden	Europe	2016	[4]
KF922979	HC11	A	human	Brazil ^a	South America	2014	[5]
KF922983	HC29	A	human	Brazil ^a	South America	2014	[5]
KF922992	HC44	A	human	Brazil ^a	South America	2014	[5]
KF923000	DC01	A	dog	Brazil ^a	South America	2014	[5]
KM190682	VANC/85/UBC/2	A	human	Canada	North America	1985	[6]
KY612245	AN1.5	A	human	Brazil ^b	South America	2014	[7]
MN629930	Z1	A	dog	Iraq	Asia	2020	[8]
XM001705373	WB C6	A	culture	USA	North America	2007	[9]
GQ329671	Sweh166	A	human	Sweden	Europe	2011	[4]
EU014384	Be2	A1	beaver	Canada	North America	2011	[10]
HQ179591	46c2	A2	human	Australia	Oceania	2011	[11]
AY072724	ISSGF7	A3	human	Italy	Europe	2002	[12]
DQ090542	STS-U	A3	human	Norway	Europe	2004	[13]
DQ116612	CBHRG9	A3	wastewater	Mexico	North America	2006	[14]
EU014381	AB	A3	human	Peru	South America	1985	[15]
FJ560577	Lim2	A3	human	France	Europe	1998	[16]
FJ971408	GLT1	A3	human	Thailand	Asia	2007	[17]
FN386484	I231104	A3	wastewater	Spain	Europe	2004	[18]
KU504738	S42	A3	human	Brazil ^c	South America	2011	[19]
DQ090523	BG-Ber2	B	human	Norway	Europe	2005	[13]
AB480877	PalH9	B	human	Palestine	Asia	2006	[20]
AB618785	GH-202	B	human	Japan	Asia	2010	[21]
FJ971440	GLT2	B	human	Thailand	Asia	2007	[17]
FJ971461	GL518	B	human	Thailand	Asia	2010	[17]
FJ971482	GL121	B	human	Thailand	Asia	2007	[17]
KF922985	HC32	B	human	Brazil ^a	South America	2014	[5]
KF922993	HC45	B	human	Brazil ^a	South America	2014	[5]
KP687755	VANC/90/UBC/44	B	wastewater	Canada	North America	1990	[6]
KP687756	"VANC/90/UBC/54	B	beaver	Canada	North America	1990	[6]
KU504702	S3C1	B	human	Brazil ^c	South America	2011	[19]
KU504704	S3C3	B	human	Brazil ^c	South America	2011	[19]
KU504707	S8	B	human	Brazil ^c	South America	2011	[19]
KU504708	S9	B	human	Brazil ^c	South America	2011	[19]
KU504709	S10C3	B	human	Brazil ^c	South America	2011	[19]
KU504712	S11C1	B	human	Brazil ^c	South America	2011	[19]
KU504713	S11C2	B	human	Brazil ^c	South America	2011	[19]
KU504714	S11C3	B	human	Brazil ^c	South America	2011	[19]
KU504715	S13C2	B	human	Brazil ^c	South America	2011	[19]
KU504720	S15	B	human	Brazil ^c	South America	2011	[19]
KU504722	S16C2	B	human	Brazil ^c	South America	2011	[19]
KU504723	S16C4	B	human	Brazil ^c	South America	2011	[19]
KU504731	S24C2	B	human	Brazil ^c	South America	2011	[19]
KU504732	S24C3	B	human	Brazil ^c	South America	2011	[19]
KY612242	S8C5	B	human	Brazil ^c	South America	2011	[19]
LC508615	K6 clone1	B	human	Kenya	Africa	2013	[22]
MT542771	5	B	human	Brazil ^c	South America	2021	[23]
AY072725	Nij5	B1	human	The Netherlands	Europe	2002	[24]
AY072726	LD18	B2	human	Belgium	Europe	2002	[24]
AY072727	BAH8	B3	human	Australia	Oceania	1999	[12]
JF422719	BRAdogD15	C	dog	Brazil ^a	South America	2009	[25]

KF923019	VET01	D	dog	Brazil ^a	South America	2014	[5]
AY072729	P15	E	pig	Czech Republic	Europe	1996	[12]
AY647264	A101	F	cat	Italy	Europe	2004	[26]

^a Atlantic Forest biome; ^b Cerrado biome; ^c Amazon biome.

Table S2. Molecular diversity indexes of *Giardia duodenalis* based on β-giardin locus (592 bp, n=106).

Assemblage	Region (N)	Statistics					
		H ± SD	Nº of haplo-types	Nº of polymorphic sites	Nº of substitutions	Nº of transitions	Nº of transversions
Assemblage A	South America (31)	0.879 ± 0.037	14	38	39	21	18
	North America (4)	0.666 ± 0.204	2	4	4	4	0
	Europe (10)	0.822 ± 0.096	5	15	15	15	0
	Asia (2)	1.000 ± 0.500	2	4	4	4	0
	All continents (49)	0.854 ± 0.029	16	50	50	33	18
	All Amazon biome (5)	0.800 ± 0.164	3	8	8	6	2
	Amazon biome* (4)	0.833 ± 0.222	3	8	8	6	2
	All Cerrado biome (8)	0.964 ± 0.077	7	30	31	15	16
	Cerrado biome* (7)	1.000 ± 0.076	7	30	31	15	16
	Caatinga biome* (5)	0.700 ± 0.218	3	2	2	2	0
	All Atlantic Forest biome (13)	0.859 ± 0.063	6	6	6	6	0
	Atlantic Forest biome* (9)	0.750 ± 0.112	4	4	4	4	0
	Atlantic Forest biome (4)	0.833 ± 0.222	3	2	2	2	0
	All Brazil (31)	0.879 ± 0.037	14	38	39	21	18
Assemblage B	South America (41)	0.918 ± 0.033	23	36	36	31	6
	North America (2)	1.000 ± 0.500	2	3	3	3	0
	Europe (3)	1.000 ± 0.272	3	7	7	7	0
	Asia (5)	1.000 ± 0.126	5	5	5	5	0
	All continents (51)	0.921 ± 0.028	28	37	37	32	6
	All Amazon biome (37)	0.899 ± 0.039	19	24	24	23	1
	Amazon biome* (21)	0.757 ± 0.086	7	8	8	7	1
	Amazon biome (16)	0.991 ± 0.025	15	21	21	21	0
	All Atlantic Forest biome (3)	1.000 ± 0.272	3	10	10	7	3
	Atlantic Forest (2)	1.000 ± 0.500	2	5	5	4	1
ALL (106) **	All Brazil (41)	0.918 ± 0.033	23	36	36	31	6
		0.951 ± 0.009	50	125	147	105	42

H ± SD: gene diversity ± standard deviation. (bp): base pair. * Sequences obtained in this study **ALL: included assemblages A, B, C, D, E and F. Further details of reference strains can be found in Supplementary Table S1. Only groups with more than 1 sequence are shown.

References

- Lebbad M, Mattsson JG, Christensson B, Ljungström B, Backhans A, Andersson JO, et al. From mouse to moose: multilocus genotyping of *Giardia* isolates from various animal species. *Vet Parasitol.* 2010;168(3-4):231-9. doi: 10.1016/j.vetpar.2009.11.003. PMID: 19969422.
- Solarczyk P, Majewska AC, Moskwa B, Cabaj W, Dabert M, Nowosad P. Multilocus genotyping of *Giardia duodenalis* isolates from red deer (*Cervus elaphus*) and roe deer (*Capreolus capreolus*) from Poland. *Folia Parasitol (Praha).* 2012;59(3):237-40. doi: 10.14411/fp.2012.032 PMID: 23136805.
- Solarczyk P, Werner A, Majewska AC. Genotypowanie izolatów *Giardia duodenalis* uzyskanych od ludzi w zachodnio-centralnej Polsce [Genotype analysis of *Giardia duodenalis* isolates obtained from humans in west-central Poland]. *Wiad Parazytol.* 2010;56(2):171-7. PMID: 20707303.
- Lebbad M, Petersson I, Karlsson L, Botero-Kleiven S, Andersson JO, Svenungsson B, et al. Multilocus genotyping of human *Giardia* isolates suggests limited zoonotic transmission and association between assemblage B and flatulence in children. *PLoS Negl Trop Dis.* 2011;5(8):e1262. doi: 10.1371/journal.pntd.0001262 . PMID: 21829745.
- Durigan M, Abreu AG, Zucchi MI, Franco RMB, de Souza AP. Genetic Diversity of *Giardia duodenalis*: Multilocus Genotyping Reveals Zoonotic Potential between Clinical and Environmental Sources in a Metropolitan Region of Brazil. *PLoS ONE.* 2014 9(12): e115489. doi:10.1371/journal.pone.0115489.
- Prystajecky N, Tsui CK, Hsiao WW, Uyaguari-Diaz MI, Ho J, Tang P, et al. *Giardia* spp. Are Commonly Found in Mixed Assemblages in Surface Water, as Revealed by Molecular and Whole-Genome Characterization. *Appl Environ Microbiol.* 2015;81(14):4827-34. doi: 10.1128/AEM.00524-15 PMID: 25956776.
- Nunes BC, Calegar DA, Pavan MG, Jaeger LH, Monteiro KJL, Dos Reis ERC, et al. Genetic diversity of *Giardia duodenalis* circulating in three Brazilian biomes. *Infect Genet Evol.* 2018;59:107-112. doi: 10.1016/j.meegid.2018.02.001 PMID: 29410226.

8. Hassan, Z.I. Unpublished. Available in {HYPERLINK “<https://www.ncbi.nlm.nih.gov/nuccore/MN629930>”}.
9. Morrison HG, McArthur AG, Gillin FD, Aley SB, Adam RD, Olsen GJ, et al. Genomic minimalism in the early diverging intestinal parasite *Giardia lamblia*. *Science*. 2007;317(5846):1921–6. doi: 10.1126/science.114383 PMID: 17901334.
10. Teodorovic S, Braverman JM, Elmendorf HG. Unusually low levels of genetic variation among *Giardia lamblia* isolates. *Eukaryot Cell*. 2007;6(8):1421–30. doi: 10.1128/EC.00138-07 PMID: 17557879; PMCID: PMC1951139.
11. Wielinga C, Ryan U, Andrew Thompson RC, Monis P. Multi-locus analysis of *Giardia duodenalis* intra-Assemblage B substitution patterns in cloned culture isolates suggests sub-Assemblage B analyses will require multi-locus genotyping with conserved and variable genes. *Int J Parasitol*. 2011;41(5):495–503. doi: 10.1016/j.ijpara.2010.11.007 PMID: 21176781.
12. Cacciò SM, De Giacomo M, Pozio E. Sequence analysis of the beta-giardin gene and development of a polymerase chain reaction-restriction fragment length polymorphism assay to genotype *Giardia duodenalis* cysts from human faecal samples. *Int J Parasitol*. 2002;32(8):1023–30. doi: 10.1016/s0020-7519(02)00068-1 PMID: 12076631.
13. Robertson LJ, Hermansen L, Gjerde BK, Strand E, Alvsvåg JO, Langeland N. Application of genotyping during an extensive outbreak of waterborne giardiasis in Bergen, Norway, during autumn and winter 2004. *Appl Environ Microbiol*. 2006;72(3):2212–7. doi: 10.1128/AEM.72.3.2212-2217.2006 PMID: 16517674.
14. Di Giovanni GD, Betancourt WQ, Hernandez J, Assadian NW, Flores Margez JP, Lopez EJ. Investigation of potential zoonothroponotic transmission of cryptosporidiosis and giardiasis through agricultural use of reclaimed wastewater. *Int J Environ Health Res*. 2006;16(6):405–18. doi: 10.1080/09603120601095100 PMID: 17164167.
15. Teodorovic S, Braverman JM, Elmendorf HG. Unusually low levels of genetic variation among *Giardia lamblia* isolates. *Eukaryot Cell*. 2007;6(8):1421–30. doi: 10.1128/EC.00138-07 PMID: 17557879.
16. Bonhomme J, Le Goff L, Lemée V, Gargala G, Ballet JJ, Favennec L. Limitations of tpi and bg genes sub-genotyping for characterization of human *Giardia duodenalis* isolates. *Parasitol Int*. 2011;60(3):327–30. doi: 10.1016/j.parint.2011.05.004 PMID: 21627998.
17. Kosuwin R, Putaporntip C, Pattanawong U, Jongwutiwes S. Clonal diversity in *Giardia duodenalis* isolates from Thailand: evidences for intragenic recombination and purifying selection at the beta giardin locus. *Gene*. 2010;449(1-2):1–8. doi: 10.1016/j.gene.2009.09.010 PMID: 19796671.
18. Alonso JL, Amorós I, Cuesta G. LNA probes in a real-time TaqMan PCR assay for genotyping of *Giardia duodenalis* in wastewaters. *J Appl Microbiol*. 2010;108(5):1594–601. doi: 10.1111/j.1365-2672.2009.04559.x PMID: 19840182.
19. Coronato Nunes B, Pavan MG, Jaeger LH, Monteiro KJ, Xavier SC, Monteiro FA, et al. Spatial and Molecular Epidemiology of *Giardia intestinalis* Deep in the Amazon, Brazil. *PLoS One*. 2016;11(7):e0158805. doi: 10.1371/journal.pone.0158805 PMID: 27392098.
20. Hussein AI, Tokoro M. Unpublished. Available in {HYPERLINK “<https://www.ncbi.nlm.nih.gov/nuccore/AB480877>”}.
21. Abe N, Teramoto I. Molecular evidence for person-to-person transmission of a novel subtype in *Giardia duodenalis* assemblage B at the rehabilitation institution for developmentally disabled people. *Parasitol Res*. 2012;110(2):1025–8. doi: 10.1007/s00436-011-2564-4 PMID: 21786066.
22. Tokoro M, Mizuno T, Hendarto J, Matey EJ, Songok EM, Ichimura H. Unpublished. Available in {HYPERLINK “<https://www.ncbi.nlm.nih.gov/nuccore/LC508615>”}.
23. Köster PC, Malheiros AF, Shaw JJ, Balasegaram S, Prendergast A, Lucaccioni H, et al. Multilocus Genotyping of *Giardia duodenalis* in Mostly Asymptomatic Indigenous People from the Tapirapé Tribe, Brazilian Amazon. *Pathogens*. 2021;10(2):206. <https://doi.org/10.3390/pathogens10020206>.
24. Paz e Silva FM, Monobe MM, Lopes RS, Araujo JP Jr. Molecular characterization of *Giardia duodenalis* in dogs from Brazil. *Parasitol Res*. 2012;110(1):325–34. doi: 10.1007/s00436-011-2492-3 PMID: 21695567.
25. Cacciò SM, De Giacomo M, Pozio E. Sequence analysis of the beta-giardin gene and development of a polymerase chain reaction-restriction fragment length polymorphism assay to genotype *Giardia duodenalis* cysts from human faecal samples. *Int J Parasitol*. 2002;32(8):1023–30. doi: 10.1016/s0020-7519(02)00068-1 PMID: 12076631.
26. Lalle M, Pozio E, Capelli G, Bruschi F, Crotti D, Cacciò SM. Genetic heterogeneity at the beta-giardin locus among human and animal isolates of *Giardia duodenalis* and identification of potentially zoonotic subgenotypes. *Int J Parasitol*. 2005;35(2):207–13. doi: 10.1016/j.ijpara.2004.10.022 PMID: 15710441.