

Supplementary Material

Modeling gastrointestinal tract wet pool size in small ruminants

Paola R. Ribeiro¹, Marcelo Gindri², Gilberto L. Macedo Junior³, Caio J. L. Herbster⁴, Elzania S. Pereira⁴, Bruno Biagioli¹, and Isabelle A. M. A. Teixeira^{1,5,*}

¹ Department of Animal Science, São Paulo State University, Jaboticabal 14884900, SP, Brazil; paola.rezenderibeiro@gmail.com (P.R.R.); bruno.biagioli@unesp.br (B.B.)

² UMR Modélisation Systémique Appliquée aux Ruminants, AgroParisTech, INRAE, Université Paris-Saclay, 91120, Palaiseau, France; gindri.marcelo@gmail.com

³ Department of Animal Science, Federal University of Uberlândia, Uberlândia 38402018, MG, Brazil; gilberto.macedo@ufu.br

⁴ Department of Animal Science, Federal University of Ceará, Fortaleza 60356000, CE, Brazil; julioherbster@gmail.com (C.J.L.H.); elzania@hotmail.com (E.S.P.)

⁵ Department of Animal Veterinary, and Food Sciences, University of Idaho, Twin Falls, ID 83301, USA

* Correspondence: izabelle@uidaho.edu




Figure S1: Studentized residuals of the linear equations proposed to describe the gastrointestinal tract wet pool size (GITwps) of small ruminants. Goats are grey and sheep are black.




Figure S2: Studentized residuals of the non-linear equations proposed to describe gastrointestinal tract wet pool size (GITwps) of small ruminants. Goats are grey and sheep are black.