

Table 1. SM. Change in the score of the NMSS and its domains between the visits of the study: V1 (N=50), V2 (N=47), V3 (N=45), V4 (N=44). .

	V1	V2	V3	V4	p ^a	p ^b	p ^c	p ^d
	97.48 ±	65.98 ±	59.40 ±	59.91 ±				
	43.70	40.10	32.34	35.49				
NMSS total score	9.58 ±	6.29 ±	5.09 ±	6.72 ±				
- Cardiovascular	2.46	10.64	8.23	11.94				
- Sleep / fatigue	36.08 ±	23.58 ±	18.33 ±	23.15 ±				
- Mood / apathy	21.77	20.97	14.36	18.12	<0.000	<0.000	<0.000	
- Perceptual symptoms	34.42 ±	19.47 ±	16.94 ±	14.49 ±	1	1	1	0.525
- Attention / memory	29.89	21.51	17.29	19.63	0.268	0.056	<0.000	0.500
- Gastrointestinal symptoms	4.33 ±	3.02 ±7.04	2.66 ± 4-	2.84 ±	0.002	<0.000	1	0.912
- Urinary symptoms	8.67	14.83 ±	32	5.88	<0.000	1	<0.000	0.604
- Sexual dysfunction	17.50 ±	18.86	14.26 ±	13.32 ±	1	<0.000	1	0.640
- Miscellaneous	17.09	13.94 ±	16.93	18.19	0.630	1	0.515	0.959
Dose of safinamide (mg/day)	19.61 ±	16.94	14.07 ±	13.13 ±	0.026	0.896	0.150	0.764
	18.01	30.32 ±	14.59	13.39	0.010	0.070	0.013	0.701
	42.72 ±	24.96	28.21 ±	30.62 ±	0.003	0.013	0.001	0.474
	30.41	24.29 ±	27.99	23.94	0.784	0.002	0.249	0.235
	28.25 ±	36.91	27.87 ±	25.28 ±	<0.000	0.867	0.001	
	35.69	22.78 ±	35.62	33.58	1	0.084		
	33.33 ±	16.29	19.12 ±	18.99 ±				
	20.73		15.01	14.03				
	N. A.	53.84 ±	96.15 ±	98.72 ±				
		13.49	13.49	8.00				

P values were computed using the Wilcoxon signed-rank test. The results represent mean ± SD. Domains of the NMSS were expressed as a percentage to be able to establish comparisons on their severity between them; p^a, V4 vs V1; p^b, V3 vs V1; p^c, V2 vs V1; p^d, V4 vs V2. N. A., Not applicable.

NMSS, Non-Motor Symptoms Scale.