

Supplementary Materials

Branched Sulfonyl Imide-Based Proton Exchange Polymer Membranes from poly(phenylenebenzopheneone)s for Fuel Cells Application

Sabuj Chandra Sutradhar ^{1,†}, Sujin Yoon ^{1,†}, Taewook Ryu ¹, Lei Jin ¹, Wei Zhang ¹, Whangi Kim ¹ and Hohyoun Jang ^{2*}

¹ Department of Applied Chemistry, Konkuk University, Chungju 27478, Korea; sabujchandra@gmail.com (S.C.S.); ysj920126@naver.com (S.Y.); gundam0924@naver.com (T.R.); jinlei8761@naver.com (L.J.); arno_zw@hotmail.com (W.Z.); wgkim@kku.ac.kr (W.K.)

² Department of Liberal Art, Konkuk University, Chungju 27478, Korea

* Correspondence: 200417450@kku.ac.kr; +82-43-840-4764 (ext. 410)

† These authors contributed equally to this work.


Scheme

Scheme S1. Synthesis route for PBP monomer.

Figures

Figure S1. ¹H NMR of PBP monomer.

Scheme



Scheme S1. Synthesis route for PBP monomer.

Figures




Figure S1. ¹H NMR of PBP monomer.