

Supplementary material

Article

Genotype-Related Clinical Characteristics and Myocardial Fibrosis and their Association with Prognosis in Hypertrophic Cardiomyopathy

Hyung Yoon Kim ^{1,†}, Jong Eun Park ^{2,†}, Sang-Chol Lee ^{3,*}, Eun-Seok Jeon ³, Young Keun On ³, Sung Mok Kim ⁴, Yeon Hyeon Choe ⁴, Chang-Seok Ki ⁵, Jong-Won Kim ⁶ and Kye Hun Kim ¹

¹ Department of Cardiovascular Medicine, Chonnam National University Medical School/Hospital, Gwangju, Korea; medoc7@gmail.com (H.Y.K.); christiankyehun@hanmail.net (K.H.K.)

² Department of Laboratory Medicine, Hanyang University Guri Hospital, Hanyang University College of Medicine, Guri, Korea; jongeun820@gmail.com

³ Department of Internal medicine, Cardiovascular Imaging Center, Heart, Vascular & Stroke Institute; eunseok.jeon@samsung.com (E.-S.J.); yk.on@samsung.com (Y.K.O.)

⁴ Department of Radiology, Cardiovascular Imaging Center, Heart, Vascular & Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea; sungmok_kim@hanmail.net (S.M.K.); ychoe11@gmail.com (Y.H.C.)

⁵ Green Cross Genome, Yongin, Korea; changski.md@gmail.com

⁶ Department of Laboratory Medicine and Genetics, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea; kimjw@skku.edu

* Correspondence: chrislee.echo@gmail.com; Tel.: +82-2-3410-3419

† These authors equally contributed to this work as co-first authors.

Supplementary Table S1. Medication prescribed during follow-up

Variables	Detected (n=55)	Not detected (n=34)	P-value
	Mean ± SD or number (%)		
Amiodarone	0	0	NS
Beta-blocker	44 (80.0)	27 (79.4)	0.946
Calcium channel blocker	8 (14.5)	6 (17.6)	0.768
ACEi	1 (1.8)	0	0.618
ARB	8 (14.5)	8 (23.5)	0.395
Statin	3 (5.5)	4 (11.8)	0.421
Warfarin	2 (3.6)	1 (2.9)	0.676
Aspirin	21 (38.2)	17 (50.0)	0.378