

Supplementary Materials

Band Structure and Energy Level Alignment of Chiral Graphene Nanoribbons on Silver Surfaces

Martina Corso ^{1,2,*}, Rodrigo E. Menchón ^{2,3}, Ignacio Piquero-Zulaica ^{1,†}, Manuel Vilas-Varela ⁴, J. Enrique Ortega ^{1,2,5}, Diego Peña ³, Aran Garcia-Lekue ^{2,6,*} and Dimas G. de Oteyza ^{1,2,6,*}

¹ Centro de Física de Materiales (MPC), CSIC-UPV/EHU, 20018 San Sebastián, Spain; ge46biq@mytum.de (I.P.-Z.); enrique.ortega@ehu.es (J.E.O.)

² Donostia International Physics Center (DIPC), 20018 San Sebastián, Spain; menchon@dipc.org

³ Dpto. Departamento de Polímeros y Materiales Avanzados: Física, Química y Tecnología, Universidad del País Vasco UPV/EHU, 20080 San Sebastián, Spain; diego.pena@usc.es

⁴ Centro Singular de Investigación en Química Biolóxica e Materiais Moleculares (CiQUS) and Departamento de Química Orgánica, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain; manuel.vilas.varela@usc.es

⁵ Dpto. Física Aplicada I, Universidad del País Vasco, 20018 San Sebastián, Spain

⁶ Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain

* Correspondence: martina.corso@ehu.eus (M.C.); wmbgalea@ehu.eus (A.G.-L.); d_g_oteyza@ehu.es (D.G.d.O.)

† Current address: Physics Department E20, Technical University of Munich, 85748 Garching, Germany.




Figure S1. ARPES raw data and their second derivative for three representative regions of the GNR-covered curved Ag crystal surface (characterized by periodic {100} steps, flat (111) surface and periodic {111} steps), along with the reference map on the clean substrate. The displayed measurements are obtained at vicinal angles of approximately -10° , 0 and 10° .




Figure S2. ARPES data obtained at five regularly spaced positions across the curved silver surface at vicinal angles of approximately -10° , -5° , 0 , 5° and 10° .