



Article

Antioxidative Activity of Soy, Wheat and Pea Protein Isolates Characterized by Multi-Enzyme Hydrolysis

Chiung-Yueh Chang ¹, Jinn-Der Jin ², Hsiao-Li Chang ², Ko-Chieh Huang ¹, Yi-Fen Chiang ¹, Mohamed Ali ³ and Shih-Min Hsia ^{1,4,5,6,*}

¹ School of Nutrition and Health Sciences, College of Nutrition, Taipei Medical University, Taipei 110, Taiwan; d507104001@tmu.edu.tw (C.-Y.C.); a910241@gmail.com (K.-C.H.); yvonne840828@gmail.com (Y.-F.C.); bryanhsia@tmu.edu.tw (S.-M.H)

² GeneFerm Biotechnology Co., Ltd., Tainan 741, Taiwan; jin168@geneferm.com (J.-D.J); gracechang@geneferm.com (H.-L.C)

³ Clinical Pharmacy Department, Faculty of Pharmacy, Ain Shams University, Cairo 11566, Egypt; mohamed.aboouf@pharma.asu.edu.eg

⁴ Graduate Institute of Metabolism and Obesity Sciences, College of Nutrition, Taipei Medical University, Taipei 110, Taiwan

⁵ School of Food and Safety, Taipei Medical University, Taipei 110, Taiwan

⁶ Nutrition Research Center, Taipei Medical University Hospital, Taipei 110, Taiwan

* Correspondence: bryanhsia@tmu.edu.tw; Tel.: +886-2-2736-1661




Figure S1. Toxicity effect of PIH on C2C12. C2C12 myoblasts (1×10^4 cells) were cultured in 96-well plates using 2% horse serum DMEM to differentiate the myotubes. After 6 days of differentiation, the samples were treated with different dosages of plant PIH for 24 hours * $p < 0.05$; **, $p < 0.01$ versus 0 mg/mL cells. PIH: protein isolate hydrolysate.