

Supplementary Materials

Efficacy of Nanofiber Sheets Incorporating Lenvatinib in a Hepatocellular Carcinoma Xenograft Model

Terufumi Yoshida ¹, Masaki Kaibori ^{1,*}, Nanami Fujisawa ², Mariko Ishizuka ¹, Fusao Sumiyama ¹, Masahiko Hatta ¹, Hisashi Kosaka ¹, Kosuke Matsui ¹, Kensuke Suzuki ³, Tomoya O. Akama ⁴, Tayo Katano ⁵, Kengo Yoshii ⁶, Mitsuhiro Ebara ² and Mitsugu Sekimoto ¹

- ¹ Department of Surgery, Kansai Medical University, 2-5-1 Shinmachi, Hirakata 573-1010, Japan; yoshiter@hirakata.kmu.ac.jp (T.Y.); ishizukm@takii.kmu.ac.jp (M.I.); sumiyamf@hirakata.kmu.ac.jp (F.S.); hattamas@hirakata.kmu.ac.jp (M.H.); kosakahi@hirakata.kmu.ac.jp (H.K.); matsuk@hirakata.kmu.ac.jp (K.M.); sekimotm@hirakata.kmu.ac.jp (M.S.)
- ² Research Center for Functional Materials, National Institute for Materials Science (NIMS), Tsukuba 305-0044, Japan; nfujisawa.tokyo@gmail.com (N.F.); ebara.mitsuhiro@nims.go.jp (M.E.)
- ³ Department of Otolaryngology, Head and Neck Surgery, Kansai Medical University, Hirakata 573-1010, Japan; suzukken@hirakata.kmu.ac.jp
- ⁴ Department of Pharmacology, Kansai Medical University, Hirakata 573-1010, Japan; akamat@hirakata.kmu.ac.jp
- ⁵ Department of Medical Chemistry, Kansai Medical University, Hirakata 573-1010, Japan; katanot@hirakata.kmu.ac.jp
- ⁶ Department of Mathematics and Statistics in Medical Sciences, Kyoto Prefectural University of Medicine, Kyoto 606-0823, Japan; yoshii-k@koto.kpu-m.ac.jp
- * Correspondence: kaibori@hirakata.kmu.ac.jp; Tel.: +81-72-804-0101 (ext. 56130)




Figure S1: DSC curves of lenvatinib incorporated nanofibers, PCL nanofibers and lenvatinib.