



*Supplementary information*

# Voltammetric Behaviour of Drug Molecules as a Predictor of Metabolic Liabilities

Hikari Fuchigami <sup>1</sup>, Mandeep K. Bal <sup>2</sup>, Dale A. C. Brownson <sup>2</sup>, Craig E. Banks <sup>2</sup> and Alan M. Jones <sup>3,\*</sup>

<sup>1</sup> School of Biosciences, University of Birmingham, Edgbaston, B15 2TT, United Kingdom;  
[HXF916@student.bham.ac.uk](mailto:HXF916@student.bham.ac.uk)

<sup>2</sup> Manchester Metropolitan University, Faculty of Science and Engineering, Chester Street, Manchester, M1 5GD, United Kingdom; [m.kaur@mmu.ac.uk](mailto:m.kaur@mmu.ac.uk) (M.K.B.); [D.Brownson@mmu.ac.uk](mailto:D.Brownson@mmu.ac.uk) (D.A.C.B.);  
[C.Banks@mmu.ac.uk](mailto:C.Banks@mmu.ac.uk) (C.E.B.)

<sup>3</sup> School of Pharmacy, University of Birmingham, Edgbaston, B15 2TT, United Kingdom

\* Correspondence: [a.m.jones.2@bham.ac.uk](mailto:a.m.jones.2@bham.ac.uk); Tel.: +44-01214-147-288

Received: 15 September 2020; Accepted: 3 October 2020; Published: date

## CONTENTS

<sup>1</sup> H and <sup>13</sup> C NMR spectra of synthesised and purchased drug molecules	page 2-7
--	----------


Figure S1.  $^1\text{H}$  NMR spectrum of **2** (400 MHz,  $\text{CDCl}_3$ ).

Figure S2.  $^{13}\text{C}$  NMR spectrum of **2** (101 MHz,  $\text{CDCl}_3$ ).





Figure S3.  $^1\text{H}$  NMR spectrum of Propanidid 400 MHz,  $\text{CDCl}_3$ ).

Figure S4.  $^{13}\text{C}$  NMR spectrum of Propanidid (101 MHz,  $\text{CDCl}_3$ ).






Figure S5.  $^1\text{H}$  NMR spectrum of Entacapone (400 MHz,  $\text{CDCl}_3$ ).

Figure S6.  $^1\text{H}$  NMR spectrum of Lidocaine (400 MHz,  $\text{CDCl}_3$ ).



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).