
Supplementary file

Bioprospecting of Essential Oil-Bearing Plants: Rapid Screening of Volatile Organic Compounds Using Headspace Bubble-in-Drop Single-Drop Microextraction for Gas Chromatography Analysis

Thabiso E. Letseka¹, Ntjana J. Sepheka¹, Ian A. Dubery^{2,*} and Mosotho J. George^{1,2,*}

¹ Department of Chemistry and Chemical Technology, National University of Lesotho, P.O. Box 180, Roma 100, Lesotho

² Department of Biochemistry, University of Johannesburg, P.O. Box 524, Auckland Park 2006, South Africa

* Correspondence: idubery@uj.ac.za (I.A.D.); jm.george@nul.ls (M.J.G.)

Table S1. Structures of terpenes and terpenoids detected by GC-MS in the VOC fractions as samples by the BID-SDME method. The numbered chemicals correspond to that used in Table 1 in the main text. Structures were obtained from Pubchem (<https://pubchem.ncbi.nlm.nih.gov>, accessed on 10 October 2022).

1. , 6. 	2. 	3. 	4. 	5.
7. 	8. 	9. , 10. 	11. 	12. , 13.
14. 	15. 	16. 	17. 	18.
19. 	20. 	21. 	22. 	