

Joint Universality of the Zeta-Functions of Cusp Forms

Renata Macaitienė 

Regional Development Institute of Šiauliai Academy, Vilnius University, P. Višinskio Str. 25,
LT-76351 Šiauliai, Lithuania; renata.macaitiene@sa.vu.lt

Abstract: Let F be the normalized Hecke-eigen cusp form for the full modular group and $\zeta(s, F)$ be the corresponding zeta-function. In the paper, the joint universality theorem on the approximation of a collection of analytic functions by shifts $(\zeta(s + ih_1\tau, F), \dots, \zeta(s + ih_r\tau, F))$ is proved. Here, h_1, \dots, h_r are algebraic numbers linearly independent over the field of rational numbers.

Keywords: Hecke-eigen cusp form; joint universality; universality; zeta-function

MSC: 11M46

1. Introduction

The series of the types

$$\sum_{m=1}^{\infty} \frac{a_m}{m^s} \quad \text{and} \quad \sum_{m=1}^{\infty} a_m e^{-\lambda_m s}, \quad s = \sigma + it,$$

where $\{\lambda_m\}$ is a nondecreasing sequence of real numbers and $\lim_{m \rightarrow \infty} \lambda_m = +\infty$ are called Dirichlet series. The majority of zeta-functions are meromorphic functions in some half-plane defined by Dirichlet series having a certain arithmetic sense. The most important of zeta-functions is the Riemann zeta-function

$$\zeta(s) = \sum_{m=1}^{\infty} \frac{1}{m^s}, \quad \sigma > 1.$$

In [1], Voronin discovered a very interesting and important property of $\zeta(s)$ to approximate a wide class of analytic functions by shifts $\zeta(s + i\tau)$, $\tau \in \mathbb{R}$, and called it universality. Later, it turned out that some other zeta-functions also are universal in the Voronin sense. This paper is devoted to the universality of zeta-functions of certain cusp forms.

Let

$$SL(2, \mathbb{Z}) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \mathbb{Z}, ad - bc = 1 \right\}$$

be the full modular group. If the function $F(z)$ is holomorphic in the upper half-plane $\text{Im} z > 0$, and for all elements of $SL(2, \mathbb{Z})$ with some $\kappa \in 2\mathbb{N}$ satisfies the functional equation

$$F\left(\frac{az + b}{cz + d}\right) = (cz + d)^{\kappa} F(z), \quad (1)$$

where $F(z)$ is called a modular form of weight κ for the full modular group. Then, $F(z)$ has Fourier series expansion

$$F(z) = \sum_{m=-\infty}^{\infty} c(m) e^{2\pi i m z}.$$



Citation: Macaitienė, R. Joint Universality of the Zeta-Functions of Cusp Forms. *Mathematics* **2021**, *9*, 2161. <https://doi.org/10.3390/math9172161>

Academic Editor: Yang-Hui He

Received: 26 July 2021

Accepted: 31 August 2021

Published: 4 September 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

If $c(m) = 0$ for all $m \leq 0$, then $F(z)$ is a cusp form of weight κ . The corresponding zeta-function (or L -function) $\zeta(s, F)$ is defined for $\sigma > \frac{\kappa+1}{2}$ by the Dirichlet series

$$\zeta(s, F) = \sum_{m=1}^{\infty} \frac{c(m)}{m^s},$$

and has the analytic continuation to an entire function. Additionally, we suppose that $F(z)$ is a simultaneous eigenfunction of all Hecke operators T_m

$$T_m F(z) = m^{\kappa-1} \sum_{\substack{a, d > 0 \\ ad=m}} \frac{1}{d^{\kappa}} \sum_{b \pmod{d}} F\left(\frac{az+b}{d}\right), \quad m \in \mathbb{N}.$$

In this case, $c(1) \neq 0$; therefore, the form $F(z)$ can be normalized, and thus, we may suppose that $c(1) = 1$.

Now, we suppose that $F(z)$ is a normalized Hecke-eigen cusp form of weight κ for the full modular group. Then, the zeta-function $\zeta(s, F)$ can be written, for $\sigma > \frac{\kappa+1}{2}$, as a product over primes

$$\zeta(s, F) = \prod_p \left(1 - \frac{\alpha(p)}{p^s}\right)^{-1} \left(1 - \frac{\beta(p)}{p^s}\right)^{-1},$$

where $\alpha(p)$ and $\beta(p)$ are conjugate complex numbers satisfying the equality $\alpha(p) + \beta(p) = c(p)$.

In the paper [2], the universality of the function $\zeta(s, F)$ was proved. Let $D_{\kappa} = \{s \in \mathbb{C} : \frac{\kappa}{2} < \sigma < \frac{\kappa+1}{2}\}$, \mathcal{K}_F be the class of compact subsets of the strip D_{κ} with connected complements, and $H_{0,F}(K)$, $K \in \mathcal{K}_F$ the class of continuous nonvanishing functions on K that are analytic in the interior of K . Moreover, let $\text{meas} A$ denote the Lebesgue measure of a measurable set $A \subset \mathbb{R}$. Then, in [2], the following theorem was obtained.

Theorem 1. Suppose that $K \in \mathcal{K}_F$ and $f(s) \in H_{0,F}(K)$. Then, for every $\varepsilon > 0$,

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{s \in K} |\zeta(s + i\tau, F) - f(s)| < \varepsilon \right\} > 0.$$

Theorem 1 shows that there are infinitely many shifts $\zeta(s + i\tau, F)$ approximating a given function $f(s) \in H_{0,F}$. In the shifts $\zeta(s + i\tau, F)$ of Theorem 1, τ takes arbitrary real values; therefore, the theorem is of continuous type. Further, discrete universality theorems for the function $\zeta(s, F)$ are known. In [3,4], the discrete universality theorems with shifts $\zeta(s + ikh, F)$, $k \in \mathbb{N}$, $h > 0$ being a fixed number, were proved. Denote by $H(D_{\kappa})$ the space of analytic on D_{κ} functions endowed with the topology of uniform convergence on compacta. The paper [5] is devoted to the universality for compositions $\Phi(\zeta(s, F))$ with certain operators $\Phi : H(D_{\kappa}) \rightarrow H(D_{\kappa})$. The results of the latter paper were applied in [6] for the functional independence of the compositions $\Phi(\zeta(s, F))$.

Let, for a fixed $l \in \mathbb{N}$,

$$\Gamma_0(l) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) : c \equiv 0 \pmod{l} \right\}$$

denote the Hecke subgroup of the group $SL(2, \mathbb{Z})$. If $F(z)$ satisfies (1) for all elements of $\Gamma_0(l)$, then $F(z)$ is called a cusp form of weight κ and level l . The form $F(z)$ is called a new form if it is not a cusp form of level $l_1 \mid l$. In [7], a universality theorem was obtained for zeta-functions of new forms.

The universality theorem of [2] was generalized in [8] for shifts $\zeta(s + i\varphi(\tau), F)$ with differentiable function $\varphi(\tau)$ satisfying the estimates $(\varphi'(\tau))^{-1} = o(\tau)$ and $\varphi(2\tau) \max_{\tau \leq t \leq 2\tau} (\varphi'(t))^{-1}$

$\ll \tau$ as $\tau \rightarrow \infty$. The discrete version of results of [8] is given in [9]. In [10], the shifts $\zeta(s + i\gamma_k, F)$, where $\{\gamma_k : k \in \mathbb{N}\}$ is the sequence of nontrivial zeros of $\zeta(s)$, are used.

The joint universality of zeta- and L -functions is a more complicated problem of analytic number theory. In this case, a collection of analytic functions are simultaneously approximated by a collection of shifts of zeta-functions. The first result in this direction also belongs to Voronin. He considered [11] the functional independence of Dirichlet L -functions $L(s, \chi)$ with pairwise nonequivalent Dirichlet characters χ and, for this, he obtained their joint universality. The paper [12] is devoted to the joint universality for zeta-functions of new forms twisted by Dirichlet characters, i.e., for the functions

$$\sum_{m=1}^{\infty} \frac{c(m)\chi(m)}{m^s}, \quad \sigma > \frac{\kappa+1}{2},$$

with pairwise nonequivalent Dirichlet characters χ_1, \dots, χ_r .

Joint universality theorems with generalized shifts $\zeta(s + i\varphi_j(k), F)$, $j = 1, \dots, r$, with some differentiable functions $\varphi_j(\tau)$ can be found in [13]. Continuous and discrete joint universality theorems for more general zeta-functions are given in [14–16].

Our aim is to obtain a joint universality theorem for zeta-functions of normalized Hecke-eigen cusp forms by using different shifts. The first of the denseness results for shifts of a universal function were discussed in [17].

The main result of the paper is the following statement.

Theorem 2. Suppose that h_1, \dots, h_r are real algebraic numbers linearly independent over the field of rational numbers \mathbb{Q} . For $j = 1, \dots, r$, let $K_j \in \mathcal{K}_F$ and $f_j(s) \in H_{0,F}(K_j)$. Then, for every $\varepsilon > 0$,

$$\liminf_{T \rightarrow \infty} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \sup_{1 \leq j \leq r} \sup_{s \in K_j} |\zeta(s + ih_j\tau, F) - f_j(s)| < \varepsilon \right\} > 0.$$

Moreover “lim inf” can be replaced by “lim” for all but at most countably many $\varepsilon > 0$.

For the proof of Theorem 2, we will apply the probabilistic approach based on a limit theorem in the space of analytic functions.

2. Mean Square Estimates

Recall the metric in the space $H(D_K)$. Let $\{K_l : l \in \mathbb{N}\} \subset D_K$ be a sequence of compact subsets such that

$$D_K = \bigcup_{l=1}^{\infty} K_l,$$

$K_l \subset K_{l+1}$ for $l \in \mathbb{N}$, and if $K \subset D_K$ is a compact, then $K \subset K_l$ for some l . For example, we can take K_l closed rectangles. Then

$$\rho(g_1, g_2) = \sum_{l=1}^{\infty} 2^{-l} \frac{\sup_{s \in K_l} |g_1(s) - g_2(s)|}{1 + \sup_{s \in K_l} |g_1(s) - g_2(s)|}, \quad g_1, g_2 \in H(D_K),$$

is a metric in $H(D_K)$ inducing the topology of uniform convergence on compacta.

Let

$$H^r(D_K) = \underbrace{(H(D_K) \times \dots \times H(D_K))}_r.$$

For $\underline{g}_j = (g_{j1}, \dots, g_{jr}) \in H^r(D_K)$, $j = 1, 2$, define

$$\rho(\underline{g}_1, \underline{g}_2) = \max_{1 \leq j \leq r} \rho(g_{1j}, g_{2j}).$$

Then, ρ is a metric in $H^r(D_\kappa)$ inducing the product topology.

Let $\theta > \frac{1}{2}$ be a fixed number, and

$$v_n(m) = \exp\left\{-\left(\frac{m}{n}\right)^\theta\right\}, \quad m, n \in \mathbb{N}.$$

Then, the series

$$\zeta_n(s, F) = \sum_{m=1}^{\infty} \frac{c(m)v_n(m)}{m^s},$$

in view of the estimate

$$c(m) \ll m^{\frac{\kappa-1}{2}+\varepsilon},$$

is absolutely convergent in every fixed half plane $\sigma > \hat{\sigma}$. However, for our aim, this convergence is sufficient only for $\sigma > \frac{\kappa}{2}$.

For brevity, let $\underline{h} = (h_1, \dots, h_r)$,

$$\underline{\zeta}(s + i\underline{h}\tau, F) = (\zeta(s + ih_1\tau, F), \dots, \zeta(s + ih_r\tau, F))$$

and

$$\underline{\zeta}_n(s + i\underline{h}\tau, F) = (\zeta_n(s + ih_1\tau, F), \dots, \zeta_n(s + ih_r\tau, F)).$$

Lemma 1. For all \underline{h} ,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho(\underline{\zeta}(s + i\underline{h}\tau, F), \underline{\zeta}_n(s + i\underline{h}\tau, F)) d\tau = 0.$$

Proof. By the definitions of the metrics ρ and $\underline{\rho}$, it suffices to show that, for every $h \in \mathbb{R}$ and compact set $K \subset D_\kappa$,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \sup_{s \in K} |\zeta(s + ih\tau, F) - \zeta_n(s + ih\tau, F)| d\tau = 0.$$

It is well known that for fixed $\frac{\kappa}{2} < \sigma < \frac{\kappa+1}{2}$,

$$\int_{-T}^T |\zeta(\sigma + it, F)|^2 dt \ll_\sigma T,$$

where \ll_σ means that the implied constant depends on σ . Therefore,

$$\int_{-T}^T |\zeta(\sigma + iht, F)|^2 dt \ll_{\sigma, h} T,$$

and, for $v \in \mathbb{R}$,

$$\frac{1}{T} \int_0^T |\zeta(\sigma + ih\tau + iv, F)|^2 dv \ll_{\sigma, h} 1 + |v|. \quad (2)$$

Let

$$l_n(s) = \frac{z}{\theta} \Gamma\left(\frac{z}{\theta}\right) n^z,$$

where $\Gamma(z)$ denotes the Euler gamma-function and θ is a number from the definition of $v_n(m)$. Using the Mellin formula

$$\frac{1}{2\pi i} \int_{\beta-i\infty}^{\beta+i\infty} \Gamma(s) \alpha^s ds = e^{-\alpha}, \quad \alpha, \beta > 0,$$

we find that

$$\exp\left\{-\left(\frac{m}{n}\right)^\theta\right\} = \frac{1}{2\pi i} \int_{\theta-i\infty}^{\theta+i\infty} \frac{1}{\theta} \Gamma\left(\frac{1}{\theta}\right) \left(\frac{m}{n}\right)^{-s} ds.$$

Therefore, in virtue of the definition of the function $v_n(m)$, we obtain that, for $\sigma > \frac{\kappa}{2}$,

$$\begin{aligned} \zeta_n(s, F) &= \frac{1}{2\pi i} \sum_{m=1}^{\infty} \frac{c(m)}{m^s} \int_{\theta-i\infty}^{\theta+i\infty} \frac{z}{\theta} \Gamma\left(\frac{1}{\theta}\right) \left(\frac{m}{n}\right)^{-z} \frac{dz}{z} \\ &= \frac{1}{2\pi i} \int_{\theta-i\infty}^{\theta+i\infty} \left(\frac{l_n(z)}{z} \sum_{m=1}^{\infty} \frac{c(m)}{m^{s+z}}\right) dz \\ &= \frac{1}{2\pi i} \int_{\theta-i\infty}^{\theta+i\infty} \zeta(s+z, F) l_n(z) \frac{dz}{z}. \end{aligned} \quad (3)$$

Let $K \in D_\kappa$ be a fixed compact set. Then, there exists $\varepsilon > 0$ such that, for all $s = \sigma + it \in K$, the inequalities $\frac{\kappa}{2} + 2\varepsilon < \sigma < \frac{\kappa+1}{2} - \varepsilon$ are satisfied. We take, for such σ ,

$$\theta_1 = \frac{\kappa}{2} + \varepsilon - \sigma.$$

Then, $\theta_1 < 0$. Therefore, by the residue theorem and (3),

$$\zeta_n(s, F) - \zeta(s, F) = \frac{1}{2\pi i} \int_{\theta_1-i\infty}^{\theta_1+i\infty} \zeta(s+z, F) l_n(z) \frac{dz}{z}.$$

Hence, for all $s \in K$,

$$\begin{aligned} \zeta(s + ih\tau, F) - \zeta_n(s + ih\tau, F) &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \zeta\left(\frac{\kappa}{2} + \varepsilon + it + ih\tau + iv, F\right) \frac{l_n\left(\frac{\kappa}{2} + \varepsilon - \sigma + iv\right)}{\frac{\kappa}{2} + \varepsilon - \sigma + iv} dv \\ &= \frac{1}{2\pi i} \int_{-\infty}^{\infty} \zeta\left(\frac{\kappa}{2} + \varepsilon + ih\tau + iv, F\right) \frac{l_n\left(\frac{\kappa}{2} + \varepsilon - s + iv\right)}{\frac{\kappa}{2} + \varepsilon - s + iv} dv \\ &\ll \frac{1}{2\pi i} \int_{-\infty}^{\infty} \left| \zeta\left(\frac{\kappa}{2} + \varepsilon + ih\tau + iv, F\right) \right| \sup_{s \in K} \left| \frac{l_n\left(\frac{\kappa}{2} + \varepsilon - s + iv\right)}{\frac{\kappa}{2} + \varepsilon - s + iv} \right| dv. \end{aligned}$$

Thus, in view of (2),

$$\begin{aligned} &\frac{1}{T} \int_0^\infty \sup_{s \in K} |\zeta(s + ih\tau, F) - \zeta_n(s + ih\tau, F)| d\tau \\ &\ll \int_{-\infty}^{\infty} \left(\left(\frac{1}{T} \int_0^\infty \left| \zeta\left(\frac{\kappa}{2} + \varepsilon + ih\tau + iv\right) \right|^2 d\tau \right)^{1/2} \sup_{s \in K} \left| \frac{l_n\left(\frac{\kappa}{2} + \varepsilon - s + iv\right)}{\frac{\kappa}{2} + \varepsilon - s + iv} \right| \right) dv \end{aligned}$$

$$\ll_{\varepsilon,h,K} n^{-\varepsilon} \int_{-\infty}^{\infty} (1 + |v|) \exp\{-c_1|v|\} dv \ll_{\varepsilon,h,K} n^{-\varepsilon} \quad (4)$$

Here, we used the estimate

$$\Gamma\left(\frac{1}{\theta}\left(\frac{\kappa}{2} + \varepsilon - s + iv\right)\right) \ll \exp\left\{-\frac{c}{\theta}|v - t|\right\} \ll_{\kappa} \exp\{-c_1|v|\}, \quad c_1 > 0.$$

Estimate (4) proves the lemma. \square

Let \mathbb{P} be the set of all prime numbers, and $\gamma_p = \{s \in \mathbb{C} : |s| = 1\}$ for all $p \in \mathbb{P}$. Define the set

$$\Omega = \prod_{p \in \mathbb{P}} \gamma_p.$$

Then, the torus Ω with product topology and pointwise multiplication is a compact topological Abelian group. Therefore, on $(\Omega, \mathcal{B}(\Omega))$ ($\mathcal{B}(\mathbb{X})$ is the Borel σ -field of the space \mathbb{X}), the probability Haar measure m_H can be defined. Moreover, let

$$\underline{\Omega} = \Omega_1 \times \cdots \times \Omega_r,$$

where $\Omega_j = \Omega$ for all $j = 1, \dots, r$. Once again, $\underline{\Omega}$ is a compact topological Abelian group. Therefore, on $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}))$ the probability Haar measure \underline{m}_H exists. This gives the probability space $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}), \underline{m}_H)$. Denote by m_{jH} the Haar measure on $(\Omega_j, \mathcal{B}(\Omega_j))$, $j = 1, \dots, r$. Then, \underline{m}_H is the product of the measures m_{1H}, \dots, m_{rH} . Now, denote by $\underline{\omega} = (\omega_1, \dots, \omega_r)$ the elements of $\underline{\Omega}$, where $\omega_j \in \Omega_j$, $j = 1, \dots, r$. Let $\omega_j(p)$ be the p th component of an element $\omega_j \in \Omega_j$, $j = 1, \dots, r$, $p \in \mathbb{P}$. Extend elements $\omega_j(p)$ to the set \mathbb{N} by the formula

$$\omega_j(m) = \prod_{\substack{p^l | m \\ p^{l+1} \nmid m}} \omega_j^l(p), \quad m \in \mathbb{N},$$

and define $H(D_K)$ -valued random element

$$\zeta(s, \omega_j, F) = \sum_{m=1}^{\infty} \frac{c(m) \omega_j(m)}{m^s}, \quad j = 1, \dots, r.$$

The later series is uniformly convergent on compact subsets of D_K for almost all ω_j . Moreover, for fixed $\sigma \in \left(\frac{\kappa}{2}, \frac{\kappa+1}{2}\right)$

$$\int_{-T}^T |\zeta(s + it, \omega_j, F)|^2 dt \ll_{\sigma} T \quad (5)$$

for almost all ω_j , $j = 1, \dots, r$ [18]. Define one more series

$$\zeta_n(s, \omega_j, F) = \sum_{m=1}^{\infty} \frac{c(m) \omega_j(m) v_n(m)}{m^s}, \quad j = 1, \dots, r,$$

which also, as $\zeta_n(s, F)$, are absolutely convergent for $\sigma > \frac{\kappa}{2}$. Let

$$\underline{\zeta}(s + i\hbar\tau, \underline{\omega}, F) = (\zeta(s + ih_1\tau, \omega_1, F), \dots, \zeta(s + ih_r\tau, \omega_r, F))$$

and

$$\underline{\zeta}_n(s + i\hbar\tau, \underline{\omega}, F) = (\zeta_n(s + ih_1\tau, \omega_1, F), \dots, \zeta_n(s + ih_r\tau, \omega_r, F)).$$

Then, repeating the proof of Lemma 1 and using estimate (5), we arrive to the following statement.

Lemma 2. For all \underline{h} and almost all $\underline{\omega}$,

$$\lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{T} \int_0^T \rho \left(\underline{\zeta}(s + i\underline{h}\tau, \underline{\omega}, F), \underline{\zeta}_n(s + i\underline{h}\tau, \underline{\omega}, F) \right) d\tau = 0.$$

3. Limit Theorems

On the probability space $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}), \underline{m}_H)$, define $H(D_\kappa)$ -valued random element

$$\underline{\zeta}(s, \underline{\omega}, F) = (\zeta(s, \omega_1, F), \dots, \zeta(s, \omega_r, F))$$

and denote by $P_{\underline{\zeta}, F}$ its distribution, i.e.,

$$P_{\underline{\zeta}, F}(A) = \underline{m}_H \left\{ \underline{\omega} \in \underline{\Omega} : \underline{\zeta}(s, \underline{\omega}, F) \in A \right\}, \quad A \in \mathcal{B}(H^r(D_\kappa)).$$

Theorem 3. Suppose that h_1, \dots, h_r are real algebraic numbers linearly independent over \mathbb{Q} , and

$$P_{T, F}(A) \stackrel{\text{def}}{=} \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \underline{\zeta}(s + i\underline{h}\tau, F) \in A \right\}, \quad A \in \mathcal{B}(H^r(D_\kappa)).$$

Then, $P_{T, F}$ converges weakly to $P_{\underline{\zeta}, F}$ as $T \rightarrow \infty$.

We divide the proof of Theorem 3 into several lemmas.

Lemma 3. Suppose that $\lambda_1, \dots, \lambda_r$ are algebraic numbers such that the system $\log \lambda_1, \dots, \log \lambda_r$ is linearly independent over \mathbb{Q} . Then, for arbitrary algebraic numbers $\beta_0, \beta_1, \dots, \beta_r$ that are not all zeros, the inequality

$$|\beta_0 + \beta_1 \log \lambda_1 + \dots + \beta_r \log \lambda_r| > h^{-c}$$

holds. Here, h denotes the height of the numbers $\beta_0, \beta_1, \dots, \beta_r$, and c is an effective constant depending on $r, \lambda_1, \dots, \lambda_r$ and maximum of degrees of the numbers $\beta_0, \beta_1, \dots, \beta_r$.

The lemma is a Baker result on linear forms of logarithm; see, for example, ref. [19]. For $A \in \mathcal{B}(\underline{\Omega})$, define

$$Q_T(A) = \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \left(\left(p^{-ih_1\tau} : p \in \mathbb{P} \right), \dots, \left(p^{-ih_r\tau} : p \in \mathbb{P} \right) \right) \in A \right\}.$$

Lemma 4. Let $\lambda_1, \dots, \lambda_r$ be the same as in Theorem 3. Then, Q_T converges weakly to the Haar measure \underline{m}_H as $T \rightarrow \infty$.

Proof. We apply the Fourier transform method. Denote by $g_T(\underline{k}_1, \dots, \underline{k}_r)$, $\underline{k}_j = \{k_{pj} : k_{pj} \in \mathbb{Z}, p \in \mathbb{P}\}$, $j = 1, \dots, r$ the Fourier transform of Q_T . By the definition of Q_T , we have

$$g_T(\underline{k}_1, \dots, \underline{k}_r) = \int_{\underline{\Omega}} \prod_{j=1}^r \prod_{p \in \mathbb{P}}^* \omega_j^{k_{pj}}(p) dQ_T$$

$$= \frac{1}{T} \int_0^T \exp \left\{ -i\tau \sum_{j=1}^r \sum_{p \in \mathbb{P}}^* h_j k_{pj} \log p \right\} d\tau, \quad (6)$$

where the star shows that only a finite number of integers k_{pj} are not zero. Obviously,

$$g_T(\underline{0}, \dots, \underline{0}) = 1. \quad (7)$$

Now, suppose that $(k_1, \dots, k_r) \neq (0, \dots, 0)$. Then, there exists a prime number p such that $k_{pj} \neq 0$ for some j . Therefore,

$$\beta_p \stackrel{\text{def}}{=} \sum_{j=1}^r h_j k_{pj} \neq 0$$

because the numbers h_1, \dots, h_r are linearly independent over \mathbb{Q} . Thus, in view of Lemma 3,

$$B_{k_1, \dots, k_r} \stackrel{\text{def}}{=} \sum_{j=1}^k \sum_{p \in \mathbb{P}}^* h_j k_{pj} \log p = \sum_{p \in \mathbb{P}}^* \beta_p \log p \neq 0.$$

This and (6) imply

$$g_T(k_1, \dots, k_r) = \frac{1 - \exp\{-iTB_{k_1, \dots, k_r}\}}{iTB_{k_1, \dots, k_r}}.$$

Therefore, by (7),

$$\lim_{T \rightarrow \infty} g_T(k_1, \dots, k_r) \stackrel{\text{def}}{=} \begin{cases} 1 & \text{if } (k_1, \dots, k_r) = (0, \dots, 0), \\ 0 & \text{if } (k_1, \dots, k_r) \neq (0, \dots, 0), \end{cases}$$

and this proves the lemma. \square

For $A \in \mathcal{B}(H^r(D_\kappa))$, define

$$P_{T,n,F}(A) = \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \zeta_n(s + i\tau, F) \in A \right\}$$

and

$$P_{T,n,\Omega,F}(A) = \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \zeta_n(s + i\tau, \underline{\omega}, F) \in A \right\}.$$

Moreover, let the mapping $u_n : \Omega \rightarrow H^r(D_\kappa)$ be given by

$$u_{n,F}(\underline{\omega}) = \zeta_n(s, \underline{\omega}, F),$$

and $V_{n,F} = \underline{m}_H u_{n,F}^{-1}$, where

$$V_{n,F}(A) = \underline{m}_H(u_{n,F}^{-1}A), \quad A \in \mathcal{B}(H^r(D_\kappa)).$$

Since the series for $\zeta_n(s, \omega_j, F)$ are absolutely convergent for $\sigma > \frac{\kappa}{2}$, the mapping $u_{n,F}$ is continuous. Moreover, by the definitions of Q_T and $P_{T,n,F}$, we have $P_{T,n,F} = Q_T u_{n,F}^{-1}$. This equality, continuity of $u_{n,F}$, Lemma 4, the well-known properties of weak convergence, and the invariance of the Haar measure \underline{m}_H lead to the following lemma.

Lemma 5. Let h_1, \dots, h_r be the same as Theorem 3. Then, $P_{T,n,F}$ and $P_{T,n,\Omega,F}$ both converge weakly to the measure $V_{n,F}$ as $T \rightarrow \infty$.

Additionally to $P_{T,n,F}$, define

$$P_{T,\Omega,F}(A) = \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \zeta(s + i\tau, \underline{\omega}, F) \in A \right\}, \quad A \in \mathcal{B}(H^r(D_\kappa)).$$

Lemma 6. Let h_1, \dots, h_r be the same as Theorem 3. Then, on $(H^r(D_\kappa), \mathcal{B}(H^r(D_\kappa)))$, there exists a probability measure P_F such that $P_{T,n,F}$ and $P_{T,\Omega,F}$ both converge weakly to P_F as $T \rightarrow \infty$.

Proof. Since the series for $\zeta_n(s, F)$ is absolutely convergent, by a standard way it follows—see, for example [14,18]—that the sequence $\{V_{n,F} : n \in \mathbb{N}\}$ is tight, i.e., for every $\varepsilon > 0$, there exists a compact set $K \subset H^r(D_\kappa)$ such that

$$V_{n,F}(K) > 1 - \varepsilon$$

for all $n \in \mathbb{N}$. Hence, by the Prokhorov theorem, see [20], the sequence $\{V_{n,F}\}$ is relatively compact, i.e., each of its subsequences contains a subsequence $\{V_{n_k,F}\}$ such that $V_{n_k,F}$ converges weakly to a certain probability measure P_F on $(H^r(D_\kappa), \mathcal{B}(H^r(D_\kappa)))$ as $k \rightarrow \infty$.

Let ζ_T be a random variable defined on a certain probability space with measure ν and uniformly distributed on $[0, T]$. Define the $H^r(D_\kappa)$ -valued random element

$$\underline{X}_{T,n,F} = \underline{X}_{T,n,F}(s) = \underline{\zeta}_n(s + i\hbar\zeta_T, F)$$

and denote by $\underline{X}_{n,F} = \underline{X}_{n,F}(s)$ the $H^r(D_\kappa)$ -valued random element having the distribution $V_{n,F}$. Then, by Lemma 5, we have

$$\underline{X}_{T,n,F} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} \underline{X}_{n,F}, \quad (8)$$

where $\xrightarrow[T \rightarrow \infty]{\mathcal{D}}$ means the convergence in distribution. Moreover, since $V_{n_k,F}$ converges weakly to P_F , the relation

$$\underline{X}_{n_k,F} \xrightarrow[k \rightarrow \infty]{\mathcal{D}} P_F \quad (9)$$

is true. Let

$$\underline{X}_{T,F} = \underline{X}_{T,F}(s) = \underline{\zeta}(s + i\hbar\zeta_T, F).$$

Then, using Lemma 1, we find that for every $\varepsilon > 0$,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \nu \left\{ \rho(\underline{X}_{T,F}, \underline{X}_{T,n,F}) \geq \varepsilon \right\} \\ & \leq \lim_{n \rightarrow \infty} \limsup_{T \rightarrow \infty} \frac{1}{\varepsilon T} \int_0^T \rho(\underline{\zeta}(s + i\hbar\tau, F), \underline{\zeta}_n(s + i\hbar\tau, F)) d\tau = 0. \end{aligned}$$

The later equality together with (8) and (9), and Theorem 4.2 of [20] lead to the relation

$$\underline{X}_{T,F} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P_F. \quad (10)$$

This proves that $P_{T,F}$ converges weakly to P_F as $T \rightarrow \infty$.

The relation (10) shows that the limit measure P_F is independent of the subsequence $\{n_k\}$. Therefore, we have

$$\underline{X}_{n,F} \xrightarrow[n \rightarrow \infty]{\mathcal{D}} P_F. \quad (11)$$

Define the $H^r(D_\kappa)$ -valued random elements

$$\underline{X}_{T,n,\Omega,F} = \underline{X}_{T,n,\Omega,F}(s) = \underline{\zeta}_n(s + i\hbar\zeta_T, \omega, F)$$

an

$$\underline{X}_{T,\Omega,F} = \underline{X}_{T,\Omega,F}(s) = \underline{\zeta}(s + i\hbar\zeta_T, \omega, F).$$

Then, repeating the above arguments using Lemmas 2 and 5, and relation (11), we obtain that

$$\underline{X}_{T,n,F} \xrightarrow[T \rightarrow \infty]{\mathcal{D}} P_F,$$

and this is equivalent to weak convergence of $P_{T,\Omega,F}$ to P_F as $T \rightarrow \infty$. The lemma is proved. \square

To prove Theorem 3, it remains to show that $P_F = P_{\zeta,F}$. For this, we will apply some elements of the ergodic theory. For brevity, let

$$\underline{h}_\tau = \left(\left(p^{-ih_1\tau} : p \in \mathbb{P} \right), \dots, \left(p^{-ih_r\tau} : p \in \mathbb{P} \right) \right), \quad \tau \in \mathbb{R}.$$

Define the transformation of $\underline{\Omega}$

$$\varphi_\tau(\underline{\omega}) = \underline{h}_\tau \underline{\omega}, \quad \underline{\omega} \in \underline{\Omega}.$$

Since the Haar measure \underline{m}_H is invariant, the transformation φ_τ is measure-preserving and $\{\varphi_\tau : \tau \in \mathbb{R}\}$ is a one-parameter group. A set $A \in \mathcal{B}(\underline{\Omega})$ is called invariant with respect to the group $\{\varphi_\tau\}$ if the sets A and $\varphi_\tau(A)$, $\tau \in \mathbb{R}$, differ one from another at most by a set of \underline{m}_H -measure zero.

Lemma 7. Let h_1, \dots, h_r be the same as Theorem 3. Then, the group $\{\varphi_\tau\}$ is ergodic, i.e., the σ -field of invariant sets consists of sets having \underline{m}_H -measure 1 or 0.

Proof. The characters χ of the group Ω are of the form

$$\chi(\underline{\omega}) = \prod_{j=1}^r \prod_{p \in \mathbb{P}}^* \omega_j^{k_{pj}}(p).$$

This fact already was used in the proof of Lemma 4. Let A be an arbitrary invariant set, I_A its indicator function, and χ be a nontrivial character. Preserving the notation of the proof of Lemma 4, we have $(k_1, \dots, k_r) \neq (0, \dots, 0)$ and $B_{k_1, \dots, k_r} \neq 0$. Therefore, there exists $\tau_0 \in \mathbb{R}$ such that

$$\chi(\underline{h}_{\tau_0}) = \exp\{-i\tau_0 B_{k_1, \dots, k_r}\} \neq 1. \quad (12)$$

Moreover, in view of the invariance of A , we have

$$I_A(\underline{h}_{\tau_0} \underline{\omega}) = I_A(\underline{\omega}) \quad (13)$$

for almost all $\underline{\omega} \in \underline{\Omega}$. Denote by \hat{I}_A the Fourier transform of I_A . Then, by (13),

$$\hat{I}_A(\chi) = \chi(\underline{h}_{\tau_0}) \int_{\underline{\Omega}} I_A(\underline{h}_{\tau_0} \underline{\omega}) \chi(\underline{\omega}) d\underline{m}_H = \chi(\underline{h}_{\tau_0}) \hat{I}_A(\chi).$$

This and (12) show that

$$\hat{I}_A(\chi) = 0. \quad (14)$$

Now, let χ_0 denote the trivial character of $\underline{\Omega}$, and suppose that $\hat{I}_A(\chi_0) = \alpha$. Then, in view of (14), we find that

$$\hat{I}_A(\chi) = \alpha \int_{\underline{\Omega}} \chi(\underline{\omega}) d\underline{m}_H = \hat{\alpha}(\chi).$$

Hence, $I_A(\underline{\omega}) = \alpha$ for almost all $\underline{\omega} \in \underline{\Omega}$. Since I_A is the indicator function, $I_A(\underline{\omega}) = 1$ or $I_A(\underline{\omega}) = 0$ for almost all $\underline{\omega}$. Thus, $\underline{m}_H(A) = 1$ or $\underline{m}_H(A) = 0$, and the lemma is proved. \square

Proof of Theorem 3. We have mentioned that it suffices to show that $P_F = P_{\zeta,F}$. By Lemma 6 and the equivalent of weak convergence in terms of continuity sets, we have

$$\lim_{T \rightarrow \infty} P_{T, \underline{\Omega}, F}(A) = P_F(A) \quad (15)$$

for a continuity set A of the measure P_F , i.e., $P_F(\partial A) = 0$, where ∂A is the boundary of A . On the probability space $(\underline{\Omega}, \mathcal{B}(\underline{\Omega}), \underline{m}_H)$, define the random variable

$$\tilde{\zeta}(\underline{\omega}) = \begin{cases} 1 & \text{if } \underline{\zeta}(s, \underline{\omega}, F) \in A, \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 7 implies the ergodicity of the random process $\tilde{\zeta}(\varphi_\tau(\underline{\omega}))$. Therefore, by the classical Birkhoff–Khinchine ergodic theorem, see, for example [21],

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T \tilde{\zeta}(\varphi_\tau(\underline{\omega})) d\tau = \mathbb{E}\tilde{\zeta} = P_{\tilde{\zeta}, F}(A), \quad (16)$$

where $\mathbb{E}\tilde{\zeta}$ is the expectation of $\tilde{\zeta}$.

However, by the definitions of φ_τ and $\tilde{\zeta}$,

$$\frac{1}{T} \int_0^T \tilde{\zeta}(\varphi_\tau(\underline{\omega})) d\tau = \frac{1}{T} \text{meas} \left\{ \tau \in [0, T] : \underline{\zeta}(s + i\hbar\tau, \underline{\omega}, F) \in A \right\} = P_{T, \underline{\Omega}, F}(A).$$

This and (16) show that

$$\lim_{T \rightarrow \infty} P_{T, \underline{\Omega}, F}(A) = P_{\tilde{\zeta}, F}(A).$$

Therefore, by (15), we obtain that $P_F(A) = P_{\tilde{\zeta}, F}(A)$ for all continuity sets A of $P_F(A)$. Hence, $P_F = P_{\tilde{\zeta}, F}$, and the theorem is proved. \square

4. Proof of Theorem 2

Recall that the support of the measure $P_{\tilde{\zeta}, F}$ is a minimal closed set $S_F \subset H^r(D_\kappa)$ such that $P_{\tilde{\zeta}, F}(S_F) = 1$.

Lemma 8. *The support of the measure $P_{\tilde{\zeta}, F}$ is the set $(\{g \in H(D_\kappa) : g(s) \neq 0 \text{ or } g(s) \equiv 0\})^r$.*

Proof. Since the space $H^r(D_\kappa)$ is separable, we have [20],

$$\mathcal{B}(H^r(D_\kappa)) = \underbrace{(\mathcal{B}(H(D_\kappa)) \times \cdots \times \mathcal{B}(H(D_\kappa)))}_r.$$

Therefore, it suffices to consider the measure $P_{\tilde{\zeta}, F}$ on the rectangular sets

$$A = A_1 \times \cdots \times A_r, \quad A_1, \dots, A_r \in H(D_\kappa).$$

Let $\zeta(s, \omega, F)$ be the $H(D_\kappa)$ -valued random element defined on the probability space $(\Omega, \mathcal{B}(\Omega), m_H)$, where m_H is the Haar measure. Then, it is known [10] that the support of the distribution of $\zeta(s, \omega, F)$ is the set $\{g \in H(D_\kappa) : g(s) \neq 0 \text{ or } g(s) \equiv 0\}$. Thus, the same set is the support of the distributions of $\zeta(s, \omega_j, F)$, $j = 1, \dots, r$. Since the measure \underline{m}_H is the product of the measures m_{jH} , $j = 1, \dots, r$, we have

$$\underline{m}_H \left\{ \underline{\omega} \in \underline{\Omega} : \underline{\zeta}(s, \underline{\omega}, F) \in A \right\} = \prod_{j=1}^r m_{jH} \left\{ \omega_j \in \Omega_j : \zeta(s, \omega_j, F) \in A_j \right\}.$$

This equality, the minimality of the support, and the support of the distributions of $\zeta(s, \omega_j, F)$ prove the lemma. \square

Proof of Theorem 2. By the Mergelyan theorem on the approximation of analytic functions by polynomials [22], there exist polynomials $p_1(s), \dots, p_r(s)$ such that

$$\sup_{1 \leq j \leq r} \sup_{s \in K_j} |f_j(s) - e^{p_j(s)}| < \frac{\varepsilon}{2}. \quad (17)$$

Define the set

$$G_\varepsilon = \left\{ (g_1, \dots, g_r) \in H^r(D_\kappa) : \sup_{1 \leq j \leq r} \sup_{s \in K_j} |g_j(s) - e^{p_j(s)}| < \frac{\varepsilon}{2} \right\}.$$

In view of Lemma 8, the set G_ε is an open neighborhood of an element $(e^{p_1(s)}, \dots, e^{p_r(s)})$ in support of the measure $P_{\zeta, F}$. Hence,

$$P_{\zeta, F}(G_\varepsilon) > 0. \quad (18)$$

This, Theorem 3 and the equivalent of weak convergence in terms of open sets, and the definitions of $P_{T, F}$ and G_ε prove the theorem with “lim inf”. Define one more set

$$\hat{G}_\varepsilon = \left\{ (g_1, \dots, g_r) \in H^r(D_\kappa) : \sup_{1 \leq j \leq r} \sup_{s \in K_j} |g_j(s) - f_j(s)| < \varepsilon \right\},$$

There $\partial \hat{G}_{\varepsilon_1} \cap \partial \hat{G}_{\varepsilon_2} = \emptyset$ for $\varepsilon_1 \neq \varepsilon_2$. This shows that $P_{\zeta, F}(\partial \hat{G}_\varepsilon) = 0$ for all but, for those countable, many $\varepsilon > 0$. Moreover, (17) and (18) imply that $P_{\zeta, F}(\hat{G}_\varepsilon) > 0$. This, Theorem 3 and the equivalent of weak convergence of probability measures in terms of continuity sets, and the definitions of $P_{T, F}$ and \hat{G}_ε prove the theorem with “lim”. \square

Funding: This research received no external funding.

Conflicts of Interest: The author declares no conflict of interest.

References

- Voronin, S.M. Theorem on the “universality” of the Riemann zeta-function. *Math. USSR Izv.* **1975**, *9*, 443–453. [\[CrossRef\]](#)
- Laurinćikas, A.; Matsumoto, K. The universality of zeta-functions attached to certain cusp forms. *Acta Arith.* **2001**, *98*, 345–359. [\[CrossRef\]](#)
- Laurinćikas, A.; Matsumoto, K.; Steuding, J. Discrete universality of L -functions for new forms. *Math. Notes* **2005**, *78*, 551–558. [\[CrossRef\]](#)
- Laurinćikas, A.; Matsumoto, K.; Steuding, J. Discrete universality of L -functions for new forms. II. *Lith. Math. J.* **2016**, *56*, 207–218. [\[CrossRef\]](#)
- Laurinćikas, A.; Matsumoto, K.; Steuding, J. Universality of some functions related to zeta-functions of certain cusp forms. *Osaka J. Math.* **2013**, *50*, 1021–1037.
- Laurinćikas, A. On the functional independence of zeta-functions of certain cusp forms. *Math. Notes* **2020**, *107*, 609–617. [\[CrossRef\]](#)
- Laurinćikas, A.; Matsumoto, K.; Steuding, J. The universality of L -functions associated with new forms. *Izv. Math.* **2003**, *67*, 77–90. [\[CrossRef\]](#)
- Vaiginytė, A. Extension of the Laurinćikas-Matsumoto theorem. *Chebyshevskii Sb.* **2019**, *20*, 82–93.
- Laurinćikas, A.; Šiaučiūnas, D.; Vaiginytė, A. Extension of the discrete universality theorem for zeta-functions of certain cusp forms. *Nonlinear Anal. Model. Control* **2018**, *23*, 961–973. [\[CrossRef\]](#)
- Balčiūnas, A.; Franckevič, V.; Garbaliuskienė, V.; Macaitienė, R.; Rimkevičienė, A. Universality of zeta-functions of cusp forms and non-trivial zeros of the Riemann zeta-function. *Math. Model. Anal.* **2021**, *26*, 82–93. [\[CrossRef\]](#)
- Voronin, S.M. On the functional independence of Dirichlet L -functions. *Acta Arith.* **1975**, *27*, 493–503. (In Russian)
- Laurinćikas, A.; Matsumoto, K. The joint universality of twisted automorphic L -functions. *J. Math. Soc. Jpn.* **2004**, *56*, 923–939. [\[CrossRef\]](#)
- Laurinćikas, A.; Šiaučiūnas, D.; Vaiginytė, A. On joint approximation of analytic functions by nonlinear shifts of zeta-functions of certain cusp forms. *Nonlin. Anal. Model. Control* **2020**, *25*, 108–125. [\[CrossRef\]](#)
- Laurinćikas, A. Joint universality of zeta-functions with periodic coefficients. *Izv. Math.* **2010**, *74*, 515–539. [\[CrossRef\]](#)
- Laurinćikas, A. Joint universality of zeta functions with periodic coefficients. II. *Sib. Math. J.* **2020**, *61*, 1064–1076. [\[CrossRef\]](#)
- Laurinćikas, A. Joint discrete universality for periodic zeta-functions. III. *Quaest. Math.* **2020**. [\[CrossRef\]](#)

-
17. Kaczorowski, J.; Laurinćikas, A.; Steuding, J. On the value distribution of shifts of universal Dirichlet series. *Monatsh. Math.* **2006**, *147*, 309–317. [[CrossRef](#)]
 18. Kačėnas, A.; Laurinćikas, A. On Dirichlet series related to certain cusp forms. *Lith. Math. J.* **1998**, *38*, 64–76. [[CrossRef](#)]
 19. Baker, A. The theory of linear forms in logarithms. In *Transcendence Theory: Advances and Applications. Proceedings of a Conference Held in Cambridge in 1976*; Baker, A., Masser, D.W., Eds.; Academic Press: Boston, MA, USA, 1977; pp. 1–27.
 20. Billingsley, P. *Convergence of Probability Measures*; Willey: New York, NY, USA, 1968.
 21. Cramér, H.; Leadbetter, M.R. *Stationary and Related Stochastic Processes*; Willey: New York, NY, USA, 1967.
 22. Mergelyan, S.N. Uniform approximations to functions of a complex variable. *Am. Math. Soc. Transl. Ser. 2* **1962**, *3*, 294–391.