

Supplementary Materials

Endothelial Progenitor Cells: Relevant Players in the Vasculopathy and Lung Fibrosis Associated with the Presence of Interstitial Lung Disease in Systemic Sclerosis Patients

Verónica Pulito-Cueto^{1*}, Sara Remuzgo-Martínez^{1*}, Fernanda Genre^{1*}, Belén Atienza-Mateo^{1,2,3}, Víctor M. Mora-Cuesta^{1,4}, David Iturbe-Fernández^{1,4}, Leticia Lera-Gómez¹, Raquel Pérez-Fernández¹, Diana Prieto-Peña^{1,3}, Virginia Portilla^{1,3}, Ricardo Blanco^{1,3}, Alfonso Corrales^{1,3}, Oreste Gualillo⁵, José M. Cifrián^{1,4,6}, Raquel López-Mejías^{1,**} and Miguel A. González-Gay^{1,3,6,7**}

Table S1. Differences of EPC frequency between SSc-ILD⁺ patients and the three comparative groups (healthy controls, SSc-ILD⁻ and IPF patients).

	EPC frequency (Mean ± SD)	p
SSc-ILD ⁺ patients vs healthy controls	0.033 ± 0.012 vs 0.013 ± 0.010	< 0.001
SSc-ILD ⁺ patients vs SSc-ILD ⁻ patients	0.033 ± 0.012 vs 0.021 ± 0.017	0.012
SSc-ILD ⁺ patients vs IPF patients	0.033 ± 0.012 vs 0.087 ± 0.041	< 0.001

EPC: endothelial progenitor cells; SSc: systemic sclerosis; ILD: interstitial lung disease; IPF: idiopathic pulmonary fibrosis; SD: standard deviation. Significant results are highlighted in **bold**.