

Supplementary Material


Immobilization of Laccase on Hybrid Super-Structured Nanomaterials for the Decolorization of Phenolic dyes

Michaela Patila ^{1,*}, Panagiotis E. Athanasiou ¹, Lampros Kortessis ¹, Georgia Potsi ², Antonios Kouloumpis ², Dimitrios Gournis ² and Haralambos Stamatis ^{1,*}


¹ Biotechnology Laboratory, Department of Biological Applications and Technology, University of Ioannina, 45110 Ioannina, Greece; p.athanasiou@uoi.gr (P.E.A.); bbroslas187@gmail.com (L.K.)

² Department of Materials Science and Engineering, University of Ioannina, 45110 Ioannina, Greece; gepotsi@gmail.com (G.P.); antoniokoul@gmail.com (A.K.); dgourni@uoi.gr (D.G.)


* Correspondence: pstm10345@uoi.gr (M.P.); hstamati@uoi.gr (H.S.)




(a)



(b)




(c)




(d)

Figure S1. FTIR spectra of (a) free TvL, (b) Clay-ADA nanomaterial and immobilized TvL on Clay-ADA, (c) Clay-CNTs nanomaterial and TvL immobilized on Clay-CNTs, (d) Clay-GO nanomaterial and immobilized TvL on Clay-GO.




(a)




(b)


Figure S2. Decolorization of different dyes (75 µM) by (a) non-covalently and (b) covalently immobilized Tvl on hybrid nanomaterials at 48 h. The reaction was conducted in acetate buffer (0.1 M, pH 4.58), at 30 °C.




(a)




(b)



(c)



(d)



(e)

Figure S3. Effect of concentration of (a) bromophenol blue, (b) bromothymol blue, (c) Coomassie brilliant blue, (d) methyl orange, (e) phenol red on the decolorization ability of immobilized TvL on hybrid nanomaterials. The reaction was conducted in acetate buffer (0.1 M, pH 4.58), at 30 °C, for 48 h.